



National Toxicology Program
U.S. Department of Health and Human Services

**Peer-Review Draft:
Report on Carcinogens Monograph on
Epstein-Barr Virus**

November 2, 2015

Office of the Report on Carcinogens
Division of the National Toxicology Program
National Institute of Environmental Health Sciences
U.S. Department of Health and Human Services

This information is distributed solely for the purpose of pre-dissemination peer review under applicable information quality guidelines. It has not been formally distributed by the National Toxicology Program. It does not represent and should not be construed to represent any NTP determination or policy.

This Page Intentionally Left Blank

Foreword

The National Toxicology Program (NTP) is an interagency program within the Public Health Service (PHS) of the Department of Health and Human Services (HHS) and is headquartered at the National Institute of Environmental Health Sciences of the National Institutes of Health (NIEHS/NIH). Three agencies contribute resources to the program: NIEHS/NIH, the National Institute for Occupational Safety and Health of the Centers for Disease Control and Prevention (NIOSH/CDC), and the National Center for Toxicological Research of the Food and Drug Administration (NCTR/FDA). Established in 1978, the NTP is charged with coordinating toxicological testing activities, strengthening the science base in toxicology, developing and validating improved testing methods, and providing information about potentially toxic substances to health regulatory and research agencies, scientific and medical communities, and the public.

The Report on Carcinogens (RoC) is prepared in response to Section 301 of the Public Health Service Act as amended. The RoC contains a list of identified substances (i) that either are *known to be human carcinogens* or are *reasonably anticipated to be human carcinogens* and (ii) to which a significant number of persons residing in the United States are exposed. The Secretary, Department of HHS, has delegated responsibility for preparation of the RoC to the NTP, which prepares the report with assistance from other Federal health and regulatory agencies and nongovernmental institutions. The most recent RoC, the 13th Edition (2014), is available at <http://ntp.niehs.nih.gov/go/roc>.

Nominations for (1) listing a new substance, (2) reclassifying the listing status for a substance already listed, or (3) removing a substance already listed in the RoC are evaluated in a scientific review process (<http://ntp.niehs.nih.gov/go/rocprocess>) with multiple opportunities for scientific and public input and using established listing criteria (<http://ntp.niehs.nih.gov/go/15209>). A list of candidate substances under consideration for listing in (or delisting from) the RoC can be obtained by accessing <http://ntp.niehs.nih.gov/go/37893>.

Overview and Introduction

This collection of monographs on selected viruses provide cancer hazard evaluations for the following human viruses: Epstein-Barr virus, Kaposi sarcoma herpesvirus, human immunodeficiency virus-1, human T-cell lymphotropic virus-1, and Merkel cell polyomavirus for potential listing in the Report on Carcinogens (RoC). Currently, there are three human oncogenic viruses listed in the RoC: human papillomaviruses: some genital-mucosal types (HPV), hepatitis B virus (HBV), and hepatitis C virus (HCV). The five viruses covered in these monographs were selected for review for the RoC based on a large database of information on these agents, including authoritative reviews, and public health concerns for disease mortality and morbidity both in the United States and worldwide because of significant numbers of infected people.

This section provides background information on the preparation of the monographs as well as a discussion of overarching issues related to evaluating the evidence for cancer from human epidemiology studies and evaluating the causation by viruses.

Background

The RoC draft monograph for each virus consists of the following components: (Part 1) the cancer evaluation component that reviews the relevant scientific information and assesses its quality, applies the RoC listing criteria to the scientific information, and recommends an RoC listing status, and (Part 2) the draft substance profile containing the NTP's preliminary listing recommendation, a summary of the scientific evidence considered key to reaching that recommendation, and information on properties, exposure, and Federal regulations and guidelines. Information reviewed in the monographs, with the exception of information on properties and exposure, comes from publicly available and peer-reviewed sources. All sections of the monographs underwent scientific and quality assurance review by independent reviewers.

The cancer evaluation component provides the following information relevant to a RoC listing recommendation: Properties and Detection (Section 1), Exposure (Section 2), Human Cancer Hazard Evaluation for specific cancer endpoints (Section 3), Mechanistic and Other Relevant Data (Section 4), and Preliminary Listing Recommendation (Section 5). Because these viruses are primarily species-specific for humans and similar to the approach used by IARC, we are including information on studies in experimental animals in the Mechanistic and Other Relevant Data section of the monographs. Also, specific details about the strains of the viruses are given only if needed to provide context, such as in the viral Properties and Detection section. The monographs relied on the information and data provided in previous IARC monographs on these five viruses in addition to newer key studies or reviews published since the IARC monographs; it is an independent assessment of available data through August 17, 2015. Literature search strategies to obtain information relevant to the cancer evaluation are in Appendix A of each virus monograph; search terms were developed in collaboration with a reference librarian.

Issues related to evaluating the evidence from human epidemiological studies

The available studies of specific cancer endpoints in the human virus studies present several challenges with respect to the evaluation of methodological strengths and limitations of the body of evidence. Large prospective cohort studies, particularly those that follow individuals for

whom infection status is documented prior to follow-up or cancer diagnosis, have several potential methodological strengths, including evidence that infection precedes cancer diagnosis, adequate statistical power and, in some studies, the ability to analyze dose-response relationships. However, there is the potential for misclassification of exposure in studies that measure the virus once, but with a long follow-up period as they may miss new infections. For most cancer endpoints, only cross-sectional or retrospective cohort studies or hospital or clinic-based case-control studies are available, which lack direct evidence of temporality and may lack power or adequate data on, e.g., viral load. However, molecular evidence from human studies and mechanistic data can be used in the evaluation of temporality, distinguishing latent infections caused by the tumor virus and causality. For some (typically rare) outcomes (e.g., cutaneous T-cell lymphoma and human T-cell lymphotropic virus type 1, or lymphoepithelial carcinoma of the salivary gland and Epstein-Barr virus), only case-comparison studies, in which selection of comparison groups may be biased, unmatched, or inadequately described, or case series, are available.

In addition, for several rare endpoints, e.g., adult T-cell leukemia/lymphoma and human T-cell lymphotropic virus type 1, or primary effusion lymphoma and Kaposi sarcoma herpesvirus, the presence of the virus in the tumor cells is used as a diagnostic criterion to define the cancer, and thus evidence of causality relies on cases defined by this criterion and molecular evidence from human studies rather than on epidemiological population-based studies of the association of the virus with a level of cancer risk.

For several viruses, e.g., Epstein-Barr virus, the population prevalence may exceed 90%, so that cohort and case-control studies must rely on the evaluation of cancer risk using measures such as Epstein-Barr virus titer or antibody levels rather than exposed and non-exposed categories of study participants, allowing for the possibility that past or current viral level could be misclassified. In addition, for a number of these viruses, e.g., Kaposi sarcoma herpesvirus, the presence of the virus may be necessary but not sufficient to increase the risk for a specific cancer endpoint and more than one virus may be associated with risk. Thus, methodologically adequate studies should include measurement of such cofactors and consider potentially confounding factors; however, relatively few studies have measured a panel of other viruses or taken into account other cofactors. In addition, while studies comparing cancer risk in treated vs. untreated populations may provide indirect evidence of the role of human immunodeficiency virus-1, these studies, in particular calendar-period analyses, may not adequately account for changes in risk attributable to improved survival rates or changes in other risk factors.

Issues related to evaluating causality of viruses

Approximately 12% of all human cancers have been attributed to viral infections; however, viruses are rarely fully oncogenic themselves and only a small percentage of infected individuals develop cancer, often decades after the initial infection (Mesri *et al.* 2014). Therefore, oncogenic viruses are generally considered necessary but not sufficient to cause cancer. Additional cofactors, such as infective organisms, chemicals, or environmental agents in conjunction with risk modifiers such as immune dysfunction or chronic inflammation can contribute to malignant transformation. Severe immunosuppression, as seen with congenital immunodeficiency syndromes, chronic human immunodeficiency virus type 1 infection, or as a result of tissue anti-rejection medication, can severely compromise the immune surveillance capabilities of the patient. In addition, some cofactors produced by other organisms or agents have been shown to

activate the oncogenic potential of some of these viruses. There are also other challenges that are somewhat unique to the evaluation of the epidemiological studies (discussed below) and thus molecular evidence is often considered in the evaluation of causality.

In light of these issues, IARC monographs and several other publications have discussed paths to evaluate causality, which are discussed below and incorporated into the NTP approach for evaluating causality of the viruses. What is important for public health in determination of causation of a health effect, such as risk for cancer, is whether that health effect is eliminated or mitigated by removal of the substance.

There have been a number of attempts to develop criteria that address causal associations. However, all of them have limitations, especially when applied to infectious agents (Moore and Chang 2010). The following sections identify factors to consider for evaluating causality, some of the limitations associated with strict application of the criteria in the context of virally induced cancers, some alternative approaches, and the NTP's approach for evaluating the role of select viral agents in human cancer.

Hill's characteristics for evaluation of epidemiological studies

Hill proposed nine characteristics to consider when evaluating causality, primarily for epidemiological studies, although they have been expanded for evaluating mechanistic and other types of data (Table 1). Several considerations—strength of the association, consistency across studies, evidence of an exposure-response gradient, and temporality of exposure (Hill 1965)—are used to help guide the RoC evaluations of the human epidemiological data (see RoC Handbook, NTP 2015). However, it should be noted that these are not criteria; with the exception of temporality, each and every element is not required in order to demonstrate causality (Rothman and Greenland 2005). Hill (1965) avoided discussing the meaning of “causation” noting that the “cause” of an illness could be immediate and direct or remote and indirect. The primary question addressed by Hill was “whether the frequency of the undesirable event B will be influenced by a change in the environmental feature A.”

Table 1. Hill's epidemiological characteristics for causality

Criterion	Description
1. Strength of association	A strong association between a virus and a cancer is most consistent with causality unless confounded by some other exposure. However, a weak association does not give evidence against causality.
2. Consistency	Consistent findings observed by different persons, in different places, circumstances and times.
3. Specificity	A viral exposure is limited to specific types of cancer (considered a weak factor because there are well-established examples in which multiple types of disease are caused by one type of exposure). However, the more specific the association, the higher the probability of a causal relationship.
4. Temporality	Exposure to the virus must occur prior to the onset of the cancer, in contrast to a “passenger infection.”
5. Biologic gradient	The virus is more likely to be found at the tumor site than at non-tumor sites.
6. Plausibility	Should be applied with caution because it is limited by current medical knowledge (e.g., an implausible mechanism may gain acceptance with increased understanding of the underlying biology).

7. Coherence	A virus-cancer association should not seriously conflict with known facts on the cancer's natural history and biology.
8. Experiment	Changing either exposure or continued infection in a randomized clinical trial should change the measure of clinical outcome (e.g., vaccination programs for HPV and HBV).
9. Analogy	Are related viruses clearly established to cause cancers in animals or humans?

Source: Moore and Chang 2010.

Consideration of mechanistic data from studies in humans

In their evaluation of the evidence for Epstein-Barr virus, the IARC working group noted that the large majority of people are latently infected with Epstein-Barr virus, thus epidemiological studies may be limited in determining whether the presence of Epstein-Barr virus in tumor tissue is a cause of the cancer or an effect of the tumor. Thus, in addition to the Hill characteristics, IARC (1997) also considered the following in their evaluation of Epstein-Barr virus, which are applicable to other viruses:

- the proportion of Epstein-Barr virus-positive cases in a given tumor entity,
- the proportion of tumor cells that carry the virus,
- the monoclonality of Epstein-Barr virus in the tumor, and
- the expression of Epstein-Barr virus proteins.

zur Hausen (2001, 1994) also noted the difficulty of applying stringent criteria to identify human tumor viruses and proposed the following:

- the regular presence and persistence of the respective viral DNA in tumor biopsies and cell lines derived from the same tumor type,
- the demonstration of growth-promoting activity of specific viral genes or of virus-modified host cell genes in tissue culture systems or in suitable animal systems,
- the demonstration that the malignant phenotype depends on the continuous expression of viral oncogenes or on the modification of host cell genes containing viral sequences,
- epidemiological evidence that the respective virus infection represents a major risk factor for cancer development.

It is difficult to prove that a virus causes cancer, and such determinations almost always generate considerable controversy and debate (Moore and Chang 2010). Viral cancers employ various mechanisms that involve both direct and indirect modes of interaction (Table 2) (zur Hausen and de Villiers 2014). Understanding and managing viral-induced cancers in humans has been hampered by a lack of suitable animal models, the disparate nature of tumor types, a long latency period between primary infection and cancer development, the different types of oncogenic viruses, and the complex nature of the virus-host cell interactions leading to cancer (Mesri *et al.* 2014, zur Hausen and de Villiers 2014).

Table 2. Direct and indirect modes of interaction of viral infections

Type	Description
Direct carcinogenesis	<ul style="list-style-type: none"> Continued presence and expression of viral oncogenes usually after viral genome integration into host cell DNA Insertional gene activation or suppression Continued episomal presence of viral nucleic acid and suppression or activation of cellular genes (e.g., by viral microRNA)
Indirect carcinogenesis	<ul style="list-style-type: none"> Induction of immunomodulation, activation of latent tumor virus genomes Induction of oxygen and nitrogen radicals Amplification of latent tumor virus DNA Induction of mutations and/or translocations Prevention of apoptosis

Source: zur Hausen and de Villiers 2014.

Multicausality issues

Although thousands of viruses are known to cause infection, only a few have been shown to cause cancer in humans (Moore and Chang 2010). An agent that is both necessary and sufficient for a disease to occur describes a complete causal effect. However, this is not a practical definition for infectious diseases that emerge from complex interactions of multiple factors and may be caused by more than a single agent. An important consideration regarding multicausality is that most of the identified causes are neither necessary nor sufficient in the absence of other factors to produce the disease; however, a cause does not have to be either necessary or sufficient for its removal to result in disease prevention (Rothman 1976, zur Hausen and de Villiers 2014). Although the known oncogenic viruses belong to different virus families, they share several common traits: (1) they are often necessary but not sufficient for tumor development; (2) viral cancers appear in the context of persistent infections and occur many years to decades after acute infection; and (3) the immune system can play a deleterious or a protective role (Mesri *et al.* 2014).

Application of causality criteria and alternative approaches

Moore and Chang (2010) investigated the difficulties associated with strict application of the Hill characteristics for two of the most recently discovered oncogenic viruses: Kaposi sarcoma herpesvirus and Merkel cell polyomavirus. Kaposi sarcoma herpesvirus was shown to fulfill Hill's characteristics for causality of Kaposi sarcoma; however, the application of the characteristics was problematic in the case of Merkel cell polyomavirus and Merkel cell carcinoma (see the monographs for Kaposi sarcoma herpesvirus and Merkel cell polyomavirus). These two examples illustrate the diversity in the patterns of tumor virus epidemiology. Some of the reasons Hill's characteristics worked for Kaposi sarcoma herpesvirus but not Merkel cell polyomavirus is that all clinical forms of Kaposi sarcoma require Kaposi sarcoma herpesvirus while most studies indicate that all forms of Merkel cell carcinoma do not require Merkel cell polyomavirus infection. Further, Kaposi sarcoma herpesvirus infection is uncommon in most parts of the world but was confirmed to be present in nearly all AIDS-associated Kaposi sarcoma cases, while widespread Merkel cell polyomavirus infection rate implies that it cannot be a specific causal factor for a rare cancer like Merkel cell carcinoma. In the case of Merkel cell polyomavirus, additional considerations, as suggested by IARC (1997) and zur Hausen (2001,

1994), provide molecular evidence of the association between Merkel cell polyomavirus and Merkel cell carcinoma, such as the tumor-causing form of the virus is mutated and monoclonally integrated into the tumor genome and that tumor cells require the presence of viral oncoproteins for cell survival and proliferation.

While causal criteria can be helpful, there are flaws and practical limitations that restrict their use in cancer biology (Moore and Chang 2010). Therefore, a more probabilistic approach may be more useful for determining whether or not certain viruses cause human cancers. For example, instead of trying to determine if virus A causes cancer B, the probabilistic approach examines if cancer B is more probable in the presence of virus A. Although a correlation does not imply causation, it can be argued that correlations that are strong, reproducible, and predictive have a similar value as a causative conclusion. In a similar fashion, zur Hausen and de Villiers (2014) also expressed concern over all attempts to summarize criteria for “causality” of infectious agents in cancer development and proposed replacing “causal factor” with “risk factor” and grading them according to their contribution to an individual’s cancer risk. This will require a greater understanding of the complexity of factors involved and their mechanistic contribution to individual cancers.

NTP’s approach

For each virus, the NTP applied the RoC listing criteria (see text box) to the body of literature to reach the preliminary listing recommendation. The level of evidence conclusion from studies in humans considers the evidence from epidemiological studies as well as clinical and molecular studies of tissues from exposed (i.e., infected) individuals. In evaluating the

RoC Listing Criteria

Known To Be Human Carcinogen:

There is sufficient evidence of carcinogenicity from studies in humans*, which indicates a causal relationship between exposure to the agent, substance, or mixture, and human cancer.

Reasonably Anticipated To Be Human Carcinogen:

There is limited evidence of carcinogenicity from studies in humans*, which indicates that causal interpretation is credible, but that alternative explanations, such as chance, bias, or confounding factors, could not adequately be excluded, OR

there is sufficient evidence of carcinogenicity from studies in experimental animals, which indicates there is an increased incidence of malignant and/or a combination of malignant and benign tumors (1) in multiple species or at multiple tissue sites, or (2) by multiple routes of exposure, or (3) to an unusual degree with regard to incidence, site, or type of tumor, or age at onset, OR

there is less than sufficient evidence of carcinogenicity in humans or laboratory animals; however, the agent, substance, or mixture belongs to a well-defined, structurally related class of substances whose members are listed in a previous Report on Carcinogens as either known to be a human carcinogen or reasonably anticipated to be a human carcinogen, or there is convincing relevant information that the agent acts through mechanisms indicating it would likely cause cancer in humans.

Conclusions regarding carcinogenicity in humans or experimental animals are based on scientific judgment, with consideration given to all relevant information. Relevant information includes, but is not limited to, dose response, route of exposure, chemical structure, metabolism, pharmacokinetics, sensitive sub-populations, genetic effects, or other data relating to mechanism of action or factors that may be unique to a given substance. For example, there may be substances for which there is evidence of carcinogenicity in laboratory animals, but there are compelling data indicating that the agent acts through mechanisms which do not operate in humans and would therefore not reasonably be anticipated to cause cancer in humans.

*This evidence can include traditional cancer epidemiology studies, data from clinical studies, and/or data derived from the study of tissues or cells from humans exposed to the substance in question that can be useful for evaluating whether a relevant cancer mechanism is operating in people.

mechanistic data and determining the preliminary recommendations for its level of evidence conclusion and overall listing recommendation, the NTP considered the principles outlined by Hill, IARC, zur Hausen, and Rothman in its assessment of causality for the five viruses reviewed. However, these factors were not used as a strict checklist to either prove or disprove a causal association but rather as guidance to assess the level of epidemiological or molecular evidence that a virus contributes to a carcinogenic effect.

CONTRIBUTORS

Office of the Report on Carcinogens (ORoC), Division of the National Toxicology Program (NTP)

Conducted technical review and evaluation and proposed the preliminary listing recommendation

Ruth Lunn, DrPH
(Director, ORoC)

Gloria D. Jahnke, DVM, DABT
(project lead)

Office of Scientific Information Management, Office of the Deputy Director, NIEHS

Assisted with literature searches

Stephanie Holmgren, MSLS, MBA
Informaticist

Integrated Laboratory Systems, Inc. (Support provided through NIEHS Contract Number HHSN273201100004C)

Conducted technical review and evaluation

Sanford Garner, PhD
(Principal Investigator)

Andrew Ewens, PhD, DABT

Stanley Atwood, MS, DABT

Jennifer Ratcliffe, PhD, MSc

Jessica Geter, MSLS

Alton Peters, MS

Provided administrative support

Ella Darden, BS

Tracy Saunders, BS

Social Scientific Systems, Inc. (Support provided through NIEHS Contract Number HHSN2732015000041)

Conducted technical review and evaluation

Whitney Arroyave, PhD

Technical Advisors

Jim Goedert, MD
Senior Investigator
Infections and Immunoepidemiology Branch
Division of Cancer Epidemiology & Genetics
National Cancer Institute

Elizabeth 'Betsy' Read-Connole, PhD
Head, Cancer Etiology Section
Cancer Immunology and Hematology Etiology Branch
Division of Cancer Biology
National Cancer Institute

Sam Mbulaiteye, MBChB, M.Phil., M.Med.
Senior Investigator
Infections and Immunoepidemiology Branch
Division of Cancer Epidemiology & Genetics
National Cancer Institute

This Page Intentionally Left Blank

Part 1

Draft Cancer Hazard Evaluation

Properties and Detection

Exposure

Human Cancer Studies

Mechanistic Data and Other Relevant Effects

Preliminary Listing Recommendation

This Page Intentionally Left Blank

Table of Contents

1	Properties and Detection Methods	1
1.1	Biological Properties	1
1.1.1	Family and type.....	1
1.1.2	Viral structure and genome.....	1
1.1.3	Infection and replication	2
1.2	Detection	5
1.3	Summary	6
2	Exposure	7
2.1	Transmission and prevalence	7
2.2	Diseases, prevention, treatment.....	7
2.3	Summary	8
3	Human Cancer Studies.....	9
3.1	Selection of the literature	9
3.2	Cancer evaluation: Burkitt lymphoma	9
3.2.1	Background information	9
3.2.2	Case-control studies	10
3.2.3	Cohort studies	13
3.2.4	Cofactors	14
3.2.5	Integration of the evidence.....	15
3.3	Cancer evaluation: Hodgkin lymphoma.....	15
3.3.1	Background information	15
3.3.2	Case-control studies	16
3.3.3	Cohort studies	19
3.3.4	Integration of the evidence.....	20
3.4	Cancer evaluation: Immunosuppression-related Non-Hodgkin Lymphoma.....	21
3.4.1	Background information	21
3.4.2	Studies and evaluation	21
3.4.3	Integration of the evidence.....	22
3.5	Cancer evaluation: Extranodal NK/T-cell lymphoma, nasal type.....	22
3.6	Cancer evaluation: Nasopharyngeal carcinoma	23
3.6.1	Background information	23
3.6.2	Case-control studies	23
3.6.3	Cohort studies	26
3.6.4	Cofactors	26
3.6.5	Integration of the evidence.....	27
3.7	Cancer evaluation: Gastric cancer	27
3.7.1	Background information on gastric cancer	27
3.7.2	Case-series analyses and pooled analyses.....	27
3.7.3	Case-control studies and case-case analyses.....	28
3.7.4	Nested case-control studies.....	29
3.7.5	Cofactors	30
3.7.6	Integration of the evidence.....	30
3.8	Cancer evaluation: Lymphoepithelial carcinoma of the salivary gland	31
3.8.1	Case-case study.....	31

3.8.2	Cofactors	31
3.8.3	Integration of the evidence.....	31
3.9	Synthesis across cancer endpoints.....	32
4	Mechanisms and Other Relevant Data.....	33
4.1	General characteristics	33
4.2	EBV latent genes and malignant transformation.....	35
4.3	Mode of action and evidence for cancer causation	36
4.3.1	Burkitt lymphoma	36
4.3.2	Hodgkin lymphoma	37
4.3.3	Immunosuppression-related non-Hodgkin lymphoma (HIV+ and post-transplant lymphoproliferative disease).....	38
4.3.4	Extranodal NK/T-cell lymphoma, nasal type	39
4.3.5	Nasopharyngeal carcinoma.....	39
4.3.6	Gastric carcinoma	40
4.3.7	Lymphoepithelial cancer of salivary glands	41
4.4	Synthesis.....	42
5	Preliminary Listing Recommendation	43
	References.....	47
	Glossary	63
	Abbreviations.....	65
	Appendix A: Literature Search Strategy.....	A-1
	General approach.....	A-1
	Search strings for EBV Searches.....	A-2
	Cancer and Epidemiology.....	A-2
	Relevant Cancers	A-3
	Draft Profile	P-1

List of Tables

Table 1-1.	EBV transcription programs in normal B cells and in associated malignancies	2
Table 1-2.	Methods of EBV detection	5
Table 3-1.	Summary of cohort and nested case-control studies of Hodgkin lymphoma and Epstein-Barr Virus.....	20
Table 3-2.	Summary of case-comparison studies of NK/T cell lymphoma, nasal type and Epstein-Barr Virus.....	23
Table 3-3.	Summary of case-control studies of gastric cancer and Epstein-Barr Virus	29
Table 3-4.	Summary of cohort and nested case-control studies of gastric cancer and Epstein-Barr Virus	30
Table 3-5.	Summary of EBV cancer endpoints and strength of the epidemiological evidence	32
Table 4-1.	Characteristics of EBV-associated neoplasms	33
Table 4-2.	Activation of oncogenic pathways by viral genes.....	35
Table 4-3.	Properties of some known signaling pathways altered with EBV-related gastric cancer	41

Table 5-1. Evidence for EBV and Burkitt lymphoma (endemic or sporadic) from human studies	43
Table 5-2. Evidence for EBV and Hodgkin lymphoma from human studies	44
Table 5-3. Evidence for EBV and immunosuppression-related non-Hodgkin lymphoma (NHL) from human studies	44
Table 5-4. Evidence for EBV and NK/T cell leukemia/lymphoma, nasal type from studies in humans.....	44
Table 5-5. Evidence for EBV and nasopharyngeal carcinoma from human studies	45
Table 5-6. Evidence for EBV and gastric cancer from human studies	45
Table 5-7. Evidence for EBV and lymphoepithelial carcinoma of the salivary gland from human studies	45
Table A-1. Major topics searched	A-1

List of Figures

Figure 1-1. Epstein-Barr Virus structure	2
Figure 3-1. Forest plot of serological case-control studies of endemic Burkitt lymphoma and Epstein-Barr virus.....	11
Figure 3-2. Forest plot of serological case-control studies of sporadic Burkitt lymphoma and Epstein Barr Virus	13
Figure 3-3. Forest plot of case-control studies of Hodgkin lymphoma and infectious mononucleosis	19
Figure 3-4. Forest plot of serological case-control studies of Hodgkin lymphoma and Epstein-Barr Virus.....	17
Figure 3-5. Forest plot of serological case-control studies of nasopharyngeal carcinoma and Epstein-Barr virus.....	25
Figure A-1: Literature processing flow.....	A-2

This Page Intentionally Left Blank

1 Properties and Detection Methods

This section reviews the biological properties (Section 1.1) and detection methods (Section 1.2) for Epstein-Barr Virus.

1.1 Biological Properties

The following section reviews the types of EBV, its structure, life cycle and course of infection.

1.1.1 Family and type

Epstein-Barr Virus (EBV) is a human herpesvirus in the gammaherpesvirus subfamily and is the prototype of the *Lymphocryptovirus* genus (IARC 2012). The formal designation of EBV is human herpesvirus 4 (HHV4). EBV was the first human tumor virus to be discovered and was detected in Burkitt lymphoma cells in 1964. The two major types of EBV (EBV-1 and EBV-2) differ in their gene sequences of nuclear antigens (EBNA-2, -3A, -3B). Different viral strains within the two major types are created by DNA polymorphisms at other DNA sites.

1.1.2 Viral structure and genome

The EBV is composed of a nucleocapsid surrounded by an envelope with glycoprotein spikes protruding from its outer surface; the nucleocapsid and envelope are separated by tegument proteins (Figure 1-1) (IARC 2012). Inside the nucleocapsid is a double stranded DNA (dsDNA) genome approximately 172 kb in length, wrapped around a protein core. The genome encodes over 85 genes; the open reading frames (ORFs) are divided into latent and lytic genes. ORF nomenclature is based on the *Bam*HI restriction fragment in which they are found. The genome has a series of 0.5 kb terminal direct repeats at each end as well as internal repeat sequences dividing the genome into long and short unique sequences that have most of the coding capacity. EBV forms multiple episomes within the cell nucleus that are circularized by joining their terminal repeats. Latent EBV infections of clonal origin have the same number of terminal repeats. EBV uses different combinations of latent viral gene expression to progress from initial infection to long-term persistence within the memory B cell pool; this is discussed in the following section. Latent EBV genes have a primary role in EBV-associated oncogenesis and are discussed in Section 4.2.2.




Figure 1-1. Epstein-Barr Virus structure

Source: Provided by Wikipedia 2015.

1.1.3 Infection and replication

Like other herpes viruses, the EBV replication cycle in B lymphocytes and epithelial cells includes a latent phase and a lytic phase, corresponding to a quiescent infection and active replication, respectively. Humans are the only natural host for EBV (IARC 2012). Latent infection is divided into several phases based on the viral genes that are expressed (latency 0, I, II, and III) (Table 1-1).

Table 1-1. EBV transcription programs in normal B cells and in associated malignancies

Transcription program ^a	Gene products	Function in normal B cell type ^b	Associated malignancies
Latency 0	EBERs	Resting peripheral memory B cells; lifetime infection	None
Latency I	EBNA-1, EBERs	Dividing peripheral memory B cells	Burkitt lymphoma and gastric carcinoma ^d
Latency II	EBNA-1, LMP-1, LMP-2A, EBERs	Activated B cell differentiates into memory cell in germinal center	NPC, Hodgkin lymphoma, NK and T cell lymphoma, nasal type ^e , LEC of the salivary gland
Latency III	EBNA-1, -2, -3A, -3B, -3C, -LP, -1, LMP-2A, LMP-2B, EBERs	Naïve B cell infected and activated	Immunosuppression related NHL (AIDS-associated, post-transplant disorder, iatrogenic)
Lytic	All 9 lytic genes	Replicates in plasma cell	May be present in some malignancies ^f

Source: Adapted from Yau *et al.* 2014, IARC 2012.

EBER = EBV-encoded small RNA; EBNA = EBV nuclear antigen; EBNA-LP = EBV nuclear antigen leader protein; LMP = latent membrane protein; LEC = lymphoepithelial cancer; NHL = non-Hodgkin lymphoma; NPC = nasopharyngeal carcinoma.

^aIn latency I, II, III micro-RNAs (at least 22) are expressed in various amounts.

^bCell types are primarily restricted to lymphoid tissue of Waldeyer ring (i.e., tonsillar tissue in the oropharynx).

^cGeneral classification of disease and heterogeneous patterns may be present in different cell populations within the same individual.

^dExpression of lytic genes and LMP-2A is variable.

^ePreviously called sinonasal angiocentric T-cell lymphoma.

^fEBV lytic genes expressed in human lymphoblastoid cell lines (Arvey *et al.* 2012).

In a primary EBV infection spreads through the saliva and infects the B cells in the tonsil, oropharyngeal epithelial mucosa, and salivary glands by a poorly understood mechanism (IARC 2012, Raab-Traub *et al.* 1991). Within the oropharyngeal epithelium the infection is primarily lytic (it is unknown whether epithelial cells support latent infection) leading to amplification of the number of viruses. Within the tonsil, EBV infects local naïve B lymphocytes entering the germinal center in the underlying lymphoid tissues by interaction of the viral envelope protein gp350/220 with the CD21 protein on B lymphocytes. Viral entry into B cells requires a complex of viral glycoproteins gH, gL, and gp42 with gp42 binding to class II human leukocyte antigen on the lymphocyte (IARC 2012, Tugizov *et al.* 2003). Virions containing only the gH-gL complex can infect epithelial cells while virions require the gH-gL-gp42 complex to infect naïve B cells (Tugizov *et al.* 2003). Inside the B lymphocytes, the virus enters the latency III phase where replication is suppressed by the expression of three EBV nuclear antigens (EBNA-3A, EBNA-3B, and EBNA-3C). The B lymphocytes then can enter a germinal center in the tonsils and, in the immunocompetent host, are destroyed by cytotoxic T cells specific for viral proteins expressed during latency III (Thorley-Lawson *et al.* 2004).

Germinal centers are structures in the lymphoid tissues where antibody affinity maturation occurs through clonal proliferation of antigen-exposed B lymphocytes, diversification of antigen affinity through somatic hypermutation of immunoglobulin genes and of antibody type through class switch recombination (Ponce *et al.* 2014, Basso and Dalla-Favera 2015). While in the germinal center, the EBV infection enters latency II, in which gene expression is focused on survival signals. Within the germinal center, lymphocytes differentiate to memory B cells and antibody-producing plasma cells, both of which enter the peripheral circulation. Within resting B memory cells, the EBV infection is in the latency 0 phase, in which no genes are expressed, except only those coding for EBV-encoded small RNAs (EBER-1 and EBER-2). Occasionally, memory B cells will replicate and EBV will enter latency I, where expression of EBV nuclear antigen 1 (EBNA1) allows for replication and segregation of the viral episome to daughter cells. B cells that are activated and differentiate into antibody-producing plasma cells allow EBV to express lytic genes and enter the lytic cycle of replication. Lytic genes comprise 3 immediate early genes that initiate the lytic cycle, 10 early genes that enable virus replication, and 9 genes that enable packaging and release of the virion from the host cell (see IARC 2012 for discussion of lytic genes). Complete viral lytic replication does not occur in tissues other than mucosa-associated lymphoid tissue (MALT) and tonsillar tissue in the oropharynx (termed Waldeyer's ring). In these tissues lytic replication occurs along with cell death and shedding of virions into the saliva or re-infection of B lymphocytes repeating the cycle (Figure 1-2). While the EBV infection is in the latent I, II, or III phase, EBV-specific cytotoxic T cells and NK cells are able to recognize the expressed viral proteins and limit the infection, but within resting memory B cells (latency 0 phase), the virus is able to evade immune detection and provide a site for maintenance of long-term EBV infection (Thorley-Lawson *et al.* 2004).




Figure 1-2. Epstein-Barr Virus life cycle

Source: Adapted from Vetsika and Callan 2004.

Although EBV usually infects B lymphocytes through a CD21-dependent pathway, it also can infect some types of epithelial cells and T cells by several CD21-independent pathways (IARC 2012). Primary epithelial cells in culture can only be infected with EBV by co-culturing with EBV-producing B lymphoblastoid cells, suggesting that cell-to-cell contact is necessary for infection of epithelial cells (Imai *et al.* 1998). It has been demonstrated that viral envelope proteins (gHgL) can interact with an integrin complex on the epithelial cell surface, triggering cell entry, and that gp42 interferes with binding to the integrin complex (Chesnokova and Hutt-Fletcher 2011, Borza and Hutt-Fletcher 2002). Additionally, virions released from epithelial cells express gp42 on their surface, whereas those released from lymphocytes do not have this protein marker. It is believed that this dual cell tropism enables the virus to shuttle between lymphocytes and epithelial cells. Other work using polarized basal epithelial cells identified another EBV glycoprotein, BMRF2, which enabled infection via interaction with epithelial cell marker $\alpha 5\beta 1$

(Tugizov *et al.* 2003). Further, they showed that EBV virions could also be transmitted by direct cell-to-cell contact of B lymphocytes to apical epithelial cell membranes, and epithelial cell-to-cell infection could occur via lateral membrane transmission.

During latency, the virus can express a variable pattern of viral genes that are essential for host adaptation and in some cases also promote carcinogenesis or disease (IARC 2012). The role of EBV in the pathogenesis of various cancers is an active area of investigation and different EBV latent gene expression patterns are associated with different types of cancers (Table 1-1).

1.2 Detection

EBV-specific serologic methods or amplification of EBV DNA from peripheral white blood cells can be used to detect EBV infection in healthy carriers (Table 1-2). In developed countries, healthy carriers are negative for EBV DNA in serum (cell-free) by the quantitative polymerase chain reaction (Q-PCR) method (IARC 2012). A positive result using the latter method indicates EBV-associated diseases or EBV reactivation. EBV infection results in production of IgG, IgM, and occasionally IgA antibodies against viral proteins and glycoproteins (e.g., viral capsid antigen [VCA] or EBNA1) and these can be detected using antigen-antibody binding assays such as enzyme immunoassays (EIA) to determine antibody titer.

Detection methods are also used to determine EBV viral load, reactivation, response to treatment, and presence in tumor cells (IARC 2012). In general, high antibody titers to some EBV antigens (e.g., VCA) are detected in patients with endemic Burkitt lymphoma. Historically, early markers used for detection of EBV in Burkitt lymphoma were antibodies to early antigen (EA) and viral capsid antigen (VCA). Determination of viral load can give an indication of the degree of infection or of response to treatment and can be determined in tumor cells or in lymphocytes by Q-PCR. For example, nasopharyngeal carcinoma brushings from patients can be used to determine viral load by Q-PCR. Reverse transcriptase PCR can detect EBNA-1 or BART1 RNA in nasopharyngeal cells in these samples, whereas no EBV RNA is found in healthy donors. In addition, *in situ* hybridization and RT-PCR are highly sensitive methods for EBV detection in Hodgkin disease.

Table 1-2. Methods of EBV detection

Biomarker	Detection Method	Examples	Tissue
Antibodies to VCA, EBNA-1, CF/S antigens, neutralizing anti-gp350, EA	Serologic, indirect IFA, EIA	Detected in healthy EBV carriers, EA detected in some carriers; higher titers with Burkitt lymphoma	Serum
EBV DNA	Polymerase chain reaction (PCR)	Detection in healthy carriers	White blood cells
Cell-free EBV DNA	Quantitative PCR	EBV-associated diseases or EBV reactivation; healthy carriers are negative	Serum
EBV DNA	Quantitative PCR	Determination of viral load per cell	Tumor cells; white blood cells

EBER-1, EBER-2 RNAs	Reverse transcription - PCR and <i>in-situ</i> hybridization	EBV infection; highly sensitive for Hodgkin disease detection	Tumor cells, tissue biopsy
EBNA-1 and BARF1 RNA	Reverse transcription-PCR	NPC detection	NPC biopsy or brushing

Source: Adapted from IARC 2012.

BARF1 = *Bam*HI A rightward fragment 1 (micro RNA); BL = Burkitt lymphoma; CF/S = complement-fixing soluble antigen; EA = early antigens (EA-D encoded by *BMRF-1* and EA-R encoded by *BHRF-1*); EBER = small nuclear EBV-encoded RNA; EBNA = EBV nuclear antigen; EIA = enzyme immunoassay; gp350 = glycoprotein 350; IFA = indirect immunofluorescence assay; LMP-1 = Latent membrane protein; NPC = nasopharyngeal carcinoma; PCR = polymerase chain reaction; VCA = viral capsid antigen.

1.3 Summary

Epstein-Barr Virus (EBV), also called human herpesvirus 4 (HHV4), is an enveloped dsDNA virus in the gammaherpesvirus subfamily and consists of two major types (EBV-1, and EBV-2). EBV predominantly enters the host through contact with saliva and infects B cells in the tonsils. The virus exists as an episome, but the genome circularizes during the lytic phase of infection when virions are produced and are released to infect new cells. The latent phase is complex and changes depending on the status of the host cells (latency 0, I, II, III). EBV proteins expressed on infected cells can be recognized by cytotoxic T cells and NK cells, which attack and destroy them. However, EBV in latency 0 phase, typically found in resting memory B cells, does not express proteins on infected cells, which enables it to evade immune recognition. EBV infection can be detected by measuring anti-EBV antibodies in serum and EBV DNA or RNA in peripheral white blood cells, which can indicate viral load.

2 Exposure

This section discusses transmission and prevalence (Section 2.1) and non-cancer diseases, prevention, and treatment for Epstein-Barr virus (EBV) (Section 2.2).

2.1 Transmission and prevalence

Transmission of EBV is primarily via saliva (IARC 2012). The presence of EBV in peripheral blood suggests that transmission via blood is also possible, and transmission among transfusion recipients and organ recipients has been reported. Infected cells, primarily resting memory B cells in peripheral blood, provide a permanent reservoir for progeny to disseminate within the body and infect other hosts.

More than 90% of adults worldwide are infected with EBV (IARC 2012). Age at primary infection varies, occurring during infancy in developing countries and during adolescence and young adulthood in developed countries, perhaps due to better hygienic conditions, and other socioeconomic and demographic factors (e.g., household size and population density), which result in later age of exposure to infected saliva (Dowd *et al.* 2013, IARC 2012, Piriou *et al.* 2012, Biggar *et al.* 1978a, 1978b). The two major types of EBV (EBV-1 and EBV-2) (see Section 1) differ in geographic distribution (IARC 2012). Immunocompromised subjects more often harbor both types; EBV-2 may be more common in Africa and in men who have sex with men.

The seroprevalence of EBV antibody in the United States based on NHANES data collected in 2009 and 2010 ranged from 50% in 6 to 8 year olds to 89% in 18 to 19 year olds (Balfour *et al.* 2013, Dowd *et al.* 2013). Lower prevalence of EBV antibody has been shown to be associated with higher socioeconomic status within race/ethnicity groups. An analysis of 782 serum samples from Minnesota children 18 months to 19.9 years old indicated that a combination of genetics, family practices, and home environment were responsible for racial/ethnic differences in EBV antibody prevalence among young children and noted that the route of EBV transmission to preadolescents remains unclear (Condon *et al.* 2014).

2.2 Diseases, prevention, treatment

Most individuals who are infected with EBV remain otherwise healthy and are asymptomatic (IARC 2012). Infection is life-long and is subclinical when it occurs in early childhood (IARC 2012); however, it results in infectious mononucleosis in at least 25% of teenagers and young adults infected with EBV (CDC 2014a). Oral hairy leukoplakia results from infection with EBV in the context of immunosuppression (e.g., human immunodeficiency virus-1 [HIV-1]) or immunosenescence (aging) (Auwaerter 2015). Chronic uncontrolled EBV (with high EBV DNA in blood) occurs frequently in Asia and South America, but occurs rarely in the United States and Europe. Its etiology is unknown, but is believed to involve environmental co-factors and/or rare genetic abnormalities that impair immune control of EBV infection (Cohen 2009, Chaigne-Delalande *et al.* 2013, Rigaud *et al.* 2006).

Because EBV transmission is associated with EBV shedding in saliva, avoiding salivary exposure (e.g., not kissing or sharing drinks, food, or personal items like toothbrushes with people who have EBV infection) may theoretically prevent transmission (CDC 2014b).

Some drugs have been reported to reduce or inhibit EBV shedding (e.g., see Auwaerter 2015); however, currently there are no FDA-approved drugs for treatment of EBV infection. Currently, there is no vaccine against EBV; efforts to develop a vaccine are ongoing (ACS 2015a, Cohen 2015, FDA 2015, Balfour 2014, CDC 2011).

2.3 Summary

The high seroprevalence rate for Epstein-Barr virus (EBV) in the U.S. population indicates that a significant number of people living in the United States are exposed to EBV. Estimates for the seroprevalence for EBV in the United States range from 50% for 6 to 8 year olds to 89% for 18 to 19 year olds, based on detection of the EBV antibody; however, these levels vary based on socioeconomic status and within race/ethnicity groups. EBV is transmitted primarily via saliva, but transmission via blood is possible since EBV is present in peripheral blood and transmission among transfusion recipients and organ recipients have been reported. The infection rate worldwide is very high, likely exceeding 90%, but the age at primary infection varies geographically, with more developed countries having higher ages for primary infection due to better hygiene. EBV infections tend to be asymptomatic for most individuals, but the infection is life-long and results in infectious mononucleosis in at least 25% of teenagers and young adults infected with EBV. Prevention of transmission of EBV can theoretically be achieved by limiting exposure to saliva from kissing or sharing drinks, food, or personal items like toothbrushes between infected and non-infected individuals. There is currently no vaccine against EBV, although efforts to develop a vaccine are ongoing.

3 Human Cancer Studies

Introduction

The NTP used the body of knowledge published by IARC monograph (1997, 2012) on EBV as the resources for studies conducted up to and including 2008, together with any new human studies identified from 2008 to 2015 to evaluate the scientific evidence for specific cancer endpoints independently of IARC's conclusions. Tumor endpoints identified in the monographs and other endpoints with sufficient data, are considered in the RoC monograph. Where available, IARC data tables of the effect estimates have informed the cancer hazard assessment (IARC 2012), with links to these tables made available in the text. When tables of individual studies were not available, forest plot summaries of the data have been provided. The seven cancer endpoints include: Burkitt lymphoma, Hodgkin lymphoma, immunosuppression-related non-Hodgkin lymphoma, extranodal NK/T-cell lymphoma, nasopharyngeal carcinoma, gastric cancer, and lymphoepithelial carcinoma of the salivary gland.

EBV exposure is detected in the human studies via serological measures or from DNA in the tumor. Earlier studies included in the evaluation primarily used serological measures, while later studies used tumor DNA. (See Section 1.2 above for details on EBV detection methods.)

This evaluation of the human cancer hazard associated with EBV is divided into three following parts. The first, a summary of the approach for selection of the studies is provided in Section 3.1. Next, the cancer hazard evaluation for each endpoint is presented in Sections 3.2 to 3.8, and lastly, a summary of the evaluations across endpoints is provided in Section 3.9.

3.1 Selection of the literature

A systematic literature search of major databases, citations, and other authoritative sources for literature from 2009 to August 2015 was conducted. The literature search strategy (including the databases, search terms, and other sources for identifying literature) and procedures and results for selecting the literature (systematic screening procedures and inclusion/exclusion criteria) are described in Appendix A. For the EBV evaluation, all post-2008 case-control and cohort studies for the seven cancer endpoints were identified and included in the evaluation. These studies may range from broadly defined, non-matched hospital or population case-control designs to formal age-, sex, and race-matched case-control designs. Previous studies reviewed by IARC were included in the overall assessment, but not evaluated in depth. Case reports and case series were excluded from further review, although summaries of multiple case report or case-series studies are noted and may be considered in the overall evaluation. In addition, new cancer endpoints with sparse databases have not been assessed.

3.2 Cancer evaluation: Burkitt lymphoma

3.2.1 Background information

Burkitt lymphoma includes three subtypes defined according to their incidence in populations: endemic, sporadic, and immunodeficiency-related Burkitt lymphoma. Endemic Burkitt lymphoma (also known as the African type) occurs primarily in children aged 5 to 9 years in equatorial Africa and Papua New Guinea. It occurs with an incidence rate of 5 to 10 cases per

100,000 in children under 16, and is responsible for 30% to 70% of all childhood cancers in equatorial Africa (IARC 1997). Among endemic Burkitt lymphoma cases, EBV is detected in the tumor in more than 95% of cases (Thompson and Kurzrock 2004). In the case-series studies reviewed in IARC (1997), there were a total of 191 cases of endemic Burkitt lymphoma, with EBV DNA or antibodies present in 185 of the cases.

Sporadic Burkitt lymphoma is found throughout the world. It is rare, with an incidence of 2 to 3 cases per million people in the United States. Sporadic Burkitt lymphoma is associated globally with a lower EBV prevalence, with approximately 20% to 30% of cases being positive in the tumor. Among the case series and reports previously reviewed by IARC (1997), there were a total of 383 cases of sporadic Burkitt lymphoma, with EBV DNA or antibodies present in 192 cases. In the United States, 15% to 30% of sporadic Burkitt lymphoma cases are associated with EBV (Thompson and Kurzrock 2004). Immunodeficiency-related Burkitt lymphoma has been identified in approximately 40% of HIV-associated lymphomas (Gloghini *et al.* 2013, Stefan *et al.* 2011) and may also be associated with other factors resulting in immune suppression such as anti-rejection therapies or congenital immunodeficiency (Carbone *et al.* 2008). EBV has been detected in 30% to 60% of immunodeficiency-related Burkitt lymphoma cases (Young and Rickinson 2004, Mbulaitege *et al.* 2014).

3.2.2 Case-control studies

Twelve case-control studies conducted since 1969 were identified that have investigated the association between EBV and Burkitt lymphoma. Eleven of these were included in the IARC (1997, 2012) reviews.

Briefly, seven case-control studies of endemic Burkitt lymphoma, with a total of 904 cases, were reviewed by IARC (2012, 1997), all showing a statistically significant, positive association between EBV and Burkitt lymphoma (Figure 3-1). These case-control studies found that those with Burkitt lymphoma were more likely to have detectible or elevated levels of EBV titers with odds ratios (ORs) ranging from 2.9 to 52 (Carpenter *et al.* 2008, Mutalima *et al.* 2008, Hirshaut *et al.* 1973, Henle *et al.* 1971, Klein *et al.* 1970, Henle *et al.* 1969). Moreover, four of the seven studies found a dose-response relationship between EBV titer levels and Burkitt lymphoma, with ORs increasing as titer levels increased (Carpenter *et al.* 2008, Mutalima *et al.* 2008, Henle *et al.* 1971, Henle *et al.* 1969). Additionally, the geometric mean titer levels of Burkitt lymphoma patients were significantly higher than controls in the three studies that reported these findings (Nkrumah *et al.* 1976, Hirshaut *et al.* 1973, Henle *et al.* 1969). These studies investigated a variety of EBV antibodies including viral capsid antigen (VCA) and early antigen (EA), while other studies (Hirshaut *et al.* 1973, Henle *et al.* 1969) did not specify the type of EBV antibody under study. Regardless of the type of EBV antibody investigated, a significant, positive relationship was seen between endemic Burkitt lymphoma and EBV. Figure 3-1 shows a forest plot with the ORs of the associations between endemic Burkitt lymphoma and EBV antibodies.




Figure 3-1. Forest plot of serological case-control studies of endemic Burkitt lymphoma and Epstein-Barr virus

Sources: All endemic Burkitt lymphoma studies were conducted in sub-Saharan Africa. (Henle *et al.* 1969) - Kenya, Uganda, Nigeria; (Klein *et al.* 1970) - country not specified; (Henle *et al.* 1971) - Uganda, Kenya; (Hirshaut *et al.* 1973) - Uganda; (Carpenter *et al.* 2008) - Uganda; (Mutalima *et al.* 2008) - Malawi; (Mulama *et al.* 2014) - Kenya. Note: One study, Nikrumah *et al.* (1976) reported mean levels of EBV antibodies only but not enough information to calculate an OR.

ORs in brackets [] were calculated by NTP; if OR could not be calculated due to a cell size of 0, 1 was added to each cell.

^aOptical density (OD) measures were used as a surrogate measure for EBV VCA antigen.

CI = confidence interval, EA = early antigen, OR = odds ratio; VCA = viral capsid antigen.

Since the publication of the second IARC review (2012), there has been one additional hospital-based case-control study in Kenya on the relationship between endemic Burkitt lymphoma and EBV. This study (Mulama *et al.* 2014) investigated the association between cellular EBV load and endemic Burkitt lymphoma among 89 children with confirmed Burkitt lymphoma with 213 controls frequency matched on age range and malaria exposure from the Naynza and Rift Valley provinces in Kenya. These control sites were chosen to approximate malarial exposure for cases because Naynza province has high malarial transmission rates while the Rift Valley province has low rates. Though not reported in the manuscript, based on individual case and control level data reported in the figures, children with endemic Burkitt lymphoma were statistically significantly

more likely to have ≥ 2 log EBV copies per μg of human DNA than controls (OR = 16.2; 95% CI = 8.0 to 32.5). The results of this study are included in Figure 3-1.

Five case-control studies of sporadic Burkitt lymphoma, with a total of 113 cases, were reviewed by IARC (1997) (Cavdar *et al.* 1994, Gottleib-Stematsky *et al.* 1976, Ablashi *et al.* 1974, Hirshaut *et al.* 1973, Levine *et al.* 1972). Of the five studies, four investigated the relationship between Burkitt lymphoma and VCA antibodies, while one was unspecified (Hirshaut *et al.* 1973). Gottleib-Stematsky *et al.* (1976) also investigated EA antibodies in their analysis. Although 4 of these 5 case-control studies reported odds ratios of at least 2.0, only one reported a statistically significant association between the presence of EBV antibodies and Burkitt lymphoma (OR = 4; 95% CI = 1.3 to 12.0) (Levine *et al.* 1972). The ODs ranged from 1 to 6.9 for those with statistically non-significant findings. In contrast to endemic Burkitt lymphoma studies, many of these studies included small numbers of cases and controls and had limited statistical power. Three of the five studies found mean EBV titers to be significantly higher in cases than in controls (Cavdar *et al.* 1994, Ablashi *et al.* 1974, Levine *et al.* 1972), while the other two studies found no statistically significant differences in means between cases and controls (Gottleib-Stematsky *et al.* 1976, Hirshaut *et al.* 1973). These studies are presented in Figure 3-2, which shows a forest plot with the ORs of the associations between sporadic Burkitt lymphoma and EBV antibodies.




Figure 3-2. Forest plot of serological case-control studies of sporadic Burkitt lymphoma and Epstein Barr Virus

Sources: Sporadic Burkitt lymphoma study locations: (Levine *et al.* 1972) – U.S.A.; (Hirshaut *et al.* 1973) – U.S.A.; (Ablashi *et al.* 1974) – U.S.A.; (Gotlieb-Stematsky *et al.* 1976) - Israel; (Cavdar *et al.* 1994) – Turkey.

CI = confidence interval, EA = early antigen, OR = odds ratio, VCA = viral capsid antigen.

No case-control studies on immunodeficiency-related Burkitt lymphoma have been published since the more recent review by IARC (2012).

3.2.3 Cohort studies

Only one cohort study has been identified that investigated the relationship between EBV and Burkitt lymphoma. This study was reviewed by IARC (1997) and is briefly reviewed here. Beginning in 1972, a large-scale prospective study was begun in a Burkitt lymphoma-endemic area of northern Uganda, which collected blood samples from approximately 42,000 healthy children under age 8. The first follow-up of this cohort (1972 to 1977) was first presented by de Thé *et al.* (1978) which reported on 14 cases (13 histologically confirmed Burkitt lymphoma cases and one unclassified lymphoma), and the second follow-up of the cohort was presented by Geser *et al.* (1982) adding 2 additional cases to the analysis (1978 to 1979). Randomly selected controls, matched on age, gender, and location, were selected for a nested case-control analysis (de Thé G *et al.* 1978). Statistically significantly higher titers of VCA antibodies were seen in the pre-diagnosis sera of the Burkitt lymphoma cases compared with the controls (geometric mean titer of Burkitt lymphoma cases = 425.5; control = 125.8, $P = 0.01$). When the additional two cases were added, the difference in VCA titers between cases and controls increased slightly ($P < 0.001$) (Geser *et al.* 1982). The mean values of EA and Epstein-Barr nuclear antigen (EBNA) in cases and matched controls were not provided; however, no statistically significant differences

were seen in the titer levels for EA and EBNA antibodies between the Burkitt lymphoma cases and the matched controls. There were no differences between pre-diagnosis cases and controls in regard to malarial parasites or antibody titers to herpes simplex virus, cytomegalovirus or measles reported by de Thé *et al.* (1978). A conditional regression analysis including all 16 cases was conducted by Geser *et al.* (1982) who found that the relative risk (RR) increased by a factor of 5.05 for each dilution for which the VCA titer is above the general population average. Moreover, the RR increased to 9.16 (95% CI not reported) when limited to cases that demonstrated the presence of EBV DNA in the tumor genome (N = 9). The results of this prospective cohort study demonstrate the temporality of the EBV/Burkitt lymphoma relationship by showing that elevated EBV antibodies, particularly VCA antibodies, precede the development and diagnosis of Burkitt lymphoma in this population.

3.2.4 Cofactors

Potential cofactors for endemic Burkitt lymphoma were reviewed by IARC (2012, 1997). These co-factors include malaria, sickle-cell trait, and ingestion of *Euphorbia tirucalli* and other medicinal plants.

In earlier ecological studies, the geographic relationship between malaria and endemic Burkitt lymphoma was noted, with Burkitt lymphoma prevalence highest in areas with the highest malaria transmission rates (Morrow 1985, as cited in IARC 1997). The relationship between malaria and endemic Burkitt lymphoma was further apparent after the decline in endemic Burkitt lymphoma incidence following large-scale malaria eradication efforts. Geser *et al.* (1989) further confirmed this relationship in an intervention study designed to reduce Burkitt lymphoma incidence by reducing the prevalence of malaria. Moreover, two case-control studies published in 2008 found an increased risk of endemic Burkitt lymphoma as antibody titers to malaria increased (Carpenter *et al.* 2008, Mutalima *et al.* 2008). Based on the evidence of these studies, endemic Burkitt lymphoma is strongly associated with both EBV and with *Plasmodium* co-infection (IARC 2012) and there may be a synergistic effect between malaria and EBV in the development of Burkitt lymphoma (Carpenter *et al.* 2008, Mutalima *et al.* 2008). There is further evidence that malaria reduces the T-cell mediated immunosurveillance of EBV-infected cells, leading to an increased viral load of EBV (Moormann *et al.* 2009, as cited in IARC 2012).

A recent study (Mulama *et al.* 2014) did not find an association between sickle-cell trait and endemic Burkitt lymphoma (OR = 0.85; 95% CI = 0.61 to 1.17). Moreover, there was no association between sickle-cell trait genotype and EBV cellular viral load, suggesting sickle-cell trait is not a cofactor in the relationship between EBV and Burkitt lymphoma.

The relationship between endemic Burkitt lymphoma and *Euphorbia tirucalli* and other medicinal plants is unclear, with few studies available to provide epidemiological evidence relevant to this relationship. Several case series and two case-control studies on these relationships were reviewed by IARC (1997). Both case-control studies found a significant positive association between plant use and Burkitt lymphoma. Other studies have demonstrated that extracts of *Euphorbia tirucalli* and other related plants can induce the expression of EA and VCA antibodies, and increase EBV replication (Lin *et al.* 1982, as cited in IARC 1997, Ito *et al.* 1981). No new studies on these relationships have been identified since the IARC (1997) publication.

3.2.5 Integration of the evidence

The epidemiological data suggest an association between EBV and both endemic and sporadic Burkitt lymphoma. All seven case-control studies and one cohort study of the relationship between endemic Burkitt lymphoma and EBV found positive, statistically significant relationships. There were 5 case-control studies on the relationship between sporadic Burkitt lymphoma and EBV. Although positive odds ratios ranging from 2.2 to 6.9 were seen for 4 of the 5 studies, only one was statistically significant. These studies on sporadic Burkitt lymphoma and EBV had limited power to detect an effect due to small numbers of cases and controls. The overall results of the individual studies are heterogeneous, particularly studies on endemic Burkitt lymphoma, for which ORs ranged from 2.9 to 30.0. One possible explanation is the variety of serological markers used by the different studies to detect EBV exposure. Another possible cause of the observed heterogeneity is the lack of variability in the exposure, particularly among cases, with up to 100% of cases exposed to EBV in some studies.

3.3 Cancer evaluation: Hodgkin lymphoma

3.3.1 Background information

Hodgkin lymphoma is categorized into four histological subtypes (lymphocyte predominance, nodular sclerosis, mixed cellularity, and lymphocyte depletion). In the United States, the age-adjusted incidence rate for all types of Hodgkin lymphoma is 2.7 cases per 100,000 person-years, with a 5-year survival of 80% (SEER 2015). The relationship between EBV and Hodgkin lymphoma was first proposed in the 1960s (MacMahon 1966). Numerous case reports and case series studies since that time have examined the association between EBV and primarily the mixed cellularity subtype, with approximately 75% of cases EBV positive; while approximately 20% of Hodgkin lymphoma cases with the nodular sclerosis subtype are EBV positive (Flavell and Murray 2000, Weiss 2000). These case reports and case series have reported that the association between EBV and Hodgkin lymphoma varies by age and geographic region. EBV-related Hodgkin lymphoma appears to be mostly highly associated with Hodgkin lymphoma incidence in middle adulthood in developed countries, while in developing countries, the EBV-Hodgkin lymphoma relationship shows a bimodal age distribution, with rates of EBV-related Hodgkin lymphoma highest in childhood and in older adults (Flavell and Murray 2000, Weiss 2000). EBV has been detected in 20% to 50% of Hodgkin lymphoma cases in North America and Europe, though the percentage differs by Hodgkin lymphoma subtype (Weiss 2000). EBV seropositivity in Asian Hodgkin lymphoma patients is around 65%, and 90% to 100% in South Americans and Africans. Additionally, Hodgkin lymphoma patients with HIV have a nearly 100% EBV infection rate (Weiss 2000).

EBV has been established as the cause of infectious mononucleosis, and a large body of studies has reported an increased risk of Hodgkin lymphoma among populations with infectious mononucleosis infection (Crawford 2001). Later (after childhood) exposure to EBV in developed countries, which is more likely to lead to symptomatic infectious mononucleosis infection, is suggested to be a risk factor for the development of Hodgkin lymphoma, particularly in younger adults (Ambinder and Cesarman 2007, Jarrett 2003, as cited in IARC 2012). Several case-control and cohort studies have investigated the relationship between EBV and Hodgkin lymphoma, as well as Hodgkin lymphoma and infectious mononucleosis, as described below.

3.3.2 Case-control studies

Thirty-eight published case-control studies were identified that investigated the relationship between Hodgkin lymphoma and EBV. All but one study was previously reviewed by IARC (2012, 1997). These studies included a total of over 7,100 Hodgkin lymphoma cases. These studies have generally fallen into two categories: those investigating the relationship between Hodgkin lymphoma and EBV serology or EBV DNA directly and those investigating this relationship indirectly via the association between infectious mononucleosis and Hodgkin lymphoma.

Hodgkin lymphoma and EBV serology or EBV DNA

Twenty-seven of a total of 38 case-control studies on EBV and Hodgkin lymphoma were previously reviewed by IARC (2012, Table 2-2; 1997, Tables 19 to 21), with 1 new study (Linabery *et al.* 2014) that examined the relationship between EBV antibodies or DNA and Hodgkin lymphoma, ORs for the 22 studies reporting ORs (or information to calculate ORs) are presented in Figure 3-4. Twenty-three studies (ORs for 19 studies are graphed in Figure 3-4) looked at the relationship between EBV serology and Hodgkin lymphoma. In general, these studies investigated the association between high VCA titer levels among Hodgkin lymphoma cases and controls, with odds ratios ranging from 0.8 to 67, with the majority reporting ORs between 4 and 19. Findings from studies that did not specify the type of EBV antibodies were consistent with the studies of specific antibodies; most studies (5/8) reported statistically significant ORs ranging from 4 to 11. Ten of the 21 total studies (on VCA or unspecified antigens) additionally analyzed the association between EA antibodies and Hodgkin lymphoma between cases and controls, with ORs ranging from 1.2 to infinity; however, when studies with no exposure variability (studies with either no unexposed cases, or no exposed controls) were excluded, the ORs ranged from 1.2 to 15. There were six case-control studies that also reported risk estimates (or information to calculate estimates) for elevated titers of EBNA and Hodgkin lymphoma. These included Lange *et al.* (1978) with an OR of 19.3 (95% CI = 5.5 to 67.6, for 15/28 exposed cases), Mochanko *et al.* (1979) with an OR of 5.4 (95% CI = 1.8 to 15.8, for 18/37 exposed cases), Merk *et al.* (1995) with an OR of 1.7 (95% CI = 0.8 to 3.6, for 16/61 exposed cases), and Berrington de Gonzalez *et al.* (2006)¹ with an OR of 0.7 (95% CI = 0.3 to 1.4, for 21/83 exposed cases) and Rocchi *et al.* (1975) and Shope *et al.* 1982 reported OR of infinity (no controls had elevated titer). Three serological case-control studies did not provide enough data to calculate an odds ratio (Lennette *et al.* 1995, Lennette *et al.* 1993, Wutzler *et al.* 1983).¹ In addition to the serology studies measuring antibody, three case-control studies reported ORs between EBV DNA in serum or lymph nodes and Hodgkin lymphoma; these studies found highly statistically significant associations, with ORs ranging from 120 to infinity; however, there was little variability in exposure (Dinand *et al.* 2007, Musacchio *et al.* 2006, Lei *et al.* 2000). Gallagher *et al.* (1999) also measured EBV DNA in serum of case and controls, but it was not possible to calculate an OR as cases were pre-classified as EBV positive or negative.

¹ This study did not report an OR for other antibodies.




Figure 3-4. Forest plot of serological case-control studies of Hodgkin lymphoma and Epstein-Barr Virus.

Source: Table 19 in [IARC 1997](#); Table 2.3 in IARC 2012. Note: Hilgers and Hilers (1976) was not included ORs in curly brackets { } were calculated by IARC working group; ORs in brackets [] were calculated by NTP. CI = confidence interval, OR = odds ratio, VCA = viral capsid antigen.

^aExposed group - any detectible EBV antibody.

^bEBV antibody exposed group ≥ 160.

^cEBV antibody exposed group ≥ 640.

^dEBV antibody exposed group ≥ 1280.

^eEBV VCA antibody exposed group ≥ 5 .

^fEBV VCA antibody exposed group ≥ 160 .

^gEBV VCA antibody exposed group ≥ 320 .

^hEBV VCA antibody exposed group ≥ 1280 .

ⁱStudy also includes data on EA antibodies.

^jStudy also includes data on EBNA antibodies.

The study by Linabery *et al.* (2014) examined a subset (69%) of the 517 cases of pediatric Hodgkin lymphoma for EBV RNA in the tumor. Overall, 16% (N = 84) were found to have tumors that were EBV RNA positive. This proportion increased in the younger age groups, with 23% (N = 5) of those aged 0 to 4 years and 29% (N = 36) of those aged 5 to 9 years EBV positive. EBV status among the controls was not available; therefore, no odds ratios could be calculated.

Hodgkin lymphoma and infectious mononucleosis

Eleven of the 38 case-control studies (10 reported by IARC and 1 new study, Linabery *et al.* [2014]) reported on the association between infectious mononucleosis and Hodgkin lymphoma. These studies generally found a positive association between infectious mononucleosis and Hodgkin lymphoma, with positive ORs ranging between 1.0 and 13.1, with most falling between 1.0 and 3.0 (Linabery *et al.* 2014, Hjalgrim *et al.* 2007, Glaser *et al.* 2005, Alexander *et al.* 2003, Alexander *et al.* 2000, Serraino *et al.* 1991, Bernard *et al.* 1987, Evans and Gutensohn 1984, Gutensohn 1982, Gutensohn and Cole 1981, Henderson *et al.* 1979). Among these studies, six found a statistically significant association between Hodgkin lymphoma and infectious mononucleosis (Bernard *et al.* (1987) and Serraino *et al.* (1991) found significant associations in subpopulations, not reported in Figure 3-4). One study (Glaser *et al.* 2005) found a non-significant OR of 0.3 in a population of women, ages 19 to 79, with Hodgkin lymphoma. These studies are summarized in Figure 3-4. A number of these studies had limited statistical power to detect an effect. Additionally, studies that stratified Hodgkin lymphoma patients by age found a stronger, statistically significant association among younger adults (Bernard *et al.* (1987) [young adult males, OR = 4.9; $P = 0.04$; 95% CI not available], Hjalgrim *et al.* 2007, Alexander *et al.* 2000). The risk for infectious mononucleosis and Hodgkin lymphoma in younger age groups resulting from infection with EBV has been suggested to be greater in people with a higher socioeconomic background (Gutensohn 1982, ACS 2015). One theory proposed for this relationship is that exposure to EBV later in life for children from more affluent families might somehow increase their risk for these diseases.

Two studies reported significant associations for cases with the nodular sclerosis subtype of Hodgkin lymphoma. Henderson (1979) reported a non-significant OR of 1.5 (95% CI not reported), while Serraino *et al.* (1991) reported a statistically significant OR of 13.1 (1.0 to 176.6). The most recent study, of children and adolescents (Linabery *et al.* 2014) reported a non-statistically significant increase in the odds of Hodgkin lymphoma among those diagnosed with infectious mononucleosis (OR = 1.5; 95% CI = 0.52 to 3.50 for 9/517 exposed cases) although power was limited by a small number of participants with a previous diagnosis of infectious mononucleosis (9 cases and 10 controls). When restricted to EBV-positive (EBV detected in the tumors) cases, they found a non-statistically significant 9-fold increase in the risk of Hodgkin lymphoma among those who had a history of infectious mononucleosis infection (OR = 9.1; 95% CI = 0.81 to 102.3 for 3/84 exposed cases).




Figure 3-4. Forest plot of case-control studies of Hodgkin lymphoma and infectious mononucleosis

*95% confidence interval unavailable.

CI = confidence interval; OR = odds ratio.

3.3.3 Cohort studies

Seven cohort studies and two nested case-control studies have investigated the association between EBV and Hodgkin lymphoma. These studies are summarized in Table 3-1. The seven cohort studies, conducted in the United States and Western Europe and published from 1973 to 2000, looked at the relationship between infectious mononucleosis (caused by EBV) and Hodgkin lymphoma. Six were reviewed in detail by IARC in 1997 (Kvale *et al.* 1979, Munoz *et al.* 1978, Carter *et al.* 1977, Connelly and Christine 1974, Rosdahl *et al.* 1974, Miller and Beebe 1973), while one additional cohort study was reviewed by IARC in 2012 (Hjalgrim *et al.* 2000). These studies included a total of over 80,000 participants with serologically confirmed infectious mononucleosis. A total of 83 cases of Hodgkin lymphoma were observed, with standardized incidence ratios (SIR) ranging from 2.0 to 5.0. All studies observed more cases of Hodgkin lymphoma than expected in the general population, with three of the seven studies reporting statistically significant SIRs (Hjalgrim *et al.* 2000, Munoz *et al.* 1978, Rosdahl *et al.* 1974) in the relationship between infectious mononucleosis and subsequent development of Hodgkin lymphoma. One study did not find a statistically significant association (Miller and Beebe 1973), and the remaining studies did not report statistical significance in their risk estimates (Kvale *et al.* 1979, Carter *et al.* 1977, Connelly and Christine 1974).

Table 3-1. Summary of cohort and nested case-control studies of Hodgkin lymphoma and Epstein-Barr Virus

Reference	Age group	RR ^a	Comments
Infectious mononucleosis studies^b			
Miller and Beebe 1973	Adults	{2.0}	NS
Connelly and Christine 1974	Adults/children	{5.0}	NR
Rosdahl <i>et al.</i> 1974	Adults/children	{2.8}	Lower CI > 1
Carter <i>et al.</i> 1977	Adults	{2.3}	NR
Munoz <i>et al.</i> 1978	Adults/children	4.0	Lower CI > 1
Kvale <i>et al.</i> 1979	Adults/children	{3.0} ^c	NR
Hjalgrim <i>et al.</i> 2000	Adults/children	2.6	Lower CI > 1
EBV serology studies^b			
Mueller <i>et al.</i> 1989	NR	VCA IgG \geq 320: 2.6	Lower CI > 1
		VCA IgA \geq 20: 3.7 ^d	Lower CI > 1
		EBNA \geq 80: 4.0	Lower CI > 1
		EA(D) \geq 5: 2.6	Lower CI > 1
Lehtinen <i>et al.</i> 1993	Adults	NR	NR

Relative risks in curly brackets { } were calculated by the IARC Working Group (2012).

CI = 95% confidence interval; EA(D) = early antigen-diffuse; EBNA = EBV nuclear antigen; NR = not reported; RR = relative risk; VCA = viral capsid antigen.

^aFor IM cohort studies, RR reported is the standardized incidence ratio (SIR).

^bAll IM studies were cohort designs, while both serology studies were nested case-control designs.

^cSIR at least one year from IM diagnosis.

^dRisk estimate was adjusted for IgM.

Two nested case-control studies investigated EBV serology prior to a Hodgkin lymphoma diagnosis (Lehtinen *et al.* 1993, Mueller *et al.* 1989). Mueller *et al.* (1989) analyzed 43 Hodgkin lymphoma patients and 96 matched controls from a cohort of over 235,000 participants. They found that titers to EBV antibodies (VCA IgA and EBNA) were statistically significantly increased among those with Hodgkin lymphoma, with relative risks equaling 2.7 and 2.5, respectively. Relative risks of antibody titers to VCA IgG and EA were elevated but not significant. Additionally, they found that these associations were generally stronger in patients with a blood draw at least three years before Hodgkin lymphoma diagnosis. Lehtinen *et al.* (1993) conducted a similar nested case-control study among 39,000 adults with a blood draw who were followed for up to 12 years. Data for the association between Hodgkin lymphoma and EBV were not shown, but the authors reported an increased risk of antibody response to EBV among Hodgkin lymphoma patients.

3.3.4 Integration of the evidence

Epidemiological data provide evidence of an association between EBV and Hodgkin lymphoma with the majority finding an increased risk, and 22 of 38 case-control studies showing statistically significant associations. Positive associations have been reported between EBV and Hodgkin lymphoma between both adults and young adults; however, the association between EBV and childhood Hodgkin lymphoma is unclear. Moreover, infectious mononucleosis (caused by EBV) is positively associated with Hodgkin lymphoma among adults, young adults and children, though evidence is inconsistent, possibly due to the fact that few infectious

mononucleosis studies provided data on EBV tumor status. The temporality of this association was seen in one nested case-control study, which reported statistically significantly elevated EBV titers prior to Hodgkin lymphoma diagnosis among cases compared with controls. The strength of the observed associations varied between studies with both indirect EBV exposure measurement (infectious mononucleosis studies), and with direct EBV measurement (serological and tumor DNA studies). ORs are generally higher in the serological studies compared to the infectious mononucleosis studies, and ORs in case-control studies using tumor DNA to detect EBV exposure were highest. The heterogeneity seen in the serology and tumor DNA case-control studies may partly be explained by differential sensitivity and specificity of these exposure markers. One serology case-control study (Gotlieb-Stematsky *et al.* 1975) had only one exposed control, while two of the four EBV DNA studies had no exposed controls, and a third had only one exposed control.

3.4 Cancer evaluation: Immunosuppression-related Non-Hodgkin Lymphoma

3.4.1 Background information

Three types of immunosuppression-related non-Hodgkin lymphomas have been reported in the EBV literature and were reviewed by IARC (2012, 1997). These lymphomas occur with severe immunosuppression and are observed with post-transplant lymphoproliferative disorders (PTLD), HIV-associated lymphoproliferative disorders, and congenital immunodeficiencies. PTLD is a complication of both solid organ transplant and hematopoietic stem cell transplant, and is one of the most common post-transplant malignancies. The incidence of PTLD varies by type of transplant, but is generally more common in children than adults, with incidence rates ranging from < 1% to 13% in children and between 1% and 8% in adults (Garfin *et al.* 2015). The 5-year survival rate for adults and children is approximately 60% (Hauke *et al.* 2001). EBV has been associated with up to 50% to 70% of PTLD cases (Jimenez 2015, Al-Mansour *et al.* 2013).

HIV- lymphoproliferative disorder is a common type of non-Hodgkin lymphoma among those diagnosed with HIV. Non-Hodgkin lymphoma is designated a defining acquired immune deficiency syndrome condition. Those with HIV are 70 times more likely to be diagnosed with non-Hodgkin lymphoma (Grulich *et al.* 2007a) while the 5-year survival rates of those with HIV-related non-Hodgkin lymphoma is low, at approximately 5% (Chow *et al.* 2001). HIV-associated lymphoproliferative disorder is discussed in further detail in the HIV monograph. EBV is present in the tumor cells in almost all cases of HIV-related primary central nervous system non-Hodgkin lymphoma, and in around 50% of HIV-related diffuse large cell and immunoblastic non-Hodgkin lymphoma (Grulich *et al.* 2007b).

3.4.2 Studies and evaluation

IARC reported several case-series studies relating EBV to immunosuppression and non-Hodgkin lymphoma. Detection methods for EBV and the percent of positive cells within the tumor varied. EBV was detected in nine case-series reports of HIV-associated primary central nervous system non-Hodgkin lymphomas and was found in all cases in five studies and in the majority of cases in the remaining four studies (Table 14 in [IARC 1997](#)). EBV was also found in HIV-associated systemic non-Hodgkin lymphomas (Table 15 in [IARC 1997](#)). Sixteen studies identified EBV in some of the cases. In the cases positive for EBV, the number of EBV-positive cells in the tumor

varied from 25% to 100%. Importantly, EBV was found to be monoclonal by terminal-repeat sequence analysis in four of these studies. In addition, seven studies investigated the presence of EBV in lymphomas from patients with congenital primary immune deficiency and all of the cases were positive for EBV (Table 16 in [IARC 1997](#)).

Only one case-control study has been identified that investigated the association between EBV and immunosuppression-related non-Hodgkin lymphoma. This study included cases with several types of lymphoma, including both Hodgkin and non-Hodgkin lymphomas. This study included 13 total cases of different cancer types, including two cases of PTLD and 35 healthy controls in Hong Kong (Lei *et al.* 2000). Plasma EBV DNA was detected in both PTLD cases, suggesting activated EBV; however, EBV plasma DNA was not detected any of the controls.

One nested case-control study and no cohort studies were identified in which EBV titers were measured in immunosuppressed populations. In a U.K.-based nested case-control study, 67 HIV-positive, non-Hodgkin lymphoma patients participated in a trial of antiretroviral therapy with 67 matched controls (Newton *et al.* 2006). Controls were randomly selected from the trial participants among those who had not developed cancer after the same period of follow-up, and matched by trial, age group, sex, HIV transmission group, treatment group, and ethnicity. Among cases, a statistically non-significant association between the risk of disease for a doubling of VCA-IgG antibodies to EBV was observed compared to controls (adjusted OR = 1.5, 95% CI = 0.9 to 2.3).

3.4.3 Integration of the evidence

Severe immunosuppression from congenital, iatrogenic, or HIV/AIDS can result in EBV-associated non-Hodgkin lymphomas. Epidemiological data primarily from case studies provide consistent evidence of an association between EBV positive non-Hodgkin lymphoma and congenital immunodeficiencies. In HIV patients, almost all non-Hodgkin lymphomas of the central nervous system and a large number of systemic non-Hodgkin lymphomas are EBV related. There were some reports of monoclonality of the virus within the tumor, although all cells within the tumor were not positive for the activated virus. Further, EBV has been shown to be associated with over half of PTLD cases. The strength of these studies varied with detection method used, as some studies used a method that would detect EBV, but not necessarily an activated form of the virus.

3.5 Cancer evaluation: Extranodal NK/T-cell lymphoma, nasal type

Extranodal NK/T-cell lymphoma, nasal type is a rare type of non-Hodgkin lymphoma (also known as sinonasal angiocentric T-cell lymphoma). NK/T cell lymphoma and NK/T-cell proliferative disease most often occur in adults, are more common in males than females, are most prevalent in Asia, South America, Central America and Mexico, and represent 7% to 40% of all non-Hodgkin lymphomas, with a 5-year survival near 50% (Suwivat *et al.* 2007, Lee *et al.* 2006, Chan *et al.* 2001).

NK/T-cell lymphoma is almost universally associated with EBV in tumor cells, irrespective of the ethnic origin of the patients; however nasal type NK/T-cell lymphomas presenting in other organ locations have been most strongly associated with EBV in Asian patients (IARC 2012, Chan *et al.* 2001). In recent case series studies in Asia and South America, the presence of EBV

DNA has been identified in nearly 100% of nasal type NK/T-cell lymphoma tumor cells (Barrionuevo *et al.* 2007, He *et al.* 2007). Overall, over 400 EBV-associated NK/T-cell lymphoma cases have been identified including both these studies and studies identified by IARC (1997, see Table 11, sinonasal angiocentric T-cell lymphoma). No co-factors have been identified.

Two case-control studies (that included 10 cases) were identified that have found a positive association between EBV (DNA in plasma or cells) and extranodal NK/T-cell lymphoma, nasal type. However, the temporal relationship between EBV and nasal extranodal NK/T-cell lymphoma has not been established. These studies are reviewed in Table 3-2.

Table 3-2. Summary of case-comparison studies of NK/T cell lymphoma, nasal type and Epstein-Barr Virus

Reference	n/N exposed cases	OR	Comments
Case-Comparison Studies			
Lei <i>et al.</i> 2000 (Hong Kong)	4/4	EBV DNA - ∞ ^a	No exposed controls
Suwivat <i>et al.</i> 2007 (Thailand)	6/6	EBV DNA - ∞ ^a	No exposed controls
	6/6	CD3 ⁺ Cells - ∞	No exposed controls
	6/6	CD3 ⁻ Cells - ∞	19/45 exposed controls, no unexposed cases.

^a EBV plasma DNA.

3.6 Cancer evaluation: Nasopharyngeal carcinoma

3.6.1 Background information

Nasopharyngeal carcinoma is the predominant type of cancer in the nasopharynx. It is a rare cancer in most parts of the world. In the United States, incidence ranges from 0.5 to 2 per 100,000, with a 5-year survival of 36% to over 60% (Lee and Ko 2005). However, in areas like Southeast Asia, the Arctic, North Africa and the Middle East, incidence rates can be much higher, ranging from 2.7 per 100,000 to 26 per 100,000 (Ferlay *et al.* 2015, Chin *et al.* 2014, Yu and Yuan 2002). Additionally, nasopharyngeal carcinoma is much more common in males than females throughout the world (Chin *et al.* 2014, Chang and Adami 2006). Nasopharyngeal carcinomas are classified into three types: keratinizing squamous-cell carcinoma, non-keratinizing carcinoma, and basaloid squamous-cell carcinoma (Chan *et al.* 2005, as cited in IARC 2012). Nasopharyngeal carcinoma is associated with EBV, especially in EBV-endemic populations, with nearly universal EBV seropositivity among cases (Adham *et al.* 2012).

3.6.2 Case-control studies

Fifteen case-control studies (1976 to 2014) have been identified that investigated the association between nasopharyngeal carcinoma and EBV. These 15 studies include over 1,900 cases of nasopharyngeal carcinoma, mostly in Southeast Asia, although 3 studies included cases from Europe, North Africa, and the United States. These studies may be divided into those that investigated the relationship between nasopharyngeal carcinoma and EBV serology, and those that investigated EBV DNA in the tumor.

Eleven of the fifteen case-control studies on the association between nasopharyngeal carcinoma and EBV were serological studies. The majority of these studies generally found a strong and statistically significant association between nasopharyngeal carcinoma and EBV, using a variety of serological markers and tumor DNA analyses. These studies are summarized in Figure 3-5. The ORs ranged from 21 to ∞ , although when studies with no exposure variability, i.e., studies with no non-exposed cases or no exposed controls, were removed, the ORs ranged from 21 to 138, all were statistically significant. These studies are summarized in Figure 3-5. Three studies did not provide enough information to calculate ORs; however, the geometric mean titers of anti-EBV antibodies were statistically significantly higher in cases than in controls (Lennette *et al.* 1995, de Thé *et al.* 1978, Hilgers and Hilgers 1976). Another study, Tiwawech *et al.* (2008) reported on 75 nasopharyngeal carcinoma cases in Thailand and 44 matched controls. They found all cases and controls were exposed to EBNA-2 and LMP-1; however, they found the LMP-1 deletion type subtype to be more common in cases than in controls (OR = 2.5; 95% CI = 1.1 to 5.8). The six case-control studies of EBV DNA in nasopharyngeal carcinoma tumors reported statistically significant increases in EBV DNA with ORs ranging from 86 to ∞ , or 86 to 820 when studies with no exposure variability are removed.




Figure 3-5. Forest plot of serological case-control studies of nasopharyngeal carcinoma and Epstein-Barr virus

CI = confidence interval; EA = early antigen; EBNA = EBV nuclear antigen; IgG/A/M = immunoglobulin G/A/M; OR = odds ratio; VCA = viral capsid antigen.

^aORs or RRs and 95% CIs in curly brackets {} were calculated by IARC (2012, 1997) working groups; ORs or RRs and 95% CIs in square brackets [] were calculated by NTP.

3.6.3 Cohort studies

Two nested case-control studies and two cohort studies were identified that investigated the association between nasopharyngeal carcinoma and EBV. The two nested case-control studies included a total of 14 nasopharyngeal carcinoma cases, with one study of 7 cases among Alaska Natives (Lanier *et al.* 1980) and the other among a general population in the United States (Chan *et al.* 1991). Both investigated whether EBV antibodies were present prior to a nasopharyngeal carcinoma diagnosis, with follow-up times ranging from 1 to 12 years. Neither study found a statistically significant relationship between EBV antibodies prior to diagnosis and nasopharyngeal carcinoma. The two prospective cohort studies (Ji *et al.* 2007, Chien *et al.* 2001) had over 51,000 participants and a total of 168 cases of nasopharyngeal carcinoma, with follow up times up to 16 years. Both studies reported statistically significant associations and nasopharyngeal carcinoma incidence between those designated sero-positive and sero-negative at baseline, with relative risks of 22 (95% CI = 7.3 to 66.9) and 9.4 (95% CI = 6.7 to 13.3), respectively. Chien *et al.* also found a statistically significant relationship between EBV DNase and nasopharyngeal carcinoma with a relative risk of 3.5 (95% CI = 1.4 to 8.7). These studies are summarized in Figure 3-5.

3.6.4 Cofactors

Two potential cofactors in the association between EBV and nasopharyngeal carcinoma were reviewed by IARC (1997): dietary factors (such as Cantonese-style salted fish, other preserved foods, and deficits of fruits and vegetables), and genetic factors. These cofactors are reviewed briefly here. An in-depth review is available in IARC (1997).

No new studies on dietary factors as cofactors in the association between EBV and nasopharyngeal carcinoma have been identified since the previous IARC reviews (2012, 1997). Earlier case-control studies found the consumption of Cantonese-style salted fish, particularly during childhood, to be associated with a diagnosis of nasopharyngeal carcinoma, with ORs ranging from 1.5 to 38, which were generally statistically significant (Zheng *et al.* 1994b, Zheng *et al.* 1994a, Sriamporn *et al.* 1992, Ning *et al.* 1990, Yu *et al.* 1989, Yu *et al.* 1988, Yu *et al.* 1986, Armstrong and Armstrong 1983, Geser *et al.* 1978, Henderson and Louie 1978). Those studies that did not find a statistically significant association did not include childhood consumption in their analysis (Lee *et al.* 1994, Chen *et al.* 1988). Other types of salted fish were not significantly associated with nasopharyngeal carcinoma; however, other preserved foods, typically salted, were associated with nasopharyngeal carcinoma in several case-control studies, with ORs ranging from 1.2 to 8.6. The results of studies investigating the relationship between deficits of fresh fruits and vegetables and nasopharyngeal carcinoma were mixed, with some finding no association and others finding statistically significantly lower levels of certain vitamins among nasopharyngeal carcinoma patients. Although none of the studies evaluated the relationship between EBV and dietary factors, they may be potential co-factors as mechanistic studies (reviewed by IARC 1997) found that Cantonese-style salted fish, along with other preservatives, contained substances capable of activating EBV in latently infected cells (Poirier *et al.* 1989, Shao *et al.* 1988, as cited in IARC 1997).

Genetic factors have also been suggested as cofactors for nasopharyngeal carcinoma, with multiple susceptible genetic loci identified. EBV interacts with the host cell genes implicated in nasopharyngeal carcinoma development, influencing cell signaling and host gene regulation, and

predisposing the cell for nasopharyngeal carcinoma (Lung *et al.* 2014, Aldred and Eng 2006, as cited in Lung *et al.* 2014). No epidemiological studies on the associations between nasopharyngeal carcinoma, EBV, and genetic factors have been identified; however, five new studies since 2008 have demonstrated that genetic variation in certain genes is associated with the risk of nasopharyngeal carcinoma. Two of these studies (Chin *et al.* 2014, Zhao *et al.* 2012c) found increased susceptibility for nasopharyngeal carcinoma associated with the HLA-A locus in Chinese cases of nasopharyngeal carcinoma. This is consistent with previous research that has found an increased risk of nasopharyngeal carcinoma associated with the HLA-A and HLA-B loci among Chinese populations. These previous studies were reviewed in IARC 1997, along with studies in non-Chinese populations. Two new case-control studies investigated the association between nasopharyngeal carcinoma and the genes for the cytokines interleukin-2 (Wei *et al.* 2010) and interleukin-18 (Nong *et al.* 2009) among Chinese populations. Significant differences were found between the genotypes and allele frequencies of these cytokines between cases and controls in both studies. Moumad *et al.* (2013) studied 492 nasopharyngeal carcinoma patients in North Africa, and found significant associations between nasopharyngeal carcinoma and polymorphisms of genes of pattern recognition receptors (Toll-like receptor TLR3, C-type lectin receptor CD209, retinoic acid-inducible gen I-like receptor [DDX58]).

3.6.5 Integration of the evidence

The epidemiological data, primarily a large body of case-control studies, consistently report statistically significant, positive associations between EBV seropositivity and nasopharyngeal carcinoma. A number of these studies were conducted among populations with a high prevalence of nasopharyngeal carcinoma. ORs of 20 or higher were seen for most case-control studies, using both serological and tumor DNA EBV detection methods. Two cohort studies have also shown positive associations between EBV seropositivity and nasopharyngeal carcinoma incidence and demonstrated the temporality of the relationship, with EBV seropositivity preceding development and diagnosis of nasopharyngeal carcinoma. Long follow-up times between study enrollment (when EBV seropositivity is determined) and diagnosis may lead to some misclassification, with those developing EBV seropositivity after enrollment but before diagnosis being misclassified as seronegative. Two cofactors, dietary factors and genetic factors, have been suggested to have an influence on the relationship between EBV and nasopharyngeal carcinoma.

3.7 Cancer evaluation: Gastric cancer

3.7.1 Background information on gastric cancer

Gastric cancer is one of the most common cancers worldwide, with 7.5 cases per 100,000 people diagnosed each year in the United States. The 5-year survival rate for gastric cancer is 20% or less (Crew and Neugut 2006). Epstein-Barr virus was first detected (via polymerase chain reaction) in a patient with gastric cancer in 1990 (Burke *et al.* 1990).

3.7.2 Case-series analyses and pooled analyses

EBV-related gastric cancer is defined as detection of EBV in the gastric cancer tumor, through a variety of detection methods, such as *in situ* hybridization or polymerase chain reaction. A case series of 138 U.S. patients (Shibata and Weiss 1992, as cited in IARC 1997) found the

prevalence of EBV-related gastric cancer to be 16%. In a systematic review of 47 case series and case-control studies, Chen *et al.* (2015) found EBV DNA in 5% to 18% of gastric cancer cases when detected by *in situ* hybridization. A pooled analysis of case reports and cases series found the global prevalence of EBV-related gastric cancer to be around 8.3% to 8.7% (Murphy *et al.* 2009, Sousa *et al.* 2008). These analyses also found the highest prevalence of EBV-related gastric cancer in North and South America (13%) and the lowest in Southeast Asia (7.8%) (Sousa *et al.* 2008). Additionally, in their meta-analysis based on 15,952 cases of gastric cancer worldwide, Murphy *et al.* (2009) noted a two-fold difference in EBV-related gastric cancer by sex, with a prevalence of 11.1% in males and 5.2% in females. Pooling studies with various EBV detection methodologies, this meta-analysis also identified differences in EBV-related gastric cancer prevalence based on anatomic location of the gastric cancer tumor. Tumors originating in the gastric cardia or corpus (body) were twice as likely to be EBV positive compared with tumors in the pyloric antrum, and EBV prevalence was 4 times higher for tumors arising in postsurgical gastric stump/remnants compared to the pooled prevalence of EBV positivity. Additionally, over 90% of lymphoepithelioma-like carcinomas of the stomach were EBV positive.

3.7.3 Case-control studies and case-case analyses

There have been three case-control studies and one case-case comparison study published that investigated the association between EBV and gastric cancer (de Aquino *et al.* 2012, Lo *et al.* 2001, Shinkura *et al.* 2000) (Table 3-3). Among the three case-control studies, there were a total of 77 EBV-positive gastric cancer cases out of 184 total gastric cancer cases. Two of these studies were reviewed by IARC (2012). In a population of 123 gastric cancer cases (64 cases with EBV-positive tumors) and 73 healthy controls in Japan, Shinkura *et al.* (2000) calculated the EBV seroprevalence of cases compared with controls. Cases were 7.1 (95% CI = 2.4 to 25.6) times more likely to be VCA-IgA positive and 19.9 (95% CI = 6.6 to 70) times more likely to be EA-IgG positive than healthy controls (ORs calculated by IARC working group, available in IARC 2012, [Table 2.9](#)). The authors also conducted a case-case comparison between patients with EBV-positive and EBV-negative tumors and found the geometric mean of EBV VCA-IgG ($P < 0.001$), EBV VCA-IgA ($P = 0.006$), and EA-IgG ($P < 0.001$) to be significantly higher among EBV-positive cases compared to EBV-negative cases; however, there was no difference in EBNA antibodies. EBV-positive cases were 3.4 (95% CI = 1.3 to 8.8) and 6.6 (95% CI = 2.7 to 16.3) times more likely to be seropositive for VCA-IgA and EA-IgG, respectively, compared to EBV-negative cases. It is noteworthy, however, that cases with EBV-negative tumors were 4.9 times more likely to be seropositive for EA-IgG than healthy controls, and the geometric mean titer of VCA-IgG was significantly higher for EBV-negative cases than in healthy controls.

Table 3-3. Summary of case-control studies of gastric cancer and Epstein-Barr Virus

Reference	OR	Comments
Case-Control Studies		
Shinkura <i>et al.</i> 2000	VCA-IgA: {7.1} ^a EA-IgG: {19.9} ^a	Lower 95% CI > 1.0 Lower 95% CI > 1.0
Lo <i>et al.</i> 2001	EBV DNA (EBV+ tumor): ∞	5/5 cases EBV+, 7/197 controls EBV+
de Aquino <i>et al.</i> 2012	EBV DNA (EBV – tumor, EBV + lymphocytes): {352.9}	Lower CI > 1.0
	EBV DNA: ∞	8/10 cases EBV+. 0/6 control EBV+

^aORs in curly brackets {} were calculated by the IARC Working Group (2012).

CI = confidence interval; EA = early antigen; IgG/A = immunoglobulin G/A; OR = odds ratio; VCA = viral capsid antigen.

Lo *et al.* (2001) investigated serum EBV DNA in 51 gastric cancer cases (5 patients EBV-encoded small RNA [EBER]-positive, and 14 EBER-negative) in Hong Kong compared with 197 healthy controls. They found that EBV DNA was detectable in all EBV-encoded small RNA (EBER)-positive cases (5/5) and in all but one EBER-negative case (13/14), compared to 3.6% of healthy controls.

One new case-control study has been published since 2008 that investigated the prevalence of EBV DNA in the tumors of 10 gastric cancer patients in Brazil compared to a convenience sample of biopsies from 6 cancer-free control subjects obtained during endoscopies (de Aquino *et al.* 2012). EBV DNA, detected by polymerase chain reaction (PCR), was present in 8 of 10 gastric cancer cases and in none of the control subjects.

Boysen *et al.* (2011) conducted a nationwide case-case comparison study in Denmark which included 18 EBER-positive EBV gastric cancer cases out of 186 total gastric cancer cases, both with and without pernicious anemia. In comparison with gastric cancer patients without pernicious anemia, gastric cancer patients with pernicious anemia (M = 8) were 2.5 (95% CI = 0.88 to 7.14) times more likely to be EBV positive when adjusting for gender, age at diagnosis, and year of diagnosis; when further adjusted for lymphocytic infiltration, those with pernicious anemia were 2.9 (95% CI = 0.99 to 8.67) times more likely to be EBV positive.

3.7.4 Nested case-control studies

Levine *et al.* (1995) examined serum samples collected and banked prior to the diagnosis of 54 cases of gastric adenocarcinoma patients of Japanese ancestry selected from the Honolulu Heart Cohort study, along with 54 controls matched on age and date of blood collection. This study found a non-statistically significant increased risk between EBV seropositivity and subsequent development of gastric adenocarcinoma (Table 3-3).

In another nested case-control study (Koshiol *et al.* 2007) among 185 cases of gastric cancer and 200 controls in China, EBV seropositivity prior to diagnosis was found to be unrelated to the incidence of gastric cancer (ORs less than 1). Follow-up time between enrollment and diagnosis

was as long as fifteen years. The study also found no difference in EBV seropositivity between cardia and non-cardia gastric cancer cases (see Table 3-4). However, this study did find that cardia gastric cancer cases with high baseline EBNA IgG titers (prior to diagnosis) had longer survival (hazard ratio = 0.46, 95% CI = 0.29 to 0.74) than either cardia gastric cancer cases with low baseline EBNA titers or all non-cardia gastric cancer cases, although no interpretation for this finding was given.

Table 3-4. Summary of cohort and nested case-control studies of gastric cancer and Epstein-Barr Virus

Reference	OR	Comments ^b
Nested Case-Control Studies		
Levine <i>et al.</i> 1995	VCA IgG: 1.4 VCA IgA: 3.9 EBNA: 0.72 EA(D): 1.2 EA(R): 1.9	NS – High (1280+) versus low (\leq 640) titers NS – High (20+) versus low (\leq 20) titers NS – High (640+) versus low (\leq 320) titers NS – High (5+) versus low (\leq 5) titers NS – High (5+) versus low (\leq 5) titers
Koshiol <i>et al.</i> 2007	EBNA: 0.46 ^a VCA IgA: 0.69 EA-D IgG: 0.95 EA-R IgG: 0.52 High EBNA: 0.91	Upper CI < 1. Survival among individuals with high vs. low EBNA. Seropositivity and development of gastric cancer. All non-significant.
Kim <i>et al.</i> 2009b	VCA IgG: 1.37 EBNA IgG: 0.90	NS. OR of gastric cancer risk of highest titer levels compared to lowest. NS. OR of gastric cancer risk of highest titer levels compared to lowest.

^aHazard Ratio reported.

CI = confidence interval; EA = early antigen; EA-D = early antigen-diffuse; EA-R = early antigen-restricted; EBNA = EBV nuclear antigen; IgG/A = immunoglobulin G/A; NS = not statistically significant; OR = odds ratio; RR = relative risk; VCA = viral capsid antigen.

In a third nested case-control study based in South Korea, Kim *et al.* (2009b) studied 100 incident gastric cancer cases from a cohort of 14,000 participants in a multi-center cohort with 200 controls matched on year of enrollment, age, gender, and area of residence. Follow-up time between enrollment and gastric cancer diagnosis ranged from 0 to 9 years. The OR between EBV antibody levels prior to diagnosis and subsequent risk of gastric cancer, including among patients, was 1.37 (95% CI = 0.62 to 3.06) with the highest titers of VCA IgG and 0.87 (95% CI = 0.51 to 1.46) for EBNA IgG, but no dose-response relationship was observed (Table 3-4).

3.7.5 Cofactors

Currently, there are no identified cofactors for the association between EBV and gastric cancer.

3.7.6 Integration of the evidence

The data from the three case-control studies suggest a positive association between EBV and gastric cancer. However, in two of those studies there was little exposure variability. In one study, all cases were EBV-positive, while in another there were no exposed controls. This lack of

variability may account for heterogeneity seen in the results. In the three nested case-control studies, EBV titer levels prior to diagnosis were not significantly associated with an increased risk of gastric cancer; however, two studies found non-significant associations for VCA and EA antigens (Kim *et al.* 2009b, Levine *et al.* 1995). None of the prospective studies typed the tumor DNA for EBV; therefore, it is unknown if those with EBV seropositivity prior to diagnosis were true EBV-positive gastric cancer cases. There were also long follow-up times between enrollment and diagnosis for some cases resulting in non-differential exposure misclassification. Those who developed EBV seropositivity inbetween enrollment and diagnosis would have been missed and thus would bias the findings towards the null.

3.8 Cancer evaluation: Lymphoepithelial carcinoma of the salivary gland

Lymphoepithelial carcinoma of the salivary gland is a rare carcinoma histologically similar to nasopharyngeal carcinoma, accounting for less than 1% of all head and neck cancers (Tsang and Chan 2005, Tsai *et al.* 1996). In populations where nasopharyngeal carcinoma is common, such as Southeast China and Greenland, lymphoepithelial carcinoma of the salivary gland has shown a positive association with EBV (Wang *et al.* 2004). As reported by the IARC (1997) review of five case reports and case series published between 1982 and 1996, 25 out of 27 total salivary gland lymphoepithelial carcinoma cases tested positive for EBV in tumor cells (IARC 1997). Additional case-series studies of lymphoepithelial carcinoma of the parotid gland were reported in the later IARC (2012) review with 208 of 209 cases of parotid gland lymphoepithelial carcinoma reporting EBV DNA in the tumor.

3.8.1 Case-case study

Only one case-case study of salivary gland lymphoepithelial carcinoma has been identified in the published literature. Wang *et al.* (2004) compared 16 cases of salivary gland lymphoepithelial carcinoma to 12 cases of other types of salivary gland tumors in Taiwan between 1977 and 2001. EBV DNA was present in the tumors of all 16 salivary gland lymphoepithelial carcinoma cases and in none of the other types of salivary gland tumors. Cases of lymphoepithelial carcinoma (all of which were EBV positive) had a better prognosis than cases of other tumor types. The lymphoepithelial carcinoma patients had a 5-year survival rate of 86% with treatment. Patients with other salivary gland tumor types who underwent treatment had a 5-year survival of only 36%.

3.8.2 Cofactors

No cofactors have been identified.

3.8.3 Integration of the evidence

The number of epidemiological studies on the association between EBV and lymphoepithelial carcinoma of the salivary gland is insufficient to evaluate this endpoint. Although there is evidence of a positive association between EBV and salivary gland lymphoepithelial carcinoma in several case reports and series, no case-control, cohort, or nested case-control studies have been identified to date.

3.9 Synthesis across cancer endpoints

A summary of the evidence for EBV infection and the different cancer endpoints from epidemiological studies is provided in Table 3-5. The preliminary level of evidence from cancer studies in humans also considers studies of tissues from humans in addition to epidemiological studies and is provided in Section 5.

Table 3-5. Summary of EBV cancer endpoints and strength of the epidemiological evidence

Cancer endpoint	Strength of evidence
Burkitt lymphoma (endemic)	<ul style="list-style-type: none"> Consistent evidence across multiple studies. All epidemiological studies report significant associations. Elevated pre-diagnosis titers seen to one EBV antigen (VCA) in one prospective study. Dose-response relationships observed in several studies.
(sporadic)	<ul style="list-style-type: none"> Generally elevated, but non-significant ORs. One study found a significant association. No prospective studies identified.
Hodgkin lymphoma	<ul style="list-style-type: none"> Consistent evidence among EBV serology studies, with only a few non-significant ORs. EBV DNA studies show a very strong association with tumor DNA. One EBV serology nested case-control studies shows temporal relationship. Evidence of infectious mononucleosis and Hodgkin lymphoma association less clear.
Immunosuppression-related non-Hodgkin lymphoma	<ul style="list-style-type: none"> Case series indicate EBV is consistently found in cases of immunosuppression related non-Hodgkin lymphoma. Two epidemiological studies, one prospective, found evidence of an association, though not statistically significant.
Extranodal NK/T cell lymphoma, nasal type	<ul style="list-style-type: none"> Consistent association of nasal NK/T-cell lymphoma with EBV in tumor cells in case series; over 400 cases. Two case-comparison studies found EBV DNA in the plasma or CD3+ (T cells) cells from cases but not from controls
Nasopharyngeal carcinoma	<ul style="list-style-type: none"> Consistent evidence of a strong association between EBV and NPC. Most ORs were 20 or higher. All case-control ORs were significant. Two cohort studies showed a temporal association between EBV and NPC.
Gastric cancer	<ul style="list-style-type: none"> Three case-control studies found strong associations with both EBV serology and DNA. Elevated but non-statistically significant increased risks were found in two of three serological nested case-control studies.
LEC of the salivary gland	<ul style="list-style-type: none"> EBV DNA was detected in tumors in almost all of the more than 200 cases of parotid salivary gland lymphoepithelial carcinoma No epidemiological studies were available for review.

LEC = lymphoepithelial carcinoma; NPC = nasopharyngeal carcinoma; NHL = non-Hodgkin lymphoma.

4 Mechanisms and Other Relevant Data

Epstein-Barr virus (EBV) was the first oncogenic virus identified in humans and was found to be associated with endemic Burkitt lymphoma over fifty years ago (Epstein *et al.* 1964). Evidence of the oncogenic potential of EBV was demonstrated by its ability to transform human B lymphocytes in cell culture and by studies in non-human primates (Shope *et al.* 1973, Pope *et al.* 1968). Over 90% of the world's adult population is infected with EBV by age 20 and, for the most part, the result is an asymptomatic life-long infection—similar to other herpesviruses—and is held in check by immune surveillance. Several patterns of gene expression during virus latency have been shown to be associated with EBV pathogenesis of some cancers (see Table 1-1). From studies of human cancer populations and refinement of molecular techniques, several types of cancer have a clear causal association with this virus such as Burkitt lymphoma, Hodgkin lymphoma, and nasopharyngeal carcinoma. However, the presence of EBV and its possible role in other cancers, such as carcinoma of the lung, skin, or various glandular tissues, is not as well understood. Investigation of the properties of the virus, as well as the cofactors and mechanisms that enable cancer formation or are protective, is on-going.

This section provides a brief review of the characteristics of EBV-associated neoplasms (Section 4.1), the roles of viral gene transcripts in malignant transformation (Section 4.2), the mode of action and evidence for cancer causation (Section 4.3), and a synthesis of this information (Section 4.4).

4.1 General characteristics

As discussed in the previous sections, seven neoplasms—four in lymphoid tissue and three in epithelial tissue—have been primarily associated with EBV in humans, although the level of evidence for each tumor varies (IARC 2012). These neoplasms include: (1) Burkitt lymphoma (2) Hodgkin lymphoma, (3) immunosuppression-related non-Hodgkin lymphoma, 4) extranodal NK/T-cell lymphoma, nasal type, 5) nasopharyngeal cancer, 6) gastric cancer, and 7) lymphoepithelial-like carcinoma of the salivary gland. General characteristics of these EBV-associated neoplasms and degree of association of the neoplasm with EBV, i.e., percentage of tumor cells containing the EBV genome are listed in Table 4-1.

Table 4-1. Characteristics of EBV-associated neoplasms

Neoplasm	Clinical presentation	Lineage and primary tumor cell; evidence of EBV clonality ^a	Percent of tumors or tumor cells EBV positive; clonality
Burkitt lymphoma	<i>Endemic form</i> - Extra nodal lymphoid tissue (jaw, kidney, bowel, adrenal gland); positive malaria titer; children <i>Sporadic form</i> - Abdomen, lymph nodes; adults and children	Neoplastic B cells with <i>c-myc</i> translocation to immunoglobulin locus- all forms of Burkitt lymphoma; monoclonal	<i>Endemic form</i> - 95% of tumor cells contain EBV genome <i>Sporadic form</i> - at most 20% of tumors contain EBV genome

Neoplasm	Clinical presentation	Lineage and primary tumor cell; evidence of EBV clonality ^a	Percent of tumors or tumor cells EBV positive; clonality
	<i>Immunodeficiency associated form</i> - anti-rejection therapy; early latency with HIV infection; also congenital cause		<i>Immunodeficiency associated form</i> - 40% of lymphomas in HIV-positive patients (Gloghini <i>et al.</i> 2013, Stefan <i>et al.</i> 2011)
Hodgkin lymphoma	Nodal tissue of neck, mediastinal, axillary and paraortic regions; lymphadenopathy with rich inflammatory background	Multinucleated, clonal Hodgkin Reed-Sternberg cells (B cell origin; CD15, CD30 markers)	EBV primarily associated with 75% of mixed-cellularity and 20% of nodular sclerosis subtypes of classical Hodgkin lymphoma; Hodgkin lymphomas in HIV positive patients are 100% EBV associated
Immunosuppression-related NHL	Lymph nodes, gastrointestinal tract, lungs, liver; associated with CNS lymphomas in HIV positive patients; Lymphomas may occur with anti-rejection therapy	Primarily (85%) B-cell origin, monoclonal and polyclonal forms	100% of CNS lymphomas in HIV positive patients; Systemic- 90% diffuse large cell lymphomas, 40% small non-cleaved cell lymphadenopathies (Burkitt lymphomas) EBV associated; 60% EBV associated in post transplant lymphoproliferative disease
Extra nodal NK/T-cell lymphoma	Lymphoid tissue; extra nodal; diffuse lymphocytic infiltrate and vascular damage	NK and T cells; most cases are NK cell neoplasms, some have a cytotoxic T-cell phenotype; monoclonal	100% in nasal variant
NPC	Epithelial tissue; strong ethnic and geographic association: Inuit, Southern Asia, China; keratinizing and non-keratinizing forms	Nasal epithelial cells; monoclonal	98% in non-keratinizing carcinoma; clonal detection in precancers (Tsao <i>et al.</i> 2015, Tsang <i>et al.</i> 2014, Pathmanathan <i>et al.</i> 1995)
Gastric cancer	Epithelial tissue; 89% with lymphocytic infiltration; proximal stomach or associated with stomach remnant	Gastric epithelial cells; monoclonal	9% of gastric cancers

Neoplasm	Clinical presentation	Lineage and primary tumor cell; evidence of EBV clonality ^a	Percent of tumors or tumor cells EBV positive; clonality
LEC of the salivary gland	Epithelial tissue and non-cancerous lymphoid cells	Salivary gland epithelial cells; monoclonal	Primarily parotid salivary gland

Sources: Information from IARC (2012, 1997) unless cited in the table.

^aEvidence of clonality of virus in tumor tissue.

LEC = lymphoepithelial cancer; NPC = nasopharyngeal carcinoma.

4.2 EBV latent genes and malignant transformation

EBV in its latent phase can express different transcription programs (0, I, II, III). In an immunocompetent host, EBV persists in memory B cells as a latent infection (latency 0). Since no proteins are produced during latency 0, the presence of the virus is not recognized by the immune system, resulting in lifetime infection. EBV-encoded small RNAs (EBERs) and microRNAs are expressed in all latency phases and approximately 22 microRNAs have been identified (IARC 2012). EBERs and microRNAs as well as the full complement of latent gene proteins are expressed in latency III. Latent gene proteins consist of six nuclear antigens (EBNAs), and three latent membrane proteins (LMPs) (Yau *et al.* 2014, IARC 2012). EBV transcripts include coding and non-coding RNAs and proteins produced during latency phases (see Table 1-1). Primarily latency II, but also latency I and III phases, promote acquisition of various cancer hallmarks, e.g., insensitivity to antigrowth signals, avoiding cell-cycle arrest, immune evasion, genetic instability, cell proliferation, resisting apoptosis, promotion of angiogenesis, and induction of genomic instability (Mesri *et al.* 2014). Table 1-1 relates their expression pattern to that found in various neoplasias. The general functions of key viral proteins and transcripts important in the pathogenesis of EBV-associated cancers are described in Table 4-2.

Table 4-2. Activation of oncogenic pathways by viral genes

EBV Transcription program and gene products	Some host pathways affected	Potential cancer property	Associated malignancies ^a
Latency 0: EBERs		Promote growth, anti apoptotic	None
Latency I: EBNA-1, EBERs	Regulation of RAG-1 and RAG-2 by EBNA-1	Resisting cell death, avoiding immune destruction, genome instability, increase in reactive oxygen species	Burkitt lymphoma and gastric carcinoma ^b
Latency II: EBNA-1, LMP-1, LMP-2A, EBERs	NFkB, JNK by LMP-1	Resisting cell death, enabling replicative immortality, angiogenesis, inflammation	NPC, Hodgkin lymphoma, NK and T cell lymphoma, nasal type ^c , LEC of the salivary gland
Latency III: EBNA-1, -2, -3A, -3B, -3C, -LP, LMP-1, LMP-2A,	PI3K-Akt-mTOR, ERK by LMP-2A	Inducing angiogenesis, sustaining proliferative signaling, deregulation of cellular pathways, activation of invasion and metastasis, enables	Immunosuppression-related NHL (AIDS-associated, post-transplant disorder, iatrogenic)

EBV Transcription program and gene products	Some host pathways affected	Potential cancer property	Associated malignancies ^a
LMP-2B, EBERs		replicative immortality, activation of host methyltransferase	
MicroRNAs ^d	Repression of translation	Immune evasion	

Sources: Grywalska and Rolinski 2015, Mesri *et al.* 2014, IARC 2012

EBER = EBV-encoded small RNA; EBNA = EBV nuclear antigen; LMP = latent membrane protein; EBNA-LP = EBV nuclear antigen leader protein; NHL = non-Hodgkin lymphoma; LEC = lymphoepithelial cancer; NPC = nasopharyngeal carcinoma.

^aGeneral classification of disease and heterogeneous patterns may be present in different cell populations in same individual.

^bExpression of lytic genes and LMP-2A variable.

^cPreviously called sinonasal angiocentric T cell lymphoma.

^dIn latency I, II, III microRNAs (at least 22) are expressed in various amounts.

4.3 Mode of action and evidence for cancer causation

Direct evidence for causality of EBV in lymphomagenesis comes from *in vitro* studies and studies in mice. EBV has been shown to transform lymphoblastoid cells in culture and can transform epithelial cells when co-cultured with transformed lymphoblastoid cells (Imai *et al.* 1998). In addition, lymphoblastoid cells transfected with activated *c-myc* genes were tumorigenic in nude mice, and infected B cells have been shown to cause B-cell lymphomas in SCID mice (Rowe *et al.* 1991, Mosier *et al.* 1989, Lombardi *et al.* 1987). Further, EBV proteins, EBNA-1, -2, -3A, -3C, and LMP-1 are all necessary for immortalization of B-lymphocytes (Grywalska and Rolinski 2015). Clearly, EBV has oncogenic potential to transform lymphoid and epithelial cells in culture and has been found associated with cancers of epithelial and lymphoid origin. EBV is a ubiquitous virus, and criteria have been outlined by IARC as well as discussed by others for judging cancer causality by viruses (zur Hausen 2001, IARC 1997) (see Overview and Introduction Section). The key causality criteria included the following: (1) the presence of the virus in the tumor, (2) monoclonality of EBV in the tumor (suggesting the presence of latent infection prior to expansion of the malignant clone), (3) and expression of viral proteins in pre-neoplastic lesions and in malignant tissue (IARC 1997). The epidemiological and experimental data show that EBV meets these criteria. The mode(s) of action and evidence linking EBV with cancer are briefly reviewed by tumor type in the following sections.

4.3.1 Burkitt lymphoma

Burkitt lymphoma is a B-cell non-Hodgkin lymphoma that presents mostly as an extranodal mass, although nodal involvement is seen in sporadic cases. In normal lymphoid tissue, B-cell response to antigenic stimulation produces B-cytoplasts that differentiate into plasma cells that produce antibodies. In Burkitt lymphoma, B-cell differentiation in response to antigen stimulus undergoes faulty DNA recombination in the lymphoid tissue germinal center resulting in immunoglobulin gene/*c-myc* translocation. This results in constitutive expression of the *c-myc* oncogene driven by immunoglobulin promoters and is the primary lesion in Burkitt lymphoma. From human cancer studies, it is known that endemic Burkitt lymphoma is associated with chronic antigenic stimulation associated with malarial infection, and B cells infected with EBV have latency I EBV gene expression pattern. EBV and malarial infections can promote B-cell hyperplasia and increase the chances that B-cell translocations will occur. Further, malarial infection has been shown to directly increase the number of germinal center

translocations occurring via deregulation of activation-induced cytidine deaminase, an enzyme responsible for class switch recombination and somatic hypermutation, in the germinal center (Torgbor *et al.* 2014, Robbianai *et al.* 2015). Normally, defective cells exiting the germinal center undergo apoptosis; however, it has been shown that EBV expression of EBNA-1 rescues defective B cells from removal by apoptosis by directly upregulating survivin production by complexing with SP1 on the *survivin* gene promoter (Lu *et al.* 2011, Kennedy *et al.* 2003). Therefore, potentially more Burkitt lymphoma cells would form in the germinal center with malarial infection, and EBV cells expressing EBNA-1 would be the ones that survive.

Other lines of evidence linking EBV and Burkitt lymphoma include the following:

- EBV is associated with all three clinical forms of Burkitt lymphoma (endemic, sporadic, immune suppression related) (IARC 2012, 1997).
- EBV DNA is present in Burkitt lymphoma cells in monoclonal form (IARC 1997).
- Both malaria and EBV cause B-cell hyperplasia, and increase in B cells could increase the probability of a faulty gene translocation (Torgbor *et al.* 2014, Robbiani *et al.* 2015).
- EBNA-1 prevents apoptosis of Burkitt lymphoma cells and enables cell survival (Lu *et al.* 2011, Kennedy *et al.* 2003).

4.3.2 Hodgkin lymphoma

Hodgkin lymphoma is a group of lymphoid neoplasms arising from a single lymph node, spreading to contiguous lymph nodes, and containing Hodgkin Reed-Sternberg cells, which are giant multinucleated neoplastic cells of clonal B-cell origin (Stein *et al.* 2008). Over 90% of the tumor is characterized by non-malignant inflammatory cells (i.e., reactive lymphocytes, macrophages, granulocytes, and fibrocytes) and fibrosis. Based on histopathology, Hodgkin lymphoma is divided into classical (with four subtypes: mixed cellularity, nodular sclerosis, lymphocyte depleted, and lymphocyte rich) and nodular lymphocyte-predominant Hodgkin lymphoma, an uncommon form that is almost always EBV negative (Stein *et al.* 2008) Hodgkin lymphomas account for approximately 30% of all lymphomas worldwide; however, as noted in Table 4-1, the degree of EBV infection varies among different subtypes of classical Hodgkin lymphoma (IARC 2012, 1997).

Unlike germinal center maturation of B lymphocytes to plasma cells or memory B lymphocytes, Hodgkin Reed-Sternberg cells have not undergone somatic hypermutation and do not express immunoglobulins or most B-cell specific genes, but do reveal deregulation of multiple signaling pathways such as NF- κ B, JAK/STAT, MAP kinase, PI3-kinase/AKT and produce an abnormal pattern of release of cytokines and chemokines resulting in a local inflammatory response. LMP-1 has been shown to be a transforming oncogene in cell culture and promotes growth and survival of Hodgkin Reed-Sternberg cells through upregulation of these pathways (Mohamed *et al.* 2014). EBV expresses type II latency genes (Table 4-2) in Hodgkin Reed-Sternberg cells (IARC 2012, 1997).

Further evidence that supports EBV infection as a causal agent in some forms of Hodgkin lymphoma is as follows (IARC 2012):

- Monoclonal EBV episomes are detected in Hodgkin Reed-Sternberg cells indicating that EBV infection has occurred before clonal expansion.
- Antibody titers to EBV viral capsid antigens are increased with Hodgkin lymphoma.
- The risk of Hodgkin lymphoma increases 4-fold with a previous history of infectious mononucleosis.

4.3.3 Immunosuppression-related non-Hodgkin lymphoma (HIV+ and post-transplant lymphoproliferative disease)

EBV infection in a severely immunocompromised host can lead to immunosuppression-related non-Hodgkin lymphoma. EBV-related non-Hodgkin lymphomas have been linked to immunosuppression therapies related to transplant of organs or cells (stem cells or bone marrow), and to severe immune suppression as a result of HIV infection (Pietersma *et al.* 2008). The link between immune suppression and non-Hodgkin lymphoma caused by EBV has been characterized from studies measuring viral load and EBV-specific cytotoxic T lymphocytes in the blood of patients with severe immune suppression (acquired immune deficiency syndrome, AIDS) or with post transplant lymphoproliferative disorders (IARC 2012, Gulley and Tang 2010, IARC 1997). Post-transplant lymphoproliferative disorder occurs in approximately 0.5% to 1% overall of kidney or bone marrow transplants and varies with the patient age and type of transplant (Vegso *et al.* 2011). Early lesions are often polyclonal and, if untreated, continued B-cell proliferation driven by active EBV will generate a neoplastic clone (Gulley and Tang 2010).

With HIV infection, perturbations in the immune system and loss of lymphocytes decrease immune surveillance. Central nervous system non-Hodgkin lymphoma may result from HIV-related profound immunosuppression; 100% of these lymphomas are associated with EBV (IARC 1997). In addition, there is evidence of clonality of EBV and production of LMP-1, an oncoprotein, by EBV in HIV-associated central nervous system non-Hodgkin lymphoma tissues (IARC 1997). EBV is also associated with a large percentage of systemic AIDS-related non-Hodgkin lymphomas with up to 90% of diffuse large cell lymphomas and in approximately 40% of small non-cleaved cell lymphadenopathies EBV associated (IARC 1997).

Patients receiving transplants but not previously infected with EBV, such as children, are at a greater risk of developing non-Hodgkin lymphoma from infected transplant tissue or from transmission of EBV in saliva from an infected person. Polymorphic post-transplant lymphoproliferative disease appears early after transplant procedure (< 1 year) in children and is associated with a primary EBV infection (Vegso *et al.* 2011). Most of the post-transplant lymphoproliferative disease lesions are of B-cell origin (85%) and > 90% are associated with EBV infection (IARC 2012, Vegso *et al.* 2011, Taylor *et al.* 2005). The most common EBV gene expression pattern with post-transplant patients is latency III growth pattern (Carbone *et al.* 2008, see Table 1.1).

Evidence that EBV is causal in some cases of immunosuppression-related non-Hodgkin lymphoma includes the following (Lim *et al.* 2006):

- Treatment with EBV-specific cytotoxic lymphocytes results in decreases in EBV viral load and in tumor size.

- Further administration of EBV-specific cytotoxic lymphocytes to an immune suppressed individual can protect against lymphoma development.
- In organ or cell transplant cases, decreasing therapy for immunosuppression can decrease viral load and tumor size. Not all cases of post transplant lymphoproliferative disease or lymphoproliferative disease with HIV are related to EBV infection, but reactivation of a latent infection or initiation of a primary EBV infection can lead to lymphoproliferative disease.

4.3.4 Extranodal NK/T-cell lymphoma, nasal type

Extranodal NK/T-cell lymphoma, previously known as angiocentric T-cell lymphoma, is a rare EBV-associated lymphoma that is more prevalent in South America, Mexico, and Asia than in the United States. NK/T-cell lymphomas are classified by location of the tumor to nasal, intestinal, and subcutaneous panniculitis-like lymphomas. The amount of association of these lymphomas with EBV varies with tumor site, geographic location, and genotype of the patient (Ambinder and Cesarman 2007).

EBV expresses type II latency genes (Table 4-2) in this cancer. LMP-1, an EBV-specific oncoprotein, has been shown to be a transforming protein in cell culture and promotes cell growth and survival through upregulation of NF- κ B, JAK/STAT, MAP kinase, PI3-kinase/AKT pathways (IARC 2012). Kanemitsu *et al.* (2012) examined 30 cases of NK/T cell lymphoma, nasal type, for expression profile of EBV-encoded protein; all tissues were positive by immunohistochemistry for EBER, none were positive for EBNA-2, and LMP-1 was positive in 22 cases and correlated with a localized disease ($P = 0.06$). Further, nuclear localization of phosphorylated RelA (NF- κ B) and phosphorylated Akt (PI3K) were observed in conjunction with positive cases ($P < 0.002$ and $P < 0.001$, respectively). RNA silencing experiments of LMP-1 correlated with decreased phosphorylation of RelA and Akt *in vitro* providing evidence that EBV LMP-1 expression was in part enhancing NF κ B and PI3K expression.

Further evidence that supports EBV infection as a causal agent in some forms of NK/T cell lymphoma is as follows:

- NK/T-cell lymphoma, nasal type presents as a localized disease with near 100% positivity for EBV (IARC 1997).
- EBV is clonal in these nasal tumors (IARC 1997).
- Viral proteins (LMP-1) associated with EBV latency II gene expression pattern are present in these tumors (IARC 2012).

4.3.5 Nasopharyngeal carcinoma

Research efforts have focused on addressing biological properties of EBV that may result in differences in nasopharyngeal cancer incidence among different populations. Initial characterization relied on the use of restriction enzymes to identify DNA polymorphisms describing EBV strain variation in different populations. Genome sequence analysis identified specific genetic variants (such as in LMP-1) that could potentially enhance transforming potential and virulence of EBV and enable variants of LMP-1 to evade immune recognition (Raab-Traub 2002). In addition, multiple EBV microRNAs have immune evasion functions (Tsao *et al.* 2015). Whole-EBV genome sequencing studies have revealed that the

nasopharyngeal carcinoma-derived EBV strains from endemic regions vary and show significant differences from the reported EBV genomes from non-endemic populations (Liu *et al.* 2011). A review by Lung *et al.* (2014) lists oncogenes and tumor suppressor genes important in nasopharyngeal carcinoma development and the effects of EBV infection on regulation of gene expression. The findings suggest the existence of disease-specific viral variations that may possess higher oncogenic properties, propensity for infection of epithelial cells, and persistence of the latency II program, or less efficiency in inducing host immune response, especially in the nasopharyngeal carcinoma endemic population (Liu *et al.* 2011). Linkage analysis of susceptible Chinese populations has demonstrated an association between nasopharyngeal carcinoma risk and allelic variations in human leukocyte antigen (HLA), a part of the MHC I complex of immunosurveillance. A consistent association between nasopharyngeal carcinoma and a Chinese HLA subtype, as compared to a Caucasian subtype, was detected by a genotyping study (Hildesheim *et al.* 2002). Multiple areas of hypermethylation are present in the nasopharyngeal carcinoma genome resulting in inactivation of tumor suppressor genes (e.g., *p16* and *RASSF1A*) and disruption of cell functions (i.e., cell-cycle controls, signal transduction, apoptosis, and DNA repair) (Kwong *et al.* 2002). As the efficiency of sample collection and genetic assays improve and are replicated, more details of host-virus interactions will likely be found.

Evidence that EBV is a primary causal factor in nasopharyngeal carcinoma is as follows:

- Clonal EBV episomes are detected in undifferentiated tumors (Liu *et al.* 2011).
- Studies showing early preneoplastic nasopharyngeal lesions (dysplastic lesions, carcinoma *in situ*) but not normal tissues are also infected with EBV and express latency II program gene products (Tsao *et al.* 2015, Tsang *et al.* 2014, IARC 2012, Raab-Traub 2002, Pathmanathan *et al.* 1995).

4.3.6 Gastric carcinoma

Gastric cancer is a common cancer, and it is estimated that EBV is a causal factor in 5% to 10% of all gastric cancers worldwide. Latency I/II patterns of EBV gene expression are found in these cancers with approximately 50% expressing LMP-2A which activates the NF- κ B surviving pathway (IARC 2012).

Molecular studies in humans provide strong evidence of an association between EBV and some gastric cancers. These studies show that EBV is found as a monoclonal form in a subset of human gastric cancers. Moreover, the virus produces oncogenic proteins in gastric cancer tissue that promote cell division, cell survival, and oncogenic transformation, and produces a unique molecular profile in genomic studies of gastric tumors. Recent comprehensive molecular work by the Cancer Genome Atlas Network (2014) has resulted in the identification of four distinct molecular profiles of gastric cancer, as described below. Primary gastric tumors from 295 untreated patients were analyzed and blood or non-malignant gastric tissue was used as reference for somatic alterations in the tumors. Non-malignant gastric tissue was also collected for DNA methylation and expression analyses and six molecular testing platforms were used to analyze the tissue (Cancer Genome Atlas Research 2014), reviewed by Gulley 2015). Four distinct molecular profiles were discerned for gastric cancer: EBV-positive, microsatellite instability (MSI), chromosome instability (extensive somatic copy-number aberrations [SCNAs], and

genomically stable. Molecular features unique to all EBV-positive gastric cancers were identified: hypermethylation of promoter regions resulting in gene down-regulation or silencing, mutations and gene amplifications, and expression of multiple noncoding viral RNAs. The EBV-positive group had a distinctive genetic profile, marked CpG methylation, including CDKN2A (*p16*, tumor suppressor gene), a strong immune-cell presence, and evidence of an IL-12-mediated signaling response. Some of the more frequent genetic changes are listed in Table 4-3. Promoter hypermethylation affected genes involved in cell-cycle regulation (CDKN2A - *p16*); DNA repair (GSTP1); cell adhesion and metastases (TIMP1); apoptosis (*bcl-2*); and signal transduction (PTEN). JAK2 was amplified in 12% of the EBV-positive tumors. A gene locus, 9p24.1, was linked to overexpression and amplification of JAK2, and CD274 (PD-L1) and PDCD1LG2 (PD-L2) in approximately 15% of the EBV tumor subgroup. PD-L1 and PD-L2 signaling cascades prevent T-cell proliferation and aid in escape of the cancer from immune surveillance. Further, EBV-positive tumors had mutations in PIK3CA (10-72%), ARID1A (47% to 55%), AKT2 (38%), TGFBR1 (25%), CCNA1 (25%), BCOR (23%) and MAP3K4 (21%), but rarely in *TP53*. In addition, multiple noncoding viral RNAs as well as viral LMP1, LMP2A, and EBNA1 were consistently expressed at low levels (Gulley 2015). From this unique molecular profile and the presence of activated EBV within gastric tissue, it is apparent that EBV is causal for tumor promotion. However, the pathogenesis of EBV activation in this form of gastric cancer has not been resolved.

Table 4-3. Properties of some known signaling pathways altered with EBV-related gastric cancer

Signalling Pathway	Effects
Micro RNAs	Unknown
CDKN2A (<i>p16</i>)	Tumor suppressor gene; slows progression from G1 to S phase
JAK2	Cell growth and division; LMP-1 activates
PI3K/Akt	Cell growth and division; inhibits apoptosis, promotes genomic instability, cytoskeleton change; LMP-2 activates
ERBB2	Cell growth and division
ARID1A	Cell-cycle progression
BCOR	Transcription and chromatin regulation
CD274 (PD-L1)	Immunosuppression
PDCD1LG2 (PD-L2)	Immunosuppression
IL-12	Immune cell stimulation in response to antigen
NF-κB	Resists apoptosis, cell proliferation; LMP-2A activates

Source: Gulley 2015

4.3.7 Lymphoepithelial cancer of salivary glands

Lymphoepithelial carcinomas and their association with EBV have been reported most frequently in gastric cancer and in cancer of the salivary glands but less frequently in some other epithelial malignancies such as lung and skin (IARC 2012, 1997). For cancer of the salivary glands, it is hypothesized that the close proximity of the salivary glands to the oropharyngeal area would increase exposure of these glands to the lytic form of EBV in the saliva. IARC (2012, 1997) reported a number of case-series studies detecting EBV DNA in lymphoepithelial carcinoma of the salivary glands and a case-case study presented evidence of EBV DNA in salivary tumors with lymphoepithelial histology, but not in salivary tumors of other histology

(Wang *et al.* 2004, as reported in IARC 2012). Since salivary glands can be a potential site of EBV replication as well as associated with carcinogenesis, it is important to determine whether these cancers have EBV in a latent or lytic form. As reported by IARC, viral EBER RNA and LMP-1 protein were detected in tumor tissue from 10 cases of lymphoepithelial carcinoma of the salivary gland from a Chinese population (Leung *et al.* 1995). Clonal EBV termini were associated with undifferentiated carcinoma of the parotid glands (six tumors from American Inuit) and EBER-1 RNA, LMP-1 RNA and BamHI-A rightward reading frame were expressed in the malignant cells (Raab-Traub *et al.* 1991). In both studies, an episomal form of EBV was detected in the tumor tissue, but no EBV was detected in normal adjacent tissue. These data indicate the latency II pattern of clonal EBV in these cancers.

4.4 Synthesis

EBV is highly prevalent and results in a lifelong latent infection that is refractory to immune recognition. Activated EBV transcription programs mimic B cell proliferation and survival and in some cases result in cancer. However, as with other oncoviruses, EBV infection alone is necessary but not sufficient for cancer development.

The latency patterns of EBV are associated with specific cancers and factors including immunosuppression, infectious agents, regional differences in diet, and host genetic susceptibility; all of which potentially have a role in cancer development. In primary EBV infection of naïve B cells, growth and differentiation occur due to the latency III transcription pattern that is highly immunogenic. Factors causing immunosuppression such as HIV infection or post-transplant therapies enable the latency III pattern to continue, resulting in dysregulation of cellular pathways leading to non-Hodgkin lymphoma or post-transplant lymphoproliferative disease.

Immune pressure promotes selection to latency II or I patterns, which are less immunogenic. Endemic Burkitt lymphoma is associated with co-infection with the malaria parasite, *P. falciparum*, which further enhances B cell proliferation and genetic instability of latency I pattern, resulting in *c-myc* translocation and overexpression. Hodgkin lymphoma, NK/T cell lymphoma, nasopharyngeal carcinoma, and some gastric and salivary gland lymphoepithelial cancers all express EBV latency II program. Expression of EBV latent genes LMP-1 and LMP-2 result in dysregulation of host cellular pathways and promotes oncogenesis. In addition, the genotypes of both the host and the virus, as well as dietary factors, have been linked to the prevalence of nasopharyngeal carcinoma.

5 Preliminary Listing Recommendation

Epstein-Barr virus (EBV) is known to be a human carcinogen based on sufficient evidence from studies in humans. This conclusion is based on epidemiological studies showing that it causes endemic Burkitt lymphoma (Table 5-1), Hodgkin lymphoma (Table 5-2), immunosuppression-related non-Hodgkin lymphoma (Table 5-3), NK/T cell leukemia/lymphoma, nasal type (Table 5-4), nasopharyngeal carcinoma (Table 5-5), and gastric cancer (Table 5-6) in humans, together with supporting evidence from mechanistic studies demonstrating the biological plausibility of its carcinogenicity in humans. There is also limited evidence for an association with Burkitt lymphoma (sporadic) (Table 5-1) and lymphoepithelial cancer of the salivary gland (Table 5-7) from studies in humans.

The following tables provide the preliminary level of evidence recommendations for the carcinogenicity of EBV for each tumor endpoint from studies in humans, including the key data from both epidemiological and molecular studies.

Table 5-1. Evidence for EBV and Burkitt lymphoma (endemic or sporadic) from human studies

Types of studies	Burkitt lymphoma (endemic)	Burkitt lymphoma (sporadic)
<i>Epidemiological</i>		
Number of studies reporting a positive association ^a	EBV antibodies or DNA: 7/7 case-control studies (993 cases) & 1 cohort study. All statistically significant; high RR/ORs Dose response with viral titer in cohort study and several case-control studies.	EBV antibodies: 4/5 case-control studies (113 cases). Mostly statistically nonsignificant; moderate ORs.
<i>Molecular (human tissue)</i>		
Clonality for EBV	Monoclonal	NA
% EBV-infected tumors	95%	20%
EBV protein expression	EBNA-1	NA
Level of evidence	Sufficient	Limited

EBNA = Epstein-Barr nuclear antigen; EBV = Epstein-Barr virus; NA = not available; OR = odds ratio; RR = risk ratio.

^aThe number of studies is based on those reporting risk estimates.

Table 5-2. Evidence for EBV and Hodgkin lymphoma from human studies

Type of studies	Hodgkin lymphoma
<i>Epidemiological</i>	EBV DNA: 3/3 case-control studies; very high ORs. EBV antibodies: 17/20 case-control studies & 1 nested case-control study; mostly statistically significant OR between 4 & 19. Infectious mononucleosis: 10/11 case-control studies and 7/7 cohort studies with modest ORs/RRs
<i>Molecular (human tissue)</i>	
Clonality for EBV	Monoclonal
% EBV-infected tumors	20%–50% North America and Europe; 65% Asia; 90%–100% Africa and South America
EBV protein expression	LMP-1, -2A in 50% of cases
Level of evidence	Sufficient

EBV = Epstein-Barr virus; LMP-1 = latent membrane protein 1; LMP-2A = latent membrane protein 2A; OR = odds ratio.

^a The number of studies is based on those reporting risk estimates.

Table 5-3. Evidence for EBV and immunosuppression-related non-Hodgkin lymphoma (NHL) from human studies

Type of studies	Immunosuppression-related NHL
<i>Epidemiological</i>	Consistent evidence in case-series studies Elevated statistically non-significant increased with EBV antibodies in one-nested case-control study
<i>Molecular (human tissue)</i>	
Clonality for EBV	Monoclonal
% EBV-infected tumors	100% (CNS + HIV) > 50% post transplant lymphoproliferative disease
EBV protein expression	LMP-1, -2A, -2B, EBNA _s ; EBV-specific cytotoxic
Other	T-cells protect against or reduce viral load and tumor size
Level of evidence	Sufficient

CNS = central nervous system; EBNA = Epstein-Barr nuclear antigen; EBV = Epstein-Barr virus; HIV = human immunodeficiency virus; LMP-1 = latent membrane protein 1; LMP-2A = latent membrane protein 2A; LMP-2B = latent membrane protein 2B.

Table 5-4. Evidence for EBV and NK/T cell leukemia/lymphoma, nasal type from studies in humans

Type of studies	NK/T cell leukemia/lymphoma, nasal type
<i>Epidemiological</i>	Consistent evidence in case series studies; at least 16 case-series with more than 400 cases. 2 case-comparison studies: EBV DNA found in plasma or CD3+ cells from cases but not from controls.
<i>Molecular (human tissue)</i>	
Clonality for EBV	Monoclonal
% EBV-infected tumors	100%
EBV protein expression	EBNA-1, LMP-1, -2A
Other	EBV found in majority of CD56+ tumors.
Level of evidence	Sufficient

EBNA = Epstein-Barr nuclear antigen; EBV = Epstein-Barr virus; LMP-1 = latent membrane protein 1; LMP-2A = latent membrane protein 2A.

Table 5-5. Evidence for EBV and nasopharyngeal carcinoma from human studies

Supporting evidence	Nasopharyngeal carcinoma
Epidemiological	EBV antibody: 11/11 case-control and 2 cohort studies; high to very high stat. sig. RRs; no association in 2 small nested case-control studies.
Number of studies reporting a positive association ^a	EBV DNA: 6/6 case-control studies; very high RR
Molecular (human tissue)	
Clonality for EBV	Monoclonal in precancerous/cancer
% EBV-infected tumors	98% in non-keratinizing tumors
EBV protein expression	EBNA-1, LMP-1, -2A
Level of evidence	Sufficient

EBNA = Epstein-Barr nuclear antigen; EBV = Epstein-Barr virus; LMP-1 = latent membrane protein 1; LMP-2A = latent membrane protein 2A; RR = risk ratio.

^a The number of studies is based on those reporting risk estimates.

Table 5-6. Evidence for EBV and gastric cancer from human studies

Type of studies	Gastric cancer
Epidemiological	3/3 case-control studies (77 EBV cases/184 gastric); statistically significant high ORs
Number of studies reporting a positive association	2/3 nested case-control studies; statistically non-significant modest ORs
Molecular (human tissue)	
Clonality for EBV	Monoclonal
% EBV-infected tumors	8%–11%
EBV protein expression	EBNA-1, LMP-1, -2A
Other	Unique molecular profile
Level of evidence	Sufficient

EBNA = Epstein-Barr nuclear antigen; EBV = Epstein-Barr virus; LMP-1 = latent membrane protein 1; LMP-2A = latent membrane protein 2A; OR = odds ratio.

Table 5-7. Evidence for EBV and lymphoepithelial carcinoma of the salivary gland from human studies

Type of study	Lymphoepithelial carcinoma/salivary gland
Epidemiological	Consistent evidence in case series studies (208/209 cases) A case-case study found EBV DNA in salivary gland lymphoepithelial carcinoma tumors but not other type of salivary gland tumors
Molecular (human tissue)	
Clonality for EBV	Monoclonal (few studies)
% EBV-infected tumors	100%
EBV protein expression	EBNA-1, LMP-1, -2A (few samples)
Other	No additional supporting mechanistic data.
Level of evidence	Limited

EBNA = Epstein-Barr nuclear antigen; EBV = Epstein-Barr virus; LMP-1 = latent membrane protein 1; LMP-2A = latent membrane protein 2A.

This Page Intentionally Left Blank

References

1. Ablashi DV, De-The GB, Easton JM, Liabeuf A, Levine PH, Armstrong GR. 1974. Antibodies to Epstein-Barr virus (EBV) antigens in sera of American Burkitt lymphoma patients. *Biomedicine* 21(7): 288-293.
2. ACS. 2015a. *Viruses that can lead to cancer*. American Cancer Society. Updated on 4/27/15. <http://www.cancer.org/cancer/cancercauses/othercarcinogens/infectiousagents/infectiousagentsandcancer/infectious-agents-and-cancer-viruses>. Accessed on 6/18/15.
3. ACS. 2015b. *What are the Risk Factors for Hodgkin's Disease?* American Cancer Society. Updated on 3/4/15. <http://www.cancer.org/cancer/hodgkindisease/detailedguide/hodgkin-disease-risk-factors>. Accessed on 10/28/15.
4. Adham M, Kurniawan AN, Muhtadi AI, Roezin A, Hermani B, Gondhowiardjo S, Tan IB, Middeldorp JM. 2012. Nasopharyngeal carcinoma in Indonesia: epidemiology, incidence, signs, and symptoms at presentation. *Chin J Cancer* 31(4): 185-196.
5. Al-Mansour Z, Nelson BP, Evens AM. 2013. Post-transplant lymphoproliferative disease (PTLD): risk factors, diagnosis, and current treatment strategies. *Curr Hematol Malig Rep* 8(3): 173-183. (as cited in Jimenez 2015)
6. Aldred MA, Eng C. 2006. SNP'ing at nasopharyngeal cancer susceptibility: for whom the bell TOLLs. *Cancer Biol Ther* 5(10): 1292-1293. (as cited in Lung *et al* 2014)
7. Alexander FE, Jarrett RF, Lawrence D, Armstrong AA, Freeland J, Gokhale DA, Kane E, Taylor GM, Wright DH, Cartwright RA. 2000. Risk factors for Hodgkin's disease by Epstein-Barr virus (EBV) status: prior infection by EBV and other agents. *Br J Cancer* 82(5): 1117-1121.
8. Alexander FE, Lawrence DJ, Freeland J, Krajewski AS, Angus B, Taylor GM, Jarrett RF. 2003. An epidemiologic study of index and family infectious mononucleosis and adult Hodgkin's disease (HD): evidence for a specific association with EBV+ve HD in young adults. *Int J Cancer* 107(2): 298-302.
9. Ambinder RF, Cesarman E. 2007. Clinical and pathological aspects of EBV and KSHV Infection. In *Human Herpesviruses*. Arvin A, Campadelli-Fiume G, Mocarski E, et al., eds. Cambridge: Cambridge University. pp. 885-903. (as cited in IARC 2012)
10. Armstrong RW, Armstrong MJ. 1983. Environmental risk factors and nasopharyngeal carcinoma in Selangor, Malaysia: a cross-ethnic perspective. *Ecol Dis* 2: 185-198. (as cited in IARC 1997)
11. Arvey A, Tempera I, Tsai K, Chen HS, Tikhmyanova N, Klichinsky M, Leslie C, Lieberman PM. 2012. An atlas of the Epstein-Barr virus transcriptome and epigenome reveals host-virus regulatory interactions. *Cell Host Microbe* 12(2): 233-245.

12. Auwaerter PG. 2015. *Johns Hopkins HIV Guide: Epstein-Barr virus*. Johns Hopkins Medicine. Updated on 6/2/2015.
http://www.hopkinsguides.com/hopkins/view/Johns_Hopkins_HIV_Guide/545067/all/Epstein_Barr_virus. Accessed on 6/17/15.
13. Balfour HH, Jr., Sifakis F, Sliman JA, Knight JA, Schmeling DO, Thomas W. 2013. Age-specific prevalence of Epstein-Barr virus infection among individuals aged 6-19 years in the United States and factors affecting its acquisition. *J Infect Dis* 208(8): 1286-1293.
14. Balfour HH, Jr. 2014. Progress, prospects, and problems in Epstein-Barr virus vaccine development. *Curr Opin Virol* 6: 1-5.
15. Barrionuevo C, Zaharia M, Martinez MT, Taxa L, Misad O, Moscol A, Sarria G, Guerrero I, Casanova L, Flores C, Zevallos-Giampietri EA. 2007. Extranodal NK/T-cell lymphoma, nasal type: study of clinicopathologic and prognosis factors in a series of 78 cases from Peru. *Appl Immunohistochem Mol Morphol* 15(1): 38-44.
16. Basso K, Dalla-Favera R. 2015. Germinal centres and B cell lymphomagenesis. *Nat Rev Immunol* 15(3): 172-184.
17. Bernard SM, Cartwright RA, Darwin CM, Richards ID, Roberts B, O'Brien C, Bird CC. 1987. Hodgkin's disease: case control epidemiological study in Yorkshire. *Br J Cancer* 55(1): 85-90.
18. Berrington de Gonzalez A, Urban MI, Sitas F, Blackburn N, Hale M, Patel M, Ruff P, Sur R, Newton R, Beral V. 2006. Antibodies against six human herpesviruses in relation to seven cancers in black South Africans: a case control study. *Infect Agent Cancer* 1: 2.
19. Biggar RJ, Henle G, Bocker J, Lennette ET, Fleisher G, Henle W. 1978a. Primary Epstein-Barr virus infections in African infants. II. Clinical and serological observations during seroconversion. *Int J Cancer* 22(3): 244-250.
20. Biggar RJ, Henle W, Fleisher G, Bocker J, Lennette ET, Henle G. 1978b. Primary Epstein-Barr virus infections in African infants. I. Decline of maternal antibodies and time of infection. *Int J Cancer* 22(3): 239-243.
21. Borza CM, Hutt-Fletcher LM. 2002. Alternate replication in B cells and epithelial cells switches tropism of Epstein-Barr virus. *Nat Med* 8(6): 594-599.
22. Boysen T, Friberg J, Stribolt K, Hamilton-Dutoit S, Goertz S, Wohlfahrt J, Melbye M. 2011. Epstein-Barr virus-associated gastric carcinoma among patients with pernicious anemia. *Int J Cancer* 129(11): 2756-2760.
23. Burke AP, Yen TS, Shekitka KM, Sobin LH. 1990. Lymphoepithelial carcinoma of the stomach with Epstein-Barr virus demonstrated by polymerase chain reaction. *Mod Pathol* 3(3): 377-380.
24. Cancer Genome Atlas Research N. 2014. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature* 513(7517): 202-209. (as cited in Gulley *et al* 2015)

25. Carbone A, Gloghini A, Dotti G. 2008. EBV-associated lymphoproliferative disorders: classification and treatment. *Oncologist* 13(5): 577-585.
26. Carpenter LM, Newton R, Casabonne D, Ziegler J, Mbulaiteye S, Mbidde E, Wabinga H, Jaffe H, Beral V. 2008. Antibodies against malaria and Epstein-Barr virus in childhood Burkitt lymphoma: a case-control study in Uganda. *Int J Cancer* 122(6): 1319-1323.
27. Carter CD, Brown TM, Jr., Herbert JT, Heath CW, Jr. 1977. Cancer incidence following infectious mononucleosis. *Am J Epidemiol* 105(1): 30-36.
28. Cavdar AO, Yavuz G, Babacan E, Gozdasoglu S, Unal E, Ertem U, Pamir A, Yucesan S, Gokcora H, Uluoglu O, et al. 1994. Burkitt's lymphoma in Turkish children: clinical, viral [EBV] and molecular studies. *Leuk Lymphoma* 14(3-4): 323-330.
29. CDC. 2011. *List of Vaccines Used in United States*. Centers for Disease Control and Prevention. Updated on 8/18/2011. <http://www.cdc.gov/vaccines/vpd-vac/vaccines-list.htm>. Accessed on 6/17/15.
30. CDC. 2014a. *About infectious mononucleosis*. Centers for Disease Control and Prevention. Updated on 1/7/2014. <http://www.cdc.gov/epstein-barr/about-mono.html>. Accessed on 6/18/15.
31. CDC. 2014b. *About Epstein-Barr virus (EBV)*. Centers for Disease Control and Prevention. Updated on 1/6/14. <http://www.cdc.gov/epstein-barr/about-ebv.html>. Accessed on 6/18/15.
32. Chaigne-Delalande B, Li FY, O'Connor GM, Lukacs MJ, Jiang P, Zheng L, Shatzer A, Biancalana M, Pittaluga S, Matthews HF, Jancel TJ, Blesing JJ, Marsh RA, Kuijpers TW, Nichols KE, Lucas CL, Nagpal S, Mehmet H, Su HC, Cohen JI, Uzel G, Lenardo MJ. 2013. Mg²⁺ regulates cytotoxic functions of NK and CD8 T cells in chronic EBV infection through NKG2D. *Science* 341(6142): 186-191.
33. Chan CK, Mueller N, Evans A, Harris NL, Comstock GW, Jellum E, Magnus K, Orentreich N, Polk BF, Vogelmann J. 1991. Epstein-Barr virus antibody patterns preceding the diagnosis of nasopharyngeal carcinoma. *Cancer Causes Control* 2(2): 125-131.
34. Chan JKC, Bray F, McCarron P, et al. 2005. Nasopharyngeal carcinoma. In *World Health Organization Classification of Tumours – Pathology and Genetics of Head and Neck Tumours*. Barnes EL, Eveson JW, Reichart P, Sidransky D, eds. Lyon, France: International Agency for Research on Cancer. pp. 85-97. (as cited in IARC 2012)
35. Chan WY, Chan EK, Chow JH. 2001. Epstein-Barr virus-associated gastric lymphomas are distinct from mucosa-associated lymphoid tissue-type lymphomas: genetic abnormalities of p53 gene. *Diagn Mol Pathol* 10(3): 153-160.
36. Chang ET, Adami HO. 2006. The enigmatic epidemiology of nasopharyngeal carcinoma. *Cancer Epidemiol Biomarkers Prev* 15(10): 1765-1777.
37. Chen C-J, Wang Y-F, Shieh T, Chen J-Y, Liu M-Y. 1988. Multifactorial etiology of nasopharyngeal carcinoma. Epstein-Barr virus, familial tendency and environmental co-

- factors. In *Head and Neck Oncology Research*. Wolf GT, Carey TE, eds. Amsterdam: Kugler Publications. pp. 469-476. (as cited in IARC 1997)
38. Chesnokova LS, Hutt-Fletcher LM. 2011. Fusion of Epstein-Barr virus with epithelial cells can be triggered by alphavbeta5 in addition to alphavbeta6 and alphavbeta8, and integrin binding triggers a conformational change in glycoproteins gHgL. *J Virol* 85(24): 13214-13223.
 39. Chien YC, Chen JY, Liu MY, Yang HI, Hsu MM, Chen CJ, Yang CS. 2001. Serologic markers of Epstein-Barr virus infection and nasopharyngeal carcinoma in Taiwanese men. *N Engl J Med* 345(26): 1877-1882.
 40. Chin Y-M, Mushiroda T, Takahashi A, Kubo M, Krishnan G, Yap L-F, Teo S-H, Lim PV-H, Yap Y-Y, Pua K-C, Kamatani N, Nakamura Y, Sam C-K, Soo-Beng Koo A, Malaysian NPC Study Group, Ng C-C. 2014. HLA-A SNPs and amino acid variants are associated with nasopharyngeal carcinoma in Malaysian Chinese. *Int J Cancer*.
 41. Chow KU, Mitrou PS, Geduldig K, Helm EB, Hoelzer D, Brodt HR. 2001. Changing incidence and survival in patients with aids-related non-Hodgkin's lymphomas in the era of highly active antiretroviral therapy (HAART). *Leuk Lymphoma* 41(1-2): 105-116.
 42. Cohen JL. 2009. Optimal treatment for chronic active Epstein-Barr virus disease. *Pediatr Transplant* 13(4): 393-396.
 43. Cohen JL. 2015. Epstein-barr virus vaccines. *Clin Transl Immunology* 4(1): e32.
 44. Condon LM, Cederberg LE, Rabinovitch MD, Liebo RV, Go JC, Delaney AS, Schmeling DO, Thomas W, Balfour HH, Jr. 2014. Age-specific prevalence of Epstein-Barr virus infection among Minnesota children: effects of race/ethnicity and family environment. *Clin Infect Dis* 59(4): 501-508.
 45. Connelly RR, Christine BW. 1974. A cohort study of cancer following infectious mononucleosis. *Cancer Res* 34(5): 1172-1178.
 46. Crawford DH. 2001. Biology and disease associations of Epstein-Barr virus. *Philos Trans R Soc Lond B Biol Sci* 356(1408): 461-473.
 47. Crew KD, Neugut AI. 2006. Epidemiology of gastric cancer. *World J Gastroenterol* 12(3): 354-362.
 48. de Aquino PF, Carvalho PC, da Gama Fischer JS, de Souza AQ, Viana JS, Chalub SR, de Souza AD, Carvalho MG. 2012. Epstein-Barr virus DNA associated with gastric adenocarcinoma and adjacent non-cancerous mucosa in patients from Manaus, Brazil. *Genet Mol Res* 11(4): 4442-4446.
 49. de The G, Lavoué M-E, Muenz L. 1978. Differences in EBV antibody titres of patients with nasopharyngeal carcinoma originating from high, intermediate and low incidence areas. *IARC Sci Publ* 20(471-481).

50. Dinand V, Dawar R, Arya LS, Unni R, Mohanty B, Singh R. 2007. Hodgkin's lymphoma in Indian children: prevalence and significance of Epstein-Barr virus detection in Hodgkin's and Reed-Sternberg cells. *Eur J Cancer* 43(1): 161-168.
51. Dowd JB, Palermo T, Brite J, McDade TW, Aiello A. 2013. Seroprevalence of Epstein-Barr virus infection in U.S. children ages 6-19, 2003-2010. *PLoS One* 8(5): e64921.
52. Epstein MA, Achong BG, Barr YM. 1964. Virus Particles in Cultured Lymphoblasts from Burkitt's Lymphoma. *Lancet* 1(7335): 702-703.
53. Evans AS, Gutensohn NM. 1984. A population-based case-control study of EBV and other viral antibodies among persons with Hodgkin's disease and their siblings. *Int J Cancer* 34(2): 149-157.
54. FDA. 2015. *Complete List of Vaccines Licensed for Immunization and Distribution in the US*. U.S. Food and Drug Administration. Updated on 6/17/15. <http://www.fda.gov/BiologicsBloodVaccines/Vaccines/ApprovedProducts/ucm093833.htm>. Accessed on 6/18/15.
55. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. 2015. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 136(5): E359-386.
56. Flavell KJ, Murray PG. 2000. Hodgkin's disease and the Epstein-Barr virus. *Mol Pathol* 53(5): 262-269.
57. Gallagher A, Armstrong AA, MacKenzie J, Shield L, Khan G, Lake A, Proctor S, Taylor P, Clements GB, Jarrett RF. 1999. Detection of Epstein-Barr virus (EBV) genomes in the serum of patients with EBV-associated Hodgkin's disease. *Int J Cancer* 84(4): 442-448.
58. Garfin PM, Twist CJ, Talavera F, Conrad ME, Shapiro MD. 2015. *Posttransplant Lymphoproliferative Disease*. Updated on 4/9/15. <http://emedicine.medscape.com/article/431364-overview-showall>. Accessed on 6/17/15.
59. Geser A, Charnay N, Day NE, de-The G, Ho HC. 1978. Environmental factors in the etiology of nasopharyngeal carcinoma: report on a case-control study in Hong Kong. *IARC Sci Publ*(20): 213-229. (as cited in IARC 1997)
60. Geser A, de The G, Lenoir G, Day NE, Williams EH. 1982. Final case reporting from the Ugandan prospective study of the relationship between EBV and Burkitt's lymphoma. *Int J Cancer* 29(4): 397-400.
61. Geser A, Brubaker G, Draper CC. 1989. Effect of a malaria suppression program on the incidence of African Burkitt's lymphoma. *Am J Epidemiol* 129(4): 740-752.
62. Glaser SL, Keegan TH, Clarke CA, Trinh M, Dorfman RF, Mann RB, DiGiuseppe JA, Ambinder RF. 2005. Exposure to childhood infections and risk of Epstein-Barr virus--defined Hodgkin's lymphoma in women. *Int J Cancer* 115(4): 599-605.

63. Gloghini A, Dolcetti R, Carbone A. 2013. Lymphomas occurring specifically in HIV-infected patients: from pathogenesis to pathology. *Semin Cancer Biol* 23(6): 457-467.
64. Gotlieb-Stematsky T, Ramot B, Vonsover B, Aghai E, Kende G, Ninio M, Modan M. 1976. Antibodies to Epstein-Barr viral capsid and early antigens associated with Burkitt's lymphoma and lymphoblastic lymphosarcoma in Israel. *J Natl Cancer Inst* 56(4): 721-723.
65. Gotlieb-Stematsky T, Vonsover A, Ramot B, Zaizov R, Nordan U, Aghal E, Kende G, Modan M. 1975. Antibodies to Epstein-Barr virus in patients with Hodgkin's disease and leukemia. *Cancer* 36(5): 1640-1645.
66. Grulich AE, van Leeuwen MT, Falster MO, Vajdic CM. 2007a. Incidence of cancers in people with HIV/AIDS compared with immunosuppressed transplant recipients: a meta-analysis. *Lancet* 370(9581): 59-67.
67. Grulich AE, Vajdic CM, Cozen W. 2007b. Altered immunity as a risk factor for non-Hodgkin lymphoma. *Cancer Epidemiol Biomarkers Prev* 16(3): 405-408.
68. Grywalska E, Rolinski J. 2015. Epstein-Barr virus-associated lymphomas. *Semin Oncol* 42(2): 291-303.
69. Gulley ML, Tang W. 2010. Using Epstein-Barr viral load assays to diagnose, monitor, and prevent posttransplant lymphoproliferative disorder. *Clin Microbiol Rev* 23(2): 350-366.
70. Gulley ML. 2015. Genomic assays for Epstein-Barr virus-positive gastric adenocarcinoma. *Exp Mol Med* 47: e134.
71. Gutensohn N, Cole P. 1981. Childhood social environment and Hodgkin's disease. *N Engl J Med* 304(3): 135-140.
72. Gutensohn NM. 1982. Social class and age at diagnosis of Hodgkin's disease: new epidemiologic evidence for the "two-disease hypothesis". *Cancer Treat Rep* 66(4): 689-695.
73. Hauke R, Smir B, Greiner T, Bierman P, Tarantolo S, Anderson J, Shaw B, Armitage J. 2001. Clinical and pathological features of posttransplant lymphoproliferative disorders: influence on survival and response to treatment. *Ann Oncol* 12(6): 831-834. (as cited in Garfin *et al* 2015)
74. He YJ, Jia XS, Hasui K, Wang EH, He AG. 2007. [Nasal and pharyngeal non-Hodgkin lymphomas and their relationship with Epstein-Barr virus: a report of 158 cases]. *Zhonghua Bing Li Xue Za Zhi* 36(2): 94-97.
75. Henderson BE, Louie E. 1978. Discussion of risk factors for nasopharyngeal carcinoma. *IARC Sci Publ*(20): 251-260. (as cited in IARC 1997)
76. Henderson BE, Dworsky R, Pike MC, Baptista J, Menck H, Preston-Martin S, Mack T. 1979. Risk factors for nodular sclerosis and other types of Hodgkin's disease. *Cancer Res* 39(11): 4507-4511.

77. Henle G, Henle W, Clifford P, Diehl V, Kafuko GW, Kirya BG, Klein G, Morrow RH, Munube GM, Pike P, Tukei PM, Ziegler JL. 1969. Antibodies to Epstein-Barr virus in Burkitt's lymphoma and control groups. *J Natl Cancer Inst* 43(5): 1147-1157.
78. Henle G, Henle W, Klein G, Gunven P, Clifford P, Morrow RH, Ziegler JL. 1971. Antibodies to early Epstein-Barr virus-induced antigens in Burkitt's lymphoma. *J Natl Cancer Inst* 46(4): 861-871.
79. Hildesheim A, Apple RJ, Chen CJ, Wang SS, Cheng YJ, Klitz W, Mack SJ, Chen IH, Hsu MM, Yang CS, Brinton LA, Levine PH, Erlich HA. 2002. Association of HLA class I and II alleles and extended haplotypes with nasopharyngeal carcinoma in Taiwan. *J Natl Cancer Inst* 94(23): 1780-1789.
80. Hilgers F, Hilgers J. 1976. An immunofluorescence technique with counterstain on fixed cells for the detection of antibodies to human herpesviruses; antibody patterns in patients with Hodgkin's disease and nasopharyngeal carcinoma. *Intervirology* 7(6): 309-327.
81. Hirshaut Y, Cohen MH, Stevens DA. 1973. Epstein-Barr-virus antibodies in American and African Burkitt's lymphoma. *Lancet* 2(7821): 114-116.
82. Hjalgrim H, Askling J, Sorensen P, Madsen M, Rosdahl N, Storm HH, Hamilton-Dutoit S, Eriksen LS, Frisch M, Ekbom A, Melbye M. 2000. Risk of Hodgkin's disease and other cancers after infectious mononucleosis. *J Natl Cancer Inst* 92(18): 1522-1528.
83. Hjalgrim H, Smedby KE, Rostgaard K, Molin D, Hamilton-Dutoit S, Chang ET, Ralfkiaer E, Sundstrom C, Adami HO, Glimelius B, Melbye M. 2007. Infectious mononucleosis, childhood social environment, and risk of Hodgkin lymphoma. *Cancer Res* 67(5): 2382-2388.
84. IARC. 1997. Epstein-Barr virus. In *Epstein-Barr Virus and Kaposi's Sarcoma Herpesvirus/Human Herpesvirus 8*. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans, vol. 70. Lyon, France: International Agency for Research on Cancer. pp. 47-373.
85. IARC. 2012. Epstein-Barr virus. In *Biological Agents*. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans, vol. 100B. Lyon, France: International Agency for Research on Cancer. pp. 49-92.
86. Imai S, Nishikawa J, Takada K. 1998. Cell-to-cell contact as an efficient mode of Epstein-Barr virus infection of diverse human epithelial cells. *J Virol* 72(5): 4371-4378.
87. Ito Y, Kawanishi M, Harayama T, Takabayashi S. 1981. Combined effect of the extracts from *Croton tiglium*, *Euphorbia lathyris* or *Euphorbia tirucalli* and n-butyrate on Epstein-Barr virus expression in human lymphoblastoid P3HR-1 and Raji cells. *Cancer Lett* 12(3): 175-180. (as cited in IARC 1997)
88. Jarrett RF. 2003. Risk factors for Hodgkin's lymphoma by EBV status and significance of detection of EBV genomes in serum of patients with EBV-associated Hodgkin's lymphoma. *Leuk Lymphoma* 44 Suppl 3: S27-32.

89. Ji MF, Wang DK, Yu YL, Guo YQ, Liang JS, Cheng WM, Zong YS, Chan KH, Ng SP, Wei WI, Chua DT, Sham JS, Ng MH. 2007. Sustained elevation of Epstein-Barr virus antibody levels preceding clinical onset of nasopharyngeal carcinoma. *Br J Cancer* 96(4): 623-630.
90. Jimenez S. 2015. Epstein-Barr virus-associated post-transplantation lymphoproliferative disorder: potential treatments and implications for nursing practice. *Clin J Oncol Nurs* 19(1): 94-98.
91. Kanemitsu N, Isobe Y, Masuda A, Momose S, Higashi M, Tamaru J, Sugimoto K, Komatsu N. 2012. Expression of Epstein-Barr virus-encoded proteins in extranodal NK/T-cell Lymphoma, nasal type (ENKL): differences in biologic and clinical behaviors of LMP1-positive and -negative ENKL. *Clin Cancer Res* 18(8): 2164-2172.
92. Kennedy G, Komano J, Sugden B. 2003. Epstein-Barr virus provides a survival factor to Burkitt's lymphomas. *Proc Natl Acad Sci U S A* 100(24): 14269-14274.
93. Kim Y, Shin A, Gwack J, Ko KP, Kim CS, Park SK, Hong YC, Kang D, Yoo KY. 2009b. Epstein-Barr virus antibody level and gastric cancer risk in Korea: a nested case-control study. *Br J Cancer* 101(3): 526-529.
94. Klein G, Geering G, Old LJ, Henle G, Henle W, Clifford P. 1970. Comparison of the anti-EBV titer and the EBV-associated membrane reactive and precipitating antibody levels in the sera of Burkitt lymphoma and nasopharyngeal carcinoma patients and controls. *Int J Cancer* 5(2): 185-194.
95. Koshiol J, Qiao YL, Mark SD, Dawsey SM, Abnet CC, Kamangar F, Lennette ET, Dong ZW, Taylor PR. 2007. Epstein-Barr virus serology and gastric cancer incidence and survival. *Br J Cancer* 97(11): 1567-1569.
96. Kvale G, Hoiby EA, Pedersen E. 1979. Hodgkin's disease in patients with previous infectious mononucleosis. *Int J Cancer* 23(5): 593-597.
97. Kwong J, Lo KW, To KF, Teo PM, Johnson PJ, Huang DP. 2002. Promoter hypermethylation of multiple genes in nasopharyngeal carcinoma. *Clin Cancer Res* 8(1): 131-137.
98. Lange B, Arbeter A, Hewetson J, Henle W. 1978. Longitudinal study of Epstein-Barr virus antibody titers and excretion in pediatric patients with Hodgkin's disease. *Int J Cancer* 22(5): 521-527.
99. Lanier AP, Henle W, Bender TR, Henle G, Talbot ML. 1980. Epstein-Barr virus-specific antibody titers in seven Alaskan natives before and after diagnosis of nasopharyngeal carcinoma. *Int J Cancer* 26(2): 133-137.
100. Lee HP, Gourley L, Duffy SW, Esteve J, Lee J, Day NE. 1994. Preserved foods and nasopharyngeal carcinoma: a case-control study among Singapore Chinese. *Int J Cancer* 59(5): 585-590. (as cited in IARC 1997)

101. Lee J, Suh C, Park YH, Ko YH, Bang SM, Lee JH, Lee DH, Huh J, Oh SY, Kwon HC, Kim HJ, Lee SI, Kim JH, Park J, Oh SJ, Kim K, Jung C, Park K, Kim WS. 2006. Extranodal natural killer T-cell lymphoma, nasal-type: a prognostic model from a retrospective multicenter study. *J Clin Oncol* 24(4): 612-618.
102. Lee JT, Ko CY. 2005. Has survival improved for nasopharyngeal carcinoma in the United States? *Otolaryngol Head Neck Surg* 132(2): 303-308.
103. Lehtinen T, Lumio J, Dillner J, Hakama M, Knekt P, Lehtinen M, Teppo L, Leinikki P. 1993. Increased risk of malignant lymphoma indicated by elevated Epstein-Barr virus antibodies--a prospective study. *Cancer Causes Control* 4(3): 187-193.
104. Lei KI, Chan LY, Chan WY, Johnson PJ, Lo YM. 2000. Quantitative analysis of circulating cell-free Epstein-Barr virus (EBV) DNA levels in patients with EBV-associated lymphoid malignancies. *Br J Haematol* 111(1): 239-246.
105. Lennette ET, Rymo L, Yadav M, Masucci G, Merk K, Timar L, Klein G. 1993. Disease-related differences in antibody patterns against EBV-encoded nuclear antigens EBNA 1, EBNA 2 and EBNA 6. *Eur J Cancer* 29A(11): 1584-1589.
106. Lennette ET, Winberg G, Yadav M, Enblad G, Klein G. 1995. Antibodies to LMP2A/2B in EBV-carrying malignancies. *Eur J Cancer* 31A(11): 1875-1878.
107. Leung SY, Chung LP, Yuen ST, Ho CM, Wong MP, Chan SY. 1995. Lymphoepithelial carcinoma of the salivary gland: in situ detection of Epstein-Barr virus. *J Clin Pathol* 48(11): 1022-1027.
108. Levine PH, O'Connor GT, Bernard CW. 1972. Antibodies to Epstein-Barr virus (EBV) in American patients with Burkitt's lymphoma. *Cancer* 30(3): 610-615. (as cited in IARC 1997)
109. Levine PH, Stemmermann G, Lennette ET, Hildesheim A, Shibata D, Nomura A. 1995. Elevated antibody titers to Epstein-Barr virus prior to the diagnosis of Epstein-Barr-virus-associated gastric adenocarcinoma. *Int J Cancer* 60(5): 642-644.
110. Lim WH, Russ GR, Coates PT. 2006. Review of Epstein-Barr virus and post-transplant lymphoproliferative disorder post-solid organ transplantation. *Nephrology (Carlton)* 11(4): 355-366.
111. Lin JC, Smith MC, Pagano JS. 1982. Effects of 12-O-tetradecanoyl-phorbol-13-acetate on cell proliferation and Epstein-Barr virus DNA replication. *Virology* 117(1): 186-194. (as cited in IARC 1997)
112. Linabery AM, Erhardt EB, Fonstad RK, Ambinder RF, Bunin GR, Ross JA, Spector LG, Grufferman S. 2014. Infectious, autoimmune and allergic diseases and risk of Hodgkin lymphoma in children and adolescents: a Children's Oncology Group study. *Int J Cancer* 135(6): 1454-1469.
113. Liu P, Fang X, Feng Z, Guo YM, Peng RJ, Liu T, Huang Z, Feng Y, Sun X, Xiong Z, Guo X, Pang SS, Wang B, Lv X, Feng FT, Li DJ, Chen LZ, Feng QS, Huang WL, Zeng MS,

- Bei JX, Zhang Y, Zeng YX. 2011. Direct sequencing and characterization of a clinical isolate of Epstein-Barr virus from nasopharyngeal carcinoma tissue by using next-generation sequencing technology. *J Virol* 85(21): 11291-11299.
114. Lo YM, Chan WY, Ng EK, Chan LY, Lai PB, Tam JS, Chung SC. 2001. Circulating Epstein-Barr virus DNA in the serum of patients with gastric carcinoma. *Clin Cancer Res* 7(7): 1856-1859.
115. Lombardi L, Newcomb EW, Dalla-Favera R. 1987. Pathogenesis of Burkitt lymphoma: expression of an activated c-myc oncogene causes the tumorigenic conversion of EBV-infected human B lymphoblasts. *Cell* 49(2): 161-170.
116. Lu J, Murakami M, Verma SC, Cai QL, Haldar S, Kaul R, Wasik MA, Middeldorp J, Robertson ES. 2011. Epstein-Barr Virus nuclear antigen 1 (EBNA1) confers resistance to apoptosis in EBV-positive B-lymphoma cells through up-regulation of survivin. *Virology* 410(1): 64-75.
117. Lung ML, Cheung AK, Ko JM, Lung HL, Cheng Y, Dai W. 2014. The interplay of host genetic factors and Epstein-Barr virus in the development of nasopharyngeal carcinoma. *Chin J Cancer* 33(11): 556-568.
118. MacMahon B. 1966. Epidemiology of Hodgkin's disease. *Cancer Res* 26(6): 1189-1201.
119. Mbulaiteye SM, Pullarkat ST, Nathwani BN, Weiss LM, Rao N, Emmanuel B, Lynch CF, Hernandez B, Neppalli V, Hawes D, Cockburn MG, Kim A, Williams M, Altekruse S, Bhatia K, Goodman MT, Cozen W. 2014. Epstein-Barr virus patterns in US Burkitt lymphoma tumors from the SEER residual tissue repository during 1979-2009. *APMIS* 122(1): 5-15.
120. Merk K, Lennette E, Holm G, Johansson B, Masucci G, Grimfors G, Klein G, Bjorkholm M. 1995. Antibodies to Epstein-Barr virus in relation to clinical characteristics of untreated patients with Hodgkin's Disease. *Cancer Res Ther Control* 4: 223-229.
121. Mesri EA, Feitelson MA, Munger K. 2014. Human viral oncogenesis: a cancer hallmarks analysis. *Cell Host Microbe* 15(3): 266-282.
122. Miller RW, Beebe GW. 1973. Infectious mononucleosis and the empirical risk of cancer. *J Natl Cancer Inst* 50(2): 315-321.
123. Mitarnun W, Suwiwat S, Pradutkanchana J, Saechan V, Ishida T, Takao S, Mori A. 2002. Epstein-Barr virus-associated peripheral T-cell and NK-cell proliferative disease/lymphoma: clinicopathologic, serologic, and molecular analysis. *Am J Hematol* 70(1): 31-38.
124. Mochanko K, Fejes M, Brezavscek DM, Suarez A, Bachmann AE. 1979. The relation between Epstein-Barr virus antibodies and clinical symptomatology and immunodeficiency in patients with Hodgkin's disease. *Cancer* 44(6): 2065-2070.

125. Mohamed G, Vrzalikova K, Cader FZ, Vockerodt M, Nagy E, Flodr P, Yap LF, Diepstra A, Kluin PM, Rosati S, Murray P. 2014. Epstein-Barr virus, the germinal centre and the development of Hodgkin's lymphoma. *J Gen Virol* 95(Pt 9): 1861-1869.
126. Moormann AM, Heller KN, Chelimo K, Embury P, Ploutz-Snyder R, Otieno JA, Oduor M, Munz C, Rochford R. 2009. Children with endemic Burkitt lymphoma are deficient in EBNA1-specific IFN-gamma T cell responses. *Int J Cancer* 124(7): 1721-1726. (as cited in IARC 2012)
127. Morrow Jr. RH. 1985. Epidemiological evidence for the role of falciparum malaria in the pathogenesis of Burkitt's lymphoma. In *Burkitt's Lymphoma: A Human Cancer Model*. IARC Scientific Publications No. 60. Lenoir G, O'Connor G, Olweny CLM, eds. Lyon, France: International Agency for Research on Cancer. pp. 177-186. (as cited in IARC 1997)
128. Mosier DE, Gulizia RJ, Baird SM, Spector S, Spector D, Kipps TJ, Fox RI, Carson DA, Cooper N, Richman DD, et al. 1989. Studies of HIV infection and the development of Epstein-Barr virus-related B cell lymphomas following transfer of human lymphocytes to mice with severe combined immunodeficiency. *Curr Top Microbiol Immunol* 152: 195-199.
129. Moumad K, Lascorz J, Bevier M, Khyatti M, Ennaji MM, Benider A, Huhn S, Lu S, Chouchane L, Corbex M, Hemminki K, Forsti A. 2013. Genetic polymorphisms in host innate immune sensor genes and the risk of nasopharyngeal carcinoma in North Africa. *G3 (Bethesda)* 3(6): 971-977.
130. Mueller N, Evans A, Harris NL, Comstock GW, Jellum E, Magnus K, Orentreich N, Polk BF, Vogelman J. 1989. Hodgkin's disease and Epstein-Barr virus. Altered antibody pattern before diagnosis. *N Engl J Med* 320(11): 689-695.
131. Mulama DH, Bailey JA, Foley J, Chelimo K, Ouma C, Jura WG, Otieno J, Vulule J, Moormann AM. 2014. Sick cell trait is not associated with endemic Burkitt lymphoma: an ethnicity and malaria endemicity-matched case-control study suggests factors controlling EBV may serve as a predictive biomarker for this pediatric cancer. *Int J Cancer* 134(3): 645-653.
132. Munoz N, Davidson RJ, Witthoff B, Ericsson JE, De-The G. 1978. Infectious mononucleosis and Hodgkin's disease. *Int J Cancer* 22(1): 10-13.
133. Murphy G, Pfeiffer R, Camargo MC, Rabkin CS. 2009. Meta-analysis shows that prevalence of Epstein-Barr virus-positive gastric cancer differs based on sex and anatomic location. *Gastroenterology* 137(3): 824-833.
134. Musacchio JG, Carvalho Mda G, Morais JC, Silva NH, Scheliga A, Romano S, Spector N. 2006. Detection of free circulating Epstein-Barr virus DNA in plasma of patients with Hodgkin's disease. *Sao Paulo Med J* 124(3): 154-157.
135. Mutalima N, Molyneux E, Jaffe H, Kamiza S, Borgstein E, Mkandawire N, Liomba G, Batumba M, Lagos D, Gratrix F, Boshoff C, Casabonne D, Carpenter LM, Newton R.

2008. Associations between Burkitt lymphoma among children in Malawi and infection with HIV, EBV and malaria: results from a case-control study. *PLoS One* 3(6): e2505.
136. Newton R, Carpenter L, Casabonne D, Beral V, Babiker A, Darbyshire J, Weller I, Weiss R, Kwan A, Bourbouli D, Munoz F, Lagos D, Boshoff C. 2006. A prospective study of Kaposi's sarcoma-associated herpesvirus and Epstein-Barr virus in adults with human immunodeficiency virus-1. *Br J Cancer* 94(10): 1504-1509.
137. Ning JP, Yu MC, Wang QS, Henderson BE. 1990. Consumption of salted fish and other risk factors for nasopharyngeal carcinoma (NPC) in Tianjin, a low-risk region for NPC in the People's Republic of China. *J Natl Cancer Inst* 82(4): 291-296. (as cited in IARC 1997)
138. Nkrumah F, Henle W, Henle G, Herberman R, Perkins V, Depue R. 1976. Burkitt's lymphoma: its clinical course in relation to immunologic reactivities to Epstein-Barr virus and tumor-related antigens. *J Natl Cancer Inst* 57(5): 1051-1056.
139. Nong LG, Luo B, Zhang L, Nong HB. 2009. Interleukin-18 gene promoter polymorphism and the risk of nasopharyngeal carcinoma in a Chinese population. *DNA Cell Biol* 28(10): 507-513.
140. Pathmanathan R, Prasad U, Sadler R, Flynn K, Raab-Traub N. 1995. Clonal proliferations of cells infected with Epstein-Barr virus in preinvasive lesions related to nasopharyngeal carcinoma. *N Engl J Med* 333(11): 693-698.
141. Pietersma F, Piriou E, van Baarle D. 2008. Immune surveillance of EBV-infected B cells and the development of non-Hodgkin lymphomas in immunocompromised patients. *Leuk Lymphoma* 49(6): 1028-1041.
142. Piriou E, Asito AS, Sumba PO, Fiore N, Middeldorp JM, Moormann AM, Ploutz-Snyder R, Rochford R. 2012. Early age at time of primary Epstein-Barr virus infection results in poorly controlled viral infection in infants from Western Kenya: clues to the etiology of endemic Burkitt lymphoma. *J Infect Dis* 205(6): 906-913.
143. Poirier S, Bouvier G, Malaveille C, Ohshima H, Shao YM, Hubert A, Zeng Y, de The G, Bartsch H. 1989. Volatile nitrosamine levels and genotoxicity of food samples from high-risk areas for nasopharyngeal carcinoma before and after nitrosation. *Int J Cancer* 44(6): 1088-1094. (as cited in IARC 1997)
144. Ponce RA, Gelzleichter T, Haggerty HG, Heidel S, Holdren MS, Lebrec H, Mellon RD, Pallardy M. 2014. Immunomodulation and lymphoma in humans. *J Immunotoxicol* 11(1): 1-12.
145. Pope JH, Horne MK, Scott W. 1968. Transformation of foetal human leukocytes in vitro by filtrates of a human leukaemic cell line containing herpes-like virus. *Int J Cancer* 3(6): 857-866.
146. Raab-Traub N, Rajadurai P, Flynn K, Lanier AP. 1991. Epstein-Barr virus infection in carcinoma of the salivary gland. *J Virol* 65(12): 7032-7036.

147. Raab-Traub N. 2002. Epstein-Barr virus in the pathogenesis of NPC. *Semin Cancer Biol* 12(6): 431-441.
148. Rigaud S, Fondaneche MC, Lambert N, Pasquier B, Mateo V, Soulas P, Galicier L, Le Deist F, Rieux-Laucat F, Revy P, Fischer A, de Saint Basile G, Latour S. 2006. XIAP deficiency in humans causes an X-linked lymphoproliferative syndrome. *Nature* 444(7115): 110-114.
149. Robbiani DF, Deroubaix S, Feldhahn N, Oleveira TY, Callen E, Wang Q, Jankovic M, Silva IT, Rommel PC, Bosque D, Eisenreich T, Nessenzweig A, Nussenzweig MC. 2015. Plasmodium infection promotes genomic instability and AID-dependent B cell lymphoma. *Cell* 162(4): 727-737.
150. Rosdahl N, Larsen SO, Clemmesen J. 1974. Hodgkin's disease in patients with previous infectious mononucleosis: 30 years' experience. *Br Med J* 2(5913): 253-256.
151. Rothschild G, Nikolai von Krusenstiern A, Basu U. 2015. Malaria-induced B cell genomic instability. *Cell* 162(4): 697-698.
152. Rowe M, Young LS, Crocker J, Stokes H, Henderson S, Rickinson AB. 1991. Epstein-Barr virus (EBV)-associated lymphoproliferative disease in the SCID mouse model: implications for the pathogenesis of EBV-positive lymphomas in man. *J Exp Med* 173(1): 147-158.
153. SEER. 2015. *SEER Stat Fact Sheets: Hodgkin Lymphoma*. Surveillance, Epidemiology, and End Results Program. <http://seer.cancer.gov/statfacts/html/hodg.html>. Accessed on 8/27/15.
154. Serraino D, Franceschi S, Talamini R, Barra S, Negri E, Carbone A, La Vecchia C. 1991. Socio-economic indicators, infectious diseases and Hodgkin's disease. *Int J Cancer* 47(3): 352-357.
155. Shao YM, Poirier S, Ohshima H, Malaveille C, Zeng Y, de The G, Bartsch H. 1988. Epstein-Barr virus activation in Raji cells by extracts of preserved food from high risk areas for nasopharyngeal carcinoma. *Carcinogenesis* 9(8): 1455-1457. (as cited in IARC 1997)
156. Shibata D, Weiss LM. 1992. Epstein-Barr virus-associated gastric adenocarcinoma. *Am J Pathol* 140(4): 769-774. (as cited in IARC 1997)
157. Shinkura R, Yamamoto N, Koriyama C, Shinmura Y, Eizuru Y, Tokunaga M. 2000. Epstein-Barr virus-specific antibodies in Epstein-Barr virus-positive and -negative gastric carcinoma cases in Japan. *J Med Virol* 60(4): 411-416.
158. Shope T, Dechairo D, Miller G. 1973. Malignant lymphoma in cottontop marmosets after inoculation with Epstein-Barr virus. *Proc Natl Acad Sci U S A* 70(9): 2487-2491.
159. Sousa H, Pinto-Correia AL, Medeiros R, Dinis-Ribeiro M. 2008. Epstein-Barr virus is associated with gastric carcinoma: the question is what is the significance? *World J Gastroenterol* 14(27): 4347-4351.

160. Sriamporn S, Vatanasapt V, Pisani P, Yongchaiyudha S, Rungpitarangsri V. 1992. Environmental risk factors for nasopharyngeal carcinoma: a case-control study in northeastern Thailand. *Cancer Epidemiol Biomarkers Prev* 1(5): 345-348. (as cited in IARC 1997)
161. Stefan DC, Wessels G, Poole J, Wainwright L, Stones D, Johnston WT, Newton R. 2011. Infection with human immunodeficiency virus-1 (HIV) among children with cancer in South Africa. *Pediatr Blood Cancer* 56(1): 77-79.
162. Stein H, Delsol G, Pileri SA, Weiss LM, Poppema S, Jaffe ES. 2008. Classical Hodgkin Lymphoma, Introduction. In *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*. Swerdlow SH, Campo E, Harris NL, et al, eds. Lyon, France: IARC Press. 326 pp.
163. Suwiwat S, Pradutkanchana J, Ishida T, Mitarnun W. 2007. Quantitative analysis of cell-free Epstein-Barr virus DNA in the plasma of patients with peripheral T-cell and NK-cell lymphomas and peripheral T-cell proliferative diseases. *J Clin Virol* 40(4): 277-283.
164. Taylor AL, Marcus R, Bradley JA. 2005. Post-transplant lymphoproliferative disorders (PTLD) after solid organ transplantation. *Crit Rev Oncol Hematol* 56(1): 155-167.
165. Thompson MP, Kurzrock R. 2004. Epstein-Barr virus and cancer. *Clin Cancer Res* 10(3): 803-821.
166. Thorley-Lawson DA and Gross A. 2004. Persistence of the Epstein-Barr virus and the origins of associated lymphomas. *New Engl J Med* 350: 1328-1337.
167. Tiwawech D, Srivatanakul P, Karalak A, Ishida T. 2008. Association between EBNA2 and LMP1 subtypes of Epstein-Barr virus and nasopharyngeal carcinoma in Thais. *J Clin Virol* 42(1): 1-6.
168. Torgbor C, Awuah P, Deitsch K, Kalantari P, Duca KA, Thorley-Lawson DA. 2014. A multifactorial role for *P. falciparum* malaria in endemic Burkitt's lymphoma pathogenesis. *PLoS Pathog* 10(5): e1004170.
169. Tsai CC, Chen CL, Hsu HC. 1996. Expression of Epstein-Barr virus in carcinomas of major salivary glands: a strong association with lymphoepithelioma-like carcinoma. *Hum Pathol* 27(3): 258-262.
170. Tsang CM, Deng W, Yip YL, Zeng MS, Lo KW, Tsao SW. 2014. Epstein-Barr virus infection and persistence in nasopharyngeal epithelial cells. *Chin J Cancer* 33(11): 549-555.
171. Tsang WYW, Chan JKC. 2005. Lymphoepithelial carcinoma. In *World Health Organization Classification of Tumours – Pathology and Genetics of Head and Neck Tumours*. Barnes EL, Eveson JW, Reichart P, Sidransky D, eds. Lyon, France: International Agency for Research on Cancer. pp. 85-97. (as cited in IARC 2012)
172. Tsao SW, Tsang CM, To KF, Lo KW. 2015. The role of Epstein-Barr virus in epithelial malignancies. *J Pathol* 235(2): 323-333.

173. Tugizov SM, Berline JW, Palefsky JM. 2003. Epstein-Barr virus infection of polarized tongue and nasopharyngeal epithelial cells. *Nat Med* 9(3): 307-314.
174. Vegso G, Hajdu M, Sebestyen A. 2011. Lymphoproliferative disorders after solid organ transplantation-classification, incidence, risk factors, early detection and treatment options. *Pathol Oncol Res* 17(3): 443-454.
175. Vetsika EK, Callan M. 2004. Infectious mononucleosis and Epstein-Barr virus. *Expert Rev Mol Med* 6(23): 1-16.
176. Wang CP, Chang YL, Ko JY, Lou PJ, Yeh CF, Sheen TS. 2004. Lymphoepithelial carcinoma versus large cell undifferentiated carcinoma of the major salivary glands. *Cancer* 101(9): 2020-2027.
177. Wei YS, Lan Y, Zhang L, Wang JC. 2010. Association of the interleukin-2 polymorphisms with interleukin-2 serum levels and risk of nasopharyngeal carcinoma. *DNA Cell Biol* 29(7): 363-368.
178. Weiss LM. 2000. Epstein-Barr virus and Hodgkin's disease. *Curr Oncol Rep* 2(2): 199-204.
179. Wikipedia. 2015. *Epstein-Barr Virus*. Wikipedia. Last updated on 8/27/15. https://en.wikipedia.org/wiki/Epstein%E2%80%93Barr_virus. Accessed on 8/27/15.
180. Wutzler P, Farber I, Sprossig M, Sauerbrei A, Wutke K, Hoche D, Rudiger KD, Wockel W, Scheibner K. 1983. Antibodies against herpesviruses in patients with Hodgkin's disease. *Arch Geschwulstforsch* 53(5): 417-422.
181. Yau TO, Tang CM, Yu J. 2014. Epigenetic dysregulation in Epstein-Barr virus-associated gastric carcinoma: disease and treatments. *World J Gastroenterol* 20(21): 6448-6456.
182. Young LS, Rickinson AB. 2004. Epstein-Barr virus: 40 years on. *Nat Rev Cancer* 4(10): 757-768.
183. Yu MC, Ho JH, Lai SH, Henderson BE. 1986. Cantonese-style salted fish as a cause of nasopharyngeal carcinoma: report of a case-control study in Hong Kong. *Cancer Res* 46(2): 956-961. (as cited in IARC 1997)
184. Yu MC, Mo CC, Chong WX, Yeh FS, Henderson BE. 1988. Preserved foods and nasopharyngeal carcinoma: a case-control study in Guangxi, China. *Cancer Res* 48(7): 1954-1959. (as cited in IARC 1997)
185. Yu MC, Huang TB, Henderson BE. 1989. Diet and nasopharyngeal carcinoma: a case-control study in Guangzhou, China. *Int J Cancer* 43(6): 1077-1082. (as cited in IARC 1997)
186. Yu MC, Yuan JM. 2002. Epidemiology of nasopharyngeal carcinoma. *Semin Cancer Biol* 12(6): 421-429.
187. Zhang L, Sung JJ, Yu J, Ng SC, Wong SH, Cho CH, Ng SS, Chan FK, Wu WK. 2014. Xenophagy in *Helicobacter pylori*- and Epstein-Barr virus-induced gastric cancer. *J Pathol* 233(2): 103-112.

188. Zhao M, Cai H, Li X, Zheng H, Yang X, Fang W, Zhang L, Wei G, Li M, Yao K, Li X. 2012c. Further evidence for the existence of major susceptibility of nasopharyngeal carcinoma in the region near HLA-A locus in Southern Chinese. *J Transl Med* 10: 57.
189. Zheng X, Yan L, Nilsson B, Eklund G, Drettner B. 1994a. Epstein-Barr virus infection, salted fish and nasopharyngeal carcinoma. A case-control study in southern China. *Acta Oncol* 33(8): 867-872.
190. Zheng YM, Tuppin P, Hubert A, Jeannel D, Pan YJ, Zeng Y, de The G. 1994b. Environmental and dietary risk factors for nasopharyngeal carcinoma: a case-control study in Zangwu County, Guangxi, China. *Br J Cancer* 69(3): 508-514. (as cited in IARC 1997)
191. zur Hausen H. 2001. Oncogenic DNA viruses. *Oncogene* 20(54): 7820-7823.

Glossary

Affinity: A measurement of the strength of interaction between an epitope and an antibody's antigen binding site.

Case report: Detailed descriptions of a few patients or clinical cases (frequently, just one sick person) with an unusual disease or complication, uncommon combinations of diseases, an unusual or misleading semiology, cause, or outcome (maybe a surprising recovery). They often are preliminary observations that are later refuted. They cannot estimate disease frequency or risk (e.g., for lack of a valid denominator).

Case series: A collection of subjects (usually, patients) with common characteristics used to describe some clinical, pathophysiological, or operational aspect of a disease, treatment, exposure, or diagnostic procedure. A case series does not include a comparison group and is often based on prevalent cases and on a sample of convenience. Common selection biases and confounding severely limit their power to make causal inferences.

Case-comparison study (case-control study, case referent study): The observational epidemiological study of persons with the disease (or another outcome variable) of interest and a suitable control group of persons without the disease (comparison group, reference group). The potential relationship of a suspected risk factor or an attribute to the disease is examined by comparing the diseased and non-diseased subjects with regard to how frequently the factor or attribute is present (or, if quantitative, the levels of the attribute) in each of the groups (diseased and non-diseased).

Convenience sample: Samples selected by easily employed but basically non-probabilistic (and probably biased) methods. "Man-in-the-street" surveys and a survey of blood pressure among volunteers who drop in at an examination booth in a public place are in this category.

Defining acquired immune deficiency syndrome (AIDS) condition: Any HIV-related illness included in the Centers for Disease Control and Prevention's (CDC) list of diagnostic criteria for AIDS. AIDS-defining conditions include opportunistic infections and cancers that are life threatening in a person with HIV.

Diagnostic criteria: The specific combination of signs, symptoms, and test results that a clinician uses to identify a person as representing a case of a particular disease or condition.

Enzyme immunoassay: An assay that uses an enzyme-bound antibody to detect antigen. The enzyme catalyzes a color reaction when exposed to substrate.

Exposure variability: A characteristic describing whether epidemiological studies included non-exposed cases or exposed controls. Lack of exposure variability may contribute to observed heterogeneity of study results.

Germinal center: Discrete areas within secondary lymphoid tissues where B-cell maturation

and memory development occur.

***In situ* hybridization:** A technique that allows for precise localization of a specific segment of nucleic acid within a histologic section. The underlying basis of *in situ* hybridization is that nucleic acids, if preserved adequately within a histologic specimen, can be detected through the application of a complementary strand of nucleic acid to which a reporter molecule is attached.

EBV latency patterns: Particular EBV viral gene expression patterns that establish distinct EBV latent infection statuses (latency types III, II, and I).

Latent phase: A phase of the virus life cycle during which the virus is not replicating.

Lytic phase: A phase of the virus life cycle during which the virus replicates within the host cell, releasing a new generation of viruses when the infected cell lyses.

Monoclonal: Pertaining to or designating a group of identical cells or organisms derived from a single cell or organism.

Nude mouse (athymic nude mouse): A type of laboratory mouse that is hairless, lacks a normal thymus gland, and has a defective immune system because of a genetic mutation. Athymic nude mice are often used in cancer research because they do not reject tumor cells, from mice or other species.

Open reading frame: A portion of a DNA molecule that, when translated into amino acids, contains no stop codons.

Optical density: The absorbance of a particular substance at a specified wavelength in an enzyme-linked immunosorbent assay.

Peripheral blood: Blood circulating throughout the body; the primary method for transporting nutrients such as oxygen and carbon dioxide through the body consisting of three primary components: erythrocytes (red blood cells), leukocytes (white blood cells), and thrombocytes (blood platelets).

Polyclonal: Pertaining to or designating a group of cells or organisms derived from several cells.

Titer: A laboratory measurement of the concentration of a substance in a solution (e.g., an antibody titer measures the presence and amount of antibodies in the blood).

Abbreviations

AIDS	Acquired Immune Deficiency Syndrome
BARF1	<i>Bam</i> HI A rightward fragment 1 (micro RNA)
BL	Burkitt lymphoma
CDC	Centers for Disease Control and Prevention
CF/S	complement-fixing soluble antigen
CI	confidence interval
CNS	central nervous system
DNA	deoxyribonucleic acid
EA	early antigen
EA(D)	early antigen-diffuse
EBER	EBV-encoded small RNA
EBNA	EBV nuclear antigen
EBNA-LP	EBV nuclear antigen leader protein
EBV	Epstein-Barr virus
EIA	enzyme immunoassays
GC	germinal center
HHV4	human herpesvirus 4
HIV	human immunodeficiency virus
IARC	International Agency for Research on Cancer
HLA	human leukocyte antigen
IFA	indirect immunofluorescence assay
IgA	immunoglobulin A
IgG	immunoglobulin G
IgM	immunoglobulin M

LEC	lymphoepithelial cancer
LMP	latent membrane protein
LMP-1	latent membrane protein 1
LMP-2	latent membrane protein 2
LMP-2B	latent membrane protein 2B
MALT	mucosa-associated lymphoid tissue
NHL	non-Hodgkin lymphoma
NPC	nasopharyngeal carcinoma
NHANES	National Health and Nutrition Examination Survey
NR	not reported
NS	not statistically significant
NTP	National Toxicology Program
OD	optical density
OR	odds ratio
ORF	open reading frames
PCR	polymerase chain reaction
PI3K	phosphorylated Akt
PTEN	phosphatase and tensin homologue
PTLD	post-transplant lymphoproliferative disorders
Q-PCR	quantitative polymerase chain reaction
RNA	ribonucleic acid
RR	relative risk
RT-PCR	reverse transcriptase-polymerase chain reaction
SCNA	somatic copy-number aberrations
SEER	Surveillance, Epidemiology, and End Results Program

SIR	standardized incidence ratios
USA	United States of America
VCA	viral caspid anigen

This Page Intentionally Left Blank

Appendix A: Literature Search Strategy

The objective of the literature search approach is to identify published literature that is relevant for evaluating the potential carcinogenicity of the Epstein-Barr Virus (EBV). As discussed in the Viruses Concept Document

(https://ntp.niehs.nih.gov/ntp/roc/concept_docs/2014/virusesconcept_508.pdf), the monograph relies on the IARC monograph and studies published since the monograph (new studies). The literature search strategy was used to identify new human cancer studies and recent reviews of mechanistic data.

General approach

Database searching encompasses selecting databases and search terms and conducting the searches. Searches of several citation databases are generally conducted using search terms for the individual viruses of interest, combined with search terms for cancer and/or specific topics, including epidemiological and mechanistic studies. A critical step in the process involves consultation with an information specialist to develop relevant search terms. These terms are used to search bibliographic databases. IARC used literature found by searching PubMed for EBV through 12/2008, so PubMed, Web of Science and Scopus were searched for new information about EBV from >2008 to August 2015. Table 1 highlights the general concepts searched with selected example terms. To review all the terms used, please refer to the full search strings below.

Table A-1. Major topics searched

Topics	Example terms
Epstein-Barr virus *NOT post-transplant	Epstein-Barr virus infections(Mesh), EBV infection, Herpes virus 4, Herpesvirus 4, human(Mesh)
General Cancer Relevant cancers	<u>Neoplasms(MESH), Tumor(s), leukemia, cancer</u> gastric cancers, Stomach neoplasms(Mesh), parotid, Salivary gland neoplasms(Mesh), lymphoepithel, LELC
Study types	case control, ecological studies, follow-up study
Epidemiology Terms	cohort, Epidemiologic Studies (Mesh), epidemiology (Subheading)

The large and complex body of literature for EBV was searched using narrowing terms for the relevant major topics within the bibliographic databases. The results were then processed in EndNote to remove duplicates and conduct a first level of screening, before being transferred to DistillerSR for additional screening.




Figure A-1. Literature processing flow

The bibliographic database search results (8438) were processed in Endnote then imported into DistillerSR for first and second tier screening. Relevant studies found through the citations of review articles and other secondary searched were also included. Tagging in DistillerSR categorized the useful articles into Human Epidemiologic literature (108) or Mechanistic literature (22).

Search strings for EBV Searches

Cancer and Epidemiology

PubMed: 2009-2015

(Epstein-Barr virus infections[mh] OR "epstein-barr virus nuclear antigens"[mh] OR "epstein-barr"[tiab] OR EBV[tiab] OR (EBV*[tiab] NOT "epstein-barr"[tiab]) OR EBVaGC[tiab] OR Herpesvirus 4, human[mh] OR "Herpes virus 4"[tiab] OR HHV-4[tiab] OR HHV4[tiab] OR "EBV infection" OR EBV-infected OR "EBV-associated" OR "Ebv-encoded" OR "EBV-driven" OR "EBV-positive") NOT (Posttransplant OR post-transplant OR transplant*[tiab])

AND

Neoplasms[mh] OR neoplas*[tiab] OR tumor[tiab] OR tumors[tiab] OR tumorigenesis[tiab] OR tumour*[tiab] OR cancer*[tiab] OR carcinogen*[tiab] OR lesion*[tiab] OR adenoma*[tiab] OR adenocarcinoma*[tiab] OR leiomyo*[tiab] OR leukemia*[tiab] OR lymphoma*[tiab] OR lymphangi*[tiab] OR sarcoma*[tiab] OR carcinoma*[tiab] OR adenocarcinoma*[tiab] OR oncogen*[tiab]

AND

Epidemiologic studies[mh] OR epidemiology[sh] OR case-control[tiab] OR case reports[ptyp] OR clinical trial[ptyp] OR cohort[tiab] OR comparative study[ptyp] OR cross-sectional[tiab] OR evaluation studies[ptyp] OR “follow-up study”[tiab] OR longitudinal[tiab] OR meta-analysis[tiab] OR meta-analysis[ptyp] OR multicenter study[ptyp] OR observational-study[tiab] OR prospective[tiab] OR randomized controlled trial[tiab] OR retrospective[tiab] OR individual*[tiab] OR man[tiab] OR men[tiab] OR patient*[tiab] OR subject*[tiab] OR woman[tiab] OR women[tiab] OR cases[mh]

Web of Science and Scopus: 2009-2015

("epstein-barr" OR EBV* OR EBVaGC OR "Herpes virus 4" OR HHV-4 OR HHV4 OR EBV-infected OR EBV-infection OR EBV-associated OR EBV-encoded OR EBV-driven OR EBV-positive) NOT (Posttransplant OR post-transplant OR transplant*)

AND

neoplas* OR tumor* OR tumour* OR cancer* OR carcinogen* OR lesion* OR adenoma* OR adenocarcinoma* OR leiomyo* OR leukemia* OR lymphoma* OR lymphangio* OR sarcoma* OR carcinoma* OR adenocarcinoma* OR oncogen*

AND

Epidemiologic* OR case-control OR “case report” OR “case reports” OR “clinical trial” OR cohort OR “comparative study” OR cross-sectional OR “evaluation studies” OR “follow-up study” OR longitudinal OR meta-analysis OR multicenter study OR observational-study OR prospective OR “randomized controlled trial” OR retrospective OR individual* OR man OR men OR patient* OR subject* OR woman OR women

Relevant Cancers

PubMed: 2009-2015

- A. (Epstein-Barr virus infections[mh] OR "epstein-barr virus nuclear antigens"[mh] OR "epstein-barr"[tiab] OR EBV[tiab] OR (EBV*[tiab] NOT "epstein-barr"[tiab]) OR EBVaGC[tiab] OR Herpesvirus 4, human[mh] OR "Herpes virus 4"[tiab] OR HHV-4[tiab] OR HHV4[tiab] OR “EBV infection” OR EBV-infected OR "EBV-associated" OR "Ebv-encoded" OR "EBV-driven" OR “EBV-positive”) NOT (Posttransplant OR post-transplant OR transplant*[tiab])

AND

Stomach neoplasms[mh] OR ((gastric OR stomach) AND (adenocarcinoma* OR adenoma*[tiab] OR adenocarcinoma*[tiab] OR cancer*[tiab] OR carcinogen* OR carcinoma* OR lesion*[tiab] OR lymphoid[tiab] OR lymphoma*[tiab] OR lymphoepithel*[tiab] OR neoplas* OR tumor[tiab] OR tumors[tiab] OR tumorigenesis OR tumour*))OR stomach/virology[mh] OR ((stomach OR gastric) AND (epithelial*[tiab] OR epithelial cells/virology[mh])) OR (gammaherpesvirus-68[tiab] OR gammaHV-68[tiab]) AND (stomach[tiab] OR intestin*[tiab] OR gastric[tiab] OR gut[tiab])

- B. (Epstein-Barr virus infections[mh] OR "epstein-barr virus nuclear antigens"[mh] OR "epstein-barr"[tiab] OR EBV[tiab] OR (EBV*[tiab] NOT "epstein-barr"[tiab]) OR EBVaGC[tiab] OR

Herpesvirus 4, human[mh] OR "Herpes virus 4"[tiab] OR HHV-4[tiab] OR HHV4[tiab] OR "EBV infection" OR EBV-infected OR "EBV-associated" OR "Ebv-encoded" OR "EBV-driven" OR "EBV-positive") NOT (Posttransplant OR post-transplant OR transplant*[tiab])

AND

Salivary gland neoplasms[mh] OR salivary glands[mh] OR salivary-gland*[tiab] OR parotid[tiab] OR sublingual[tiab] OR submandibular[tiab] OR salivary-duct*[tiab] OR "von Ebner"[tiab] AND (Lymphoproliferative OR lymphoepithel* OR LEC OR LELC)

- C. (Epstein-Barr virus infections[mh] OR "epstein-barr virus nuclear antigens"[mh] OR "epstein-barr"[tiab] OR EBV[tiab] OR (EBV*[tiab] NOT "epstein-barr"[tiab]) OR EBVaGC[tiab] OR Herpesvirus 4, human[mh] OR "Herpes virus 4"[tiab] OR HHV-4[tiab] OR HHV4[tiab] OR "EBV infection" OR EBV-infected OR "EBV-associated" OR "Ebv-encoded" OR "EBV-driven" OR "EBV-positive") NOT (Posttransplant OR post-transplant OR transplant*[tiab])

AND

lymphoepithel*[tiab] OR LEC[tiab] OR LELC[tiab]

Web of Science and Scopus: 2009-2015

- A. ("epstein-barr" OR EBV* OR EBVaGC OR "Herpes virus 4" OR HHV-4 OR HHV4 OR EBV-infected OR EBV-infection OR EBV-associated OR EBV-encoded OR EBV-driven OR EBV-positive) NOT (Posttransplant OR post-transplant OR transplant*)

AND

((gastric OR gut OR intestin* OR stomach) AND (adenocarcinoma* OR cancer* OR carcinogen* OR carcinoma* OR neoplas* OR tumor* OR tumour*)) OR ((gammaherpesvirus-68 OR gammaHV-68) AND (stomach OR intestin* OR gastric OR gut))

- B. ("epstein-barr" OR EBV* OR EBVaGC OR "Herpes virus 4" OR HHV-4 OR HHV4 OR EBV-infected OR EBV-infection OR EBV-associated OR EBV-encoded OR EBV-driven OR EBV-positive) NOT (Posttransplant OR post-transplant OR transplant*)

AND

"Salivary gland" OR "salivary glands" OR parotid OR sublingual OR submandibular OR "salivary duct" OR "salivary ducts" OR "von Ebner" AND (Lymphoproliferative OR lymphoepithel* OR LEC OR LELC)

- C. ("epstein-barr" OR EBV* OR EBVaGC OR "Herpes virus 4" OR HHV-4 OR HHV4 OR EBV-infected OR EBV-infection OR EBV-associated OR EBV-encoded OR EBV-driven OR EBV-positive) NOT (Posttransplant OR post-transplant OR transplant*)

AND

lymphoepithel* OR LEC OR LELC

Part 2

Draft Profile

This Page Intentionally Left Blank

Epstein-Barr Virus

CAS No.: none assigned

Known to be a human carcinogen²

Also known as EBV or human herpesvirus 4 (HHV-4)

Carcinogenicity

Epstein-Barr virus (EBV) is *known to be a human carcinogen* based on sufficient evidence from studies in humans. This conclusion is based on evidence from epidemiological, clinical, and molecular studies, which show that Epstein-Barr virus causes endemic Burkitt lymphoma, Hodgkin lymphoma, immune-suppression-related non-Hodgkin lymphoma, extranodal natural killer (NK)/T-cell lymphoma nasal type, nasopharyngeal carcinoma, and some forms of stomach cancer. There is also limited evidence for an association with sporadic Burkitt lymphoma and lymphoepithelial cancer of the salivary gland.

Results from clinical and mechanistic studies indicate that transcription programs activated during viral latency mimic signals that induce B-cell proliferation and survival, resulting in cancer in some cases. Furthermore, in most EBV-related cancers, a large proportion of tumor cells contain the virus, and it is monoclonal, suggesting that the viral infection preceded the tumor. In addition, it has been shown that the Epstein-Barr nuclear antigen (EBNA) proteins EBNA-1, -2, -3A, and -3C and latent membrane protein 1 (LMP-1) all are necessary for immortalization of B lymphocytes.

Cancer Studies in Humans

EBV was the first oncogenic human virus to be discovered; its association with Burkitt lymphoma was recognized over 50 years ago (Epstein *et al.* 1964). Since then, numerous studies have explored the relationship of EBV to various types of cancer. The majority of studies have focused on Burkitt lymphoma, Hodgkin lymphoma, nasopharyngeal cancer, and some forms of non-Hodgkin lymphoma, as well as less common lymphomas and lymphoepithelial cancers.

Burkitt Lymphoma

Burkitt lymphoma has three subtypes: (1) endemic, which occurs primarily in children in equatorial Africa and Papua, New Guinea, (2) sporadic, which is found throughout the world, and (3) immunodeficiency-related. The body of analytical epidemiological studies on Burkitt lymphoma consists of twelve case-control studies (seven of which reported risk estimates, with a total of 904 cases), one cohort study of the endemic type, and five case-control studies of the sporadic type, with a total of 113 cases (see section 3, Human Cancer Studies, Cancer Hazard Evaluation Component, IARC 1997, 2012, Mulama *et al.* 2014). Most of the studies measured antibodies to EBV antigens (such as viral capsid antigen [VCA] and early antigen) in sera; one study (Mulama *et al.* 2014) measured EBV DNA isolated from plasma and peripheral blood cells.

There is a credible evidence for an association between EBV infection and endemic Burkitt lymphoma, based on consistent findings of increased risk in several case-control and cohort studies and evidence of a dose-response relationship with viral infection. All of the studies of endemic Burkitt lymphoma and EBV infection found statistically significant positive

²NTP preliminary listing recommendation proposed for the RoC.

relationships, with risks ranging from 2.9 to 52.1. Moreover, several of the case-control studies and the cohort study found increasing risk of endemic Burkitt lymphoma with increasing viral titers (primarily by measurement of antibodies to VCA) (Henle *et al.* 1969, 1971, Carpenter *et al.* 2008, Mutalima *et al.* 2008). Evidence for an association between EBV infection and sporadic EBV is somewhat weaker: positive associations were found in four of the five case-control studies, one of which was statistically significant, and no studies evaluated dose-response relationships. These studies had limited power to detect an effect because of the small numbers of EBV-infected case and control subjects.

These findings are supported by molecular and clinical studies in humans. Evidence of monoclonal EBV infection is found in approximately 95% of endemic Burkitt lymphomas (Thompson and Kurzrock 2004) and EBV infection is found in approximately 20% of sporadic Burkitt lymphomas (IARC 1997). All forms of Burkitt lymphoma are characterized by dysregulation of the gene *c-myc* by chromosomal translocation to an immunoglobulin promoter, enhancing B-cell proliferation. In all Burkitt lymphomas, most cells express only the EBNA-1 protein. Normally, defective B cells exiting the germinal center undergo apoptosis; however, it has been shown from *in vitro* studies that viral expression of EBNA-1 rescues defective B cells from apoptosis by directly enhancing production of the apoptosis-inhibiting protein survivin; EBNA-1 complexes with Sp1, a transcription factor, on the *survivin* gene promoter (Lu *et al.* 2011).

Malarial infection has been shown to be a cofactor for endemic Burkitt lymphoma, increasing risk synergistically, in part by inducing a strong and long-lasting expansion of germinal center B cells (Robbiani *et al.* 2015), reducing T-cell-mediated immunosurveillance and increasing viral load of EBV-infected cells (Moormann *et al.* 2009, as cited in IARC 2012)

Hodgkin Lymphoma

There is credible evidence for an association between EBV infection and Hodgkin lymphoma, based on consistent findings of statistically significant increased risk in numerous case-control and cohort studies. The database of epidemiological studies consists of over 45 case-control and cohort or nested case-control studies that measured EBV infection either directly via serology or DNA or indirectly via history of infectious mononucleosis (IARC 1997, 2012, Linabery *et al.* 2014). The strongest evidence for an association comes from the collective findings of over 20 case-control studies measuring EBV infection directly. The majority of the studies that measured VCA titer reported an association between high VCA titer and Hodgkin lymphoma, with odds ratios between 4 and 19. Strong associations were also observed in studies of EBV DNA or antibodies to other viral agents. Most of the case-control studies on infectious mononucleosis and cohort studies on either infectious mononucleosis or EBV serology also found significantly increased risk, though of lower magnitude (generally between 1.3 and 5). A few studies did not find a positive association, and not all studies found significantly elevated risk. However, the strength of the database (i.e., the large number of studies using different study designs, conducted in different geographical locations, and measuring EBV by different methods) and the consistent findings of statistically significant risk of relatively large magnitude, provide evidence of a causal association between EBV infection and Hodgkin lymphoma.

These findings are supported by molecular studies of human tissue or cell lines. Some forms of Hodgkin lymphoma are associated with monoclonal EBV infection. Hodgkin Reed-Sternberg cells are cancerous lymphocytes that contain EBV. Viral LMP-1 and -2A promote growth and survival of Hodgkin Reed-Sternberg cells through enhancement of multiple growth

and survival signaling pathways, such as the NF- κ B, JAK/STAT, MAP kinase, and PI3-kinase/AKT pathways, causing an abnormal pattern of release of cytokines and chemokines resulting in a local inflammatory response (IARC 1997, 2012, Mohamed *et al.* 2014).

Immune-Suppression-Related Non-Hodgkin Lymphoma

Severe immunosuppression from congenital, iatrogenic, or HIV-related immunodeficiency can result in EBV-associated non-Hodgkin lymphoma. Evidence that these lymphomas are associated with EBV infection comes primarily from case-series and clinical studies demonstrating the presence of EBV in these lymphomas. EBV has been found in all case reports of lymphoma in patients with congenital immune deficiency, almost all case reports of non-Hodgkin lymphoma of the central nervous system in HIV patients, and 50% of patients with post-transplant lymphoproliferative disease (IARC 1997, 2012). Furthermore, EBV DNA was detected in the plasma of patients with post-transplant lymphoproliferative disease but not in 35 healthy control subjects (Lei *et al.* 2000). A nested case-control study provided limited evidence for an association between EBV and non-Hodgkin lymphoma in HIV-positive patients, based on titer of antibodies to VCA (Newton *et al.* 2006). Other studies in humans have shown that EBV in the tumors is monoclonal and produces oncogenic proteins (LMP-1, -2A, and -2B and EBNA) that promote cell division, cell survival, and oncogenic transformation. Finally, treatment of immune-suppressed patients with T cells sensitized to EBV has been shown to reduce viral load and reduce or protect against the formation of this tumor (Taylor *et al.* 2005, Vegso *et al.* 2011, IARC 2012).

Extranodal NK/T Cell Lymphoma, Nasal-Type

Extranodal NK/T-cell lymphoma, nasal type is a rare type of non-Hodgkin lymphoma of NK-cell neoplasms; however, some cases show a cytotoxic T-cell phenotype. Evidence that this type of lymphoma is associated with EBV infection comes primarily from more than a dozen case-series studies, including over 400 cases (IARC 1997, Barrionuevo *et al.* 2007, He *et al.* 2007), showing its detection in 100% of these tumors, and from other clinical and molecular studies in humans. Two case-comparison studies found EBV DNA in the plasma or CD3+ (T) cells from case subjects but not from control subjects (Lei *et al.* 2000, Suwiwat *et al.* 2007). Other studies in humans have shown that EBV is monoclonal in the tumors and produces oncogenic proteins (LMP-1 and -2A and EBNA-1) that promote cell division, survival, and oncogenic transformation (IARC 2012).

Nasopharyngeal Carcinoma

There is credible evidence for an association between EBV infection and nasopharyngeal carcinoma, based on consistent findings of highly increased risk in numerous case-control and cohort studies. The database of epidemiological studies consists of over 20 case-control and cohort or nested case-control studies that measured antibodies to EBV antigens (VCA and early antigen), EBV DNase, or EBV DNA (IARC 1997, 2012). The strongest evidence comes from the collective findings of 11 case-control studies, all of which reported statistically significant increased risks, with most estimates ranging from 21 to 138 in the serology studies and 41 to 820 in the DNA or DNase studies. Increased risks were also found in two cohort studies (totaling 168 cases with up to 16 years of follow-up) but not in two small nested case-control studies in the United States (where nasopharyngeal cancer is rare, one of the two studies was among Alaska Natives). Some studies did not calculate risk estimates but reported statistically significant higher

mean titers of anti-EBV in case subjects than in control subjects. Most of these studies were conducted in Asia, but a few also included cases from Europe, North Africa, and the United States.

These findings are supported by data from molecular studies. A monoclonal form of EBV is associated with precancerous lesions and with 98% of non-keratinizing nasopharyngeal carcinomas (Pathmanathan *et al.* 1995, Liu *et al.* 2011, Tsang *et al.* 2014, Tsao *et al.* 2015), and EBV has been shown to produce oncogenic proteins (LMP-1 and -2A and EBNA-1) that promote cell division, cell survival, and oncogenic transformation (Raab-Traub 2002).

Stomach (Gastric) Cancer

Epidemiological, case-series, clinical, and molecular studies provide credible evidence of an association between EBV infection and a specific type of stomach cancer; EBV is found in 8% to 11% of stomach tumors, and EBV-related tumors are more likely to originate in the gastric cardia (where the contents of the esophagus empty into the stomach), corpus (body), or postsurgical gastric stump remnants than are other types of stomach tumors. The available epidemiological studies consist of three case-control studies (Shinkura *et al.* 2000, Lo *et al.* 2001, De Aquino *et al.* 2012), including 184 cases of EBV-related cancer, and three nested case-control studies, primarily in Asian countries, which measured either antibodies to viral proteins or EBV DNA (Levine *et al.* 1995, Koshiol *et al.* 2007, Kim *et al.* 2009). Positive associations were found in the three case-control studies and in two of the nested case-control studies (Levine *et al.* 1995, Kim *et al.* 2009), but the results were not statistically significant in the nested case-control studies.

Molecular studies provide strong evidence of an association between EBV and some stomach cancers. These studies show that EBV is found as a monoclonal form in a subset of human stomach cancers and produces oncogenic proteins in stomach-cancer tissue (LMP-1 and -2A and EBNA-1) that promote cell division, cell survival, and oncogenic transformation. EBV produces a unique molecular profile in genomic studies, characterized by extreme DNA hypermethylation and methylation of CDKN2A+, deregulating the *p16* and *p14* tumor-suppressor genes. EBV-infected stomach tumors also display recurrent *PIK3CA* mutations and amplification of *JAK2*, *CD274* (also known as *PD-L1*), and *PDCD1LG2* (also known as *PD-L2*) (Cancer Genome Atlas Research Network 2014, Gulley 2015).

Lymphoepithelial Cancer of the Salivary Gland

Lymphoepithelial carcinoma of the salivary gland is rare; histologically similar to nasopharyngeal carcinoma, it accounts for less than 1% of all head and neck cancers (Tsai *et al.* 1996, Tsang and Chan 2005). Overall, case reports, case series, and clinical studies provide limited evidence for causal association between EBV infection and lymphoepithelial carcinoma of the salivary gland. Across the body of case-series studies, EBV DNA was detected in tumors in almost all of the over 200 cases of parotid salivary gland lymphoepithelial carcinoma (IARC 1997, 2012). A study comparing EBV infection in different types of salivary-gland cancer found EBV DNA present in the tumors of all 16 salivary-gland lymphoepithelial carcinoma patients but not in any other salivary-gland tumors (Wang *et al.* 2004). Other studies found a monoclonal form of EBV in 100% of the lymphoepithelial cancers tested but not in normal adjacent tissue; however, only a few samples were tested. EBV proteins associated with cell proliferation, cell survival, and oncogenesis were produced; EBNA-1, LMP-1, and *BamHI*-A rightward reading frame were expressed in the malignant cells (Raab-Traub *et al.* 1991, IARC 2012). These data,

although limited, indicate an oncogenic pattern of clonal EBV in these cancers. No additional supporting mechanistic data specific for this cancer were identified.

Studies on Mechanisms of Carcinogenesis

Direct evidence that EBV causes lymphoma comes from studies of human cells *in vitro* and of mice *in vivo*. EBV has been shown to transform lymphoblastoid cells in culture and can transform epithelial cells when co-cultured with transformed lymphoblastoid cells (Imai *et al.* 1998). In addition, lymphoblastoid cells transfected with activated *c-myc* genes were tumorigenic in nude mice, and infected B cells were shown to cause B-cell lymphoma in severe combined immunodeficient mice (Lombardi *et al.* 1987, Mosier *et al.* 1989, Rowe *et al.* 1991).

Biological Properties

Epstein-Barr virus is an enveloped double-stranded DNA gamma-1 herpesvirus (IARC 2012). The 172-kb EBV genome encodes over 85 genes, categorized as latent or lytic. The two major types of EBV (EBV-1 and EBV-2) differ in the gene sequences of their nuclear antigens (EBNA-2, -3A, -3B), and each type has several strains. Infection of epithelial cells is primarily lytic (resulting in destruction of the infected cell), whereas infection of B cells is primarily latent. Antibody-producing B cells, also known as plasma cells, allow EBV to enter the lytic phase and replicate (Ponce *et al.* 2014). EBV proteins expressed on the membranes of infected cells can be recognized by cytotoxic T cells and NK cells, which attack and destroy them (Thorley-Lawson and Gross 2004). However, EBV in latency 0 phase, typically found in resting memory B cells, does not express proteins on infected cells, allowing EBV to evade immune recognition.

Detection

EBV infection can be detected by measuring anti-EBV antibodies in serum or EBV DNA or RNA in peripheral white blood cells (IARC 2012). Measurement of EBV DNA or RNA can indicate viral load, reactivation, response to treatment, and presence in tumor cells. DNA and RNA detection methods include quantitative polymerase chain reaction (PCR), reverse transcriptase PCR, or *in situ* hybridization. Healthy carriers of EBV in developed countries do not have detectable DNA or RNA in cell-free serum, so positive results indicate EBV-associated disease or EBV reactivation.

Exposure

Prevalence studies measuring antibodies to EBV have shown that a significant number of people in the United States are exposed to EBV.

Transmission

Transmission of EBV is primarily via saliva (IARC 2012). The presence of EBV in peripheral blood suggests that transmission via blood is also possible, and transmission has been reported among transfusion and organ transplant recipients. Infected cells, primarily resting memory B cells in peripheral blood, provide a permanent reservoir from which progeny viruses can disseminate within the body and infect other hosts.

Seroprevalence Studies

EBV seroprevalence in the United States, based on National Health and Nutrition Examination Survey data collected in 2009 and 2010, ranged from 50% in 6- to 8-year-olds to 89% in 18- to 19-year-olds (Balfour *et al.* 2013, Dowd *et al.* 2013). Lower EBV seroprevalence has been shown to be associated with higher socioeconomic status within race and ethnicity groups. Analysis of 782 serum samples from Minnesota children aged 18 months to 19.9 years indicated that a combination of genetics, family practices, and home environment were responsible for racial and ethnic differences in EBV prevalence among young children; it was noted that the route of EBV transmission to preadolescents remains unclear (Condon *et al.* 2014).

More than 90% of adults worldwide are infected with EBV (IARC 2012). Age at primary infection varies, occurring at a higher rate during infancy in developing countries than in developed countries, perhaps as a result of better hygienic conditions and other socioeconomic and demographic factors that result in later age of exposure to infected saliva (e.g., household size and population density) (Biggar *et al.* 1978a,b, IARC 2012, Piriou *et al.* 2012, Dowd *et al.* 2013).

Diseases (Non-Cancer), Prevention, and Treatment

Most individuals are infected with EBV but remain otherwise healthy and asymptomatic (IARC 2012). Infection is lifelong and is subclinical when it occurs in early childhood (IARC 2012); however, it results in infectious mononucleosis in at least 25% of infected teenagers and young adults (CDC 2014b). Oral hairy leukoplakia results from infection with EBV in the context of immunosuppression (e.g., caused by HIV-1) or immunosenescence (from aging) (Auwaerter 2015). Chronic active EBV occurs frequently in Asia and South America, but rarely in the United States and Europe. Its etiology is unknown, but is believed to involve rare genetic abnormalities that impair immune control of EBV infection (Rigaud *et al.* 2006, Cohen 2009, Chaigne-Delalande *et al.* 2013).

EBV transmission is associated with EBV shedding in saliva; therefore, avoiding salivary exposure (e.g., via kissing or sharing food, drink, or toothbrushes) may theoretically prevent transmission (CDC 2014a).

Some drugs have been reported to reduce or inhibit EBV shedding (e.g., see Auwaerter 2015); however, currently there are no FDA-approved drugs for treatment of EBV infection. There is no vaccine against EBV, but vaccine development efforts are ongoing (Balfour 2014, ACS 2015, CDC 2015, Cohen 2015, FDA 2015b).

Regulations

Department of Transportation (DOT)

Infectious substances are considered hazardous materials, and special requirements have been set for marking, labeling, and transporting these materials.

Food and Drug Administration (FDA)

21 CFR 866 identifies Epstein-Barr virus serological reagents (i.e., devices that consist of antigens and antisera used in serological tests to identify antibodies to EBV in serum) as Class I medical devices requiring premarket notification for FDA clearance to market.

Occupational Safety and Health Administration (OSHA)

Comprehensive regulations have been developed for employers to develop and adhere to exposure control plans for bloodborne pathogens.

All work-related needlestick injuries and cuts from sharp objects that are contaminated with another person's blood or other potentially infectious material must be recorded.

First-aid training program trainees must have adequate instruction in the value of universal precautions for preventing infectious diseases.

Guidelines

Food and Drug Administration (FDA)

The FDA has issued numerous guidance documents prescribing procedures (e.g., use of standardized labels, abbreviated donor screening questionnaires) for reducing the risk of virus transmission by blood and blood products (FDA 2015a).

References

- ACS. 2015. *Viruses that Can Lead to Cancer*. American Cancer Society. <http://www.cancer.org/cancer/cancercauses/othercarcinogens/infectiousagents/infectiousagentsandcancer/infectious-agents-and-cancer-viruses>. Last updated: 4/27/15.
- Auwaerter PG. 2015. *Epstein-Barr Virus*. Johns Hopkins HIV Guide. Johns Hopkins Medicine. http://www.hopkinsguides.com/hopkins/view/Johns_Hopkins_HIV_Guide/545067/all/Epstein_Barr_virus. Last updated: 6/2/2015.
- Balfour HH Jr, Sifakis F, Sliman JA, Knight JA, Schmeling DO, Thomas W. 2013. Age-specific prevalence of Epstein-Barr virus infection among individuals aged 6-19 years in the United States and factors affecting its acquisition. *J Infect Dis* 208(8): 1286-1293.
- Balfour HH Jr. 2014. Progress, prospects, and problems in Epstein-Barr virus vaccine development. *Curr Opin Virol* 6: 1-5.
- Barrionuevo C, Zaharia M, Martinez MT, Taxa L, Misad O, Moscol A, *et al.* 2007. Extranodal NK/T-cell lymphoma, nasal type: study of clinicopathologic and prognosis factors in a series of 78 cases from Peru. *Appl Immunohistochem Mol Morphol* 15(1): 38-44.
- Biggar RJ, Henle W, Fleisher G, Bocker J, Lennette ET, Henle G. 1978a. Primary Epstein-Barr virus infections in African infants. I. Decline of maternal antibodies and time of infection. *Int J Cancer* 22(3): 239-243.
- Biggar RJ, Henle G, Bocker J, Lennette ET, Fleisher G, Henle W. 1978b. Primary Epstein-Barr virus infections in African infants. II. Clinical and serological observations during seroconversion. *Int J Cancer* 22(3): 244-250.
- Cancer Genome Atlas Research Network. 2014. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature* 513(7517): 202-209. (as cited in Gulley 2015)
- Carpenter LM, Newton R, Casabonne D, Ziegler J, Mbulaiteye S, Mbidde E, Wabinga H, Jaffe H, Beral V. 2008. Antibodies against malaria and Epstein-Barr virus in childhood Burkitt lymphoma: a case-control study in Uganda. *Int J Cancer* 122(6): 1319-1323.

- CDC. 2014a. *About Epstein-Barr virus (EBV)*. Centers for Disease Control and Prevention. <http://www.cdc.gov/epstein-barr/about-ebv.html>. Last updated: 1/6/14.
- CDC. 2014b. *About Infectious Mononucleosis*. Centers for Disease Control and Prevention. <http://www.cdc.gov/epstein-barr/about-mono.html>. Last updated: 1/7/2014.
- CDC. 2015. *List of Vaccines Used in United States*. Centers for Disease Control and Prevention. <http://www.cdc.gov/vaccines/vpd-vac/vaccines-list.htm>. Last updated: 9/3/15.
- Chaigne-Delalande B, Li FY, O'Connor GM, Lukacs MJ, Jiang P, Zheng L, *et al.* 2013. Mg²⁺ regulates cytotoxic functions of NK and CD8 T cells in chronic EBV infection through NKG2D. *Science* 341(6142): 186-191.
- Cohen JI. 2009. Optimal treatment for chronic active Epstein-Barr virus disease. *Pediatr Transplant* 13(4): 393-396.
- Cohen JI. 2015. Epstein-barr virus vaccines. *Clin Transl Immunology* 4(1): e32.
- Condon LM, Cederberg LE, Rabinovitch MD, Liebo RV, Go JC, Delaney AS, Schmelting DO, Thomas W, Balfour HH Jr. 2014. Age-specific prevalence of Epstein-Barr virus infection among Minnesota children: effects of race/ethnicity and family environment. *Clin Infect Dis* 59(4): 501-508.
- De Aquino PF, Carvalho PC, da Gama Fischer JS, de Souza AQ, Viana JS, Chalub SR, de Souza AD, Carvalho MG. 2012. Epstein-Barr virus DNA associated with gastric adenocarcinoma and adjacent non-cancerous mucosa in patients from Manaus, Brazil. *Genet Mol Res* 11(4): 4442-4446.
- Dowd JB, Palermo T, Brite J, McDade TW, Aiello A. 2013. Seroprevalence of Epstein-Barr virus infection in U.S. children ages 6-19, 2003-2010. *PLoS One* 8(5): e64921.
- Epstein MA, Achong BG, Barr YM. 1964. Virus Particles in Cultured Lymphoblasts from Burkitt's Lymphoma. *Lancet* 1(7335): 702-703.
- FDA. 2015a. *Blood Guidances*. U.S. Food and Drug Administration. <http://www.fda.gov/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/Blood/ucm2008053.htm>. Last updated: 5/12/15.
- FDA. 2015b. *Complete List of Vaccines Licensed for Immunization and Distribution in the US*. U.S. Food and Drug Administration. <http://www.fda.gov/BiologicsBloodVaccines/Vaccines/ApprovedProducts/ucm093833.htm>. Last updated: 6/17/15.
- Gulley ML. 2015. Genomic assays for Epstein-Barr virus-positive gastric adenocarcinoma. *Exp Mol Med* 47: e134.
- He YJ, Jia XS, Hasui K, Wang EH, He AG. 2007. [Nasal and pharyngeal non-Hodgkin lymphomas and their relationship with Epstein-Barr virus: a report of 158 cases]. [In Chinese]. *Zhonghua Bing Li Xue Za Zhi* 36(2): 94-97.
- Henle G, Henle W, Clifford P, Diehl V, Kafuko GW, Kirya BG, *et al.* 1969. Antibodies to Epstein-Barr virus in Burkitt's lymphoma and control groups. *J Natl Cancer Inst* 43(5): 1147-1157.

- Henle G, Henle W, Klein G, Gunven P, Clifford P, Morrow RH, Ziegler JL. 1971. Antibodies to early Epstein-Barr virus-induced antigens in Burkitt's lymphoma. *J Natl Cancer Inst* 46(4): 861-871.
- IARC. 1997. Epstein-Barr virus. In *Epstein-Barr Virus and Kaposi's Sarcoma Herpesvirus/Human Herpesvirus 8*. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans, vol. 70. Lyon, France: International Agency for Research on Cancer. pp. 47-373.
- IARC. 2012. Epstein-Barr virus. In *Biological Agents*. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans, vol. 100B. Lyon, France: International Agency for Research on Cancer. pp. 49-92.
- Imai S, Nishikawa J, Takada K. 1998. Cell-to-cell contact as an efficient mode of Epstein-Barr virus infection of diverse human epithelial cells. *J Virol* 72(5): 4371-4378.
- Kim Y, Shin A, Gwack J, Ko KP, Kim CS, Park SK, Hong YC, Kang D, Yoo KY. 2009. Epstein-Barr virus antibody level and gastric cancer risk in Korea: a nested case-control study. *Br J Cancer* 101(3): 526-529.
- Koshiol J, Qiao YL, Mark SD, Dawsey SM, Abnet CC, Kamangar F, Lennette ET, Dong ZW, Taylor PR. 2007. Epstein-Barr virus serology and gastric cancer incidence and survival. *Br J Cancer* 97(11): 1567-1569.
- Lei KI, Chan LY, Chan WY, Johnson PJ, Lo YM. 2000. Quantitative analysis of circulating cell-free Epstein-Barr virus (EBV) DNA levels in patients with EBV-associated lymphoid malignancies. *Br J Haematol* 111(1): 239-246.
- Levine PH, Stemmermann G, Lennette ET, Hildesheim A, Shibata D, Nomura A. 1995. Elevated antibody titers to Epstein-Barr virus prior to the diagnosis of Epstein-Barr-virus-associated gastric adenocarcinoma. *Int J Cancer* 60(5): 642-644.
- Linabery AM, Erhardt EB, Fonstad RK, Ambinder RF, Bunin GR, Ross JA, Spector LG, Grufferman S. 2014. Infectious, autoimmune and allergic diseases and risk of Hodgkin lymphoma in children and adolescents: a Children's Oncology Group study. *Int J Cancer* 135(6): 1454-1469.
- Liu P, Fang X, Feng Z, Guo YM, Peng RJ, Liu T, *et al.* 2011. Direct sequencing and characterization of a clinical isolate of Epstein-Barr virus from nasopharyngeal carcinoma tissue by using next-generation sequencing technology. *J Virol* 85(21): 11291-11299.
- Lo YM, Chan WY, Ng EK, Chan LY, Lai PB, Tam JS, Chung SC. 2001. Circulating Epstein-Barr virus DNA in the serum of patients with gastric carcinoma. *Clin Cancer Res* 7(7): 1856-1859.
- Lombardi L, Newcomb EW, Dalla-Favera R. 1987. Pathogenesis of Burkitt lymphoma: expression of an activated c-myc oncogene causes the tumorigenic conversion of EBV-infected human B lymphoblasts. *Cell* 49(2): 161-170.
- Lu J, Murakami M, Verma SC, Cai QL, Haldar S, Kaul R, Wasik MA, Middeldorp J, Robertson ES. 2011. Epstein-Barr Virus nuclear antigen 1 (EBNA1) confers resistance to apoptosis in EBV-positive B-lymphoma cells through up-regulation of survivin. *Virology* 410(1): 64-75.

- Mohamed G, Vrzalikova K, Cader FZ, Vockerodt M, Nagy E, Flodr P, *et al.* 2014. Epstein-Barr virus, the germinal centre and the development of Hodgkin's lymphoma. *J Gen Virol* 95(Pt 9): 1861-1869.
- Moormann AM, Heller KN, Chelimo K, Embury P, Ploutz-Snyder R, Otieno JA, Oduor M, Munz C, Rochford R. 2009. Children with endemic Burkitt lymphoma are deficient in EBNA1-specific IFN-gamma T cell responses. *Int J Cancer* 124(7): 1721-1726. (as cited in IARC 2012).
- Mosier DE, Gulizia RJ, Baird SM, Spector S, Spector D, Kipps TJ, *et al.* 1989. Studies of HIV infection and the development of Epstein-Barr virus-related B cell lymphomas following transfer of human lymphocytes to mice with severe combined immunodeficiency. *Curr Top Microbiol Immunol* 152: 195-199.
- Mulama DH, Bailey JA, Foley J, Chelimo K, Ouma C, Jura WG, Otieno J, Vulule J, Moormann AM. 2014. Sick cell trait is not associated with endemic Burkitt lymphoma: an ethnicity and malaria endemicity-matched case-control study suggests factors controlling EBV may serve as a predictive biomarker for this pediatric cancer. *Int J Cancer* 134(3): 645-653.
- Mutalima N, Molyneux E, Jaffe H, Kamiza S, Borgstein E, Mkandawire N, *et al.* 2008. Associations between Burkitt lymphoma among children in Malawi and infection with HIV, EBV and malaria: results from a case-control study. *PLoS One* 3(6): e2505.
- Newton R, Carpenter L, Casabonne D, Beral V, Babiker A, Darbyshire J, *et al.* 2006. A prospective study of Kaposi's sarcoma-associated herpesvirus and Epstein-Barr virus in adults with human immunodeficiency virus-1. *Br J Cancer* 94(10): 1504-1509.
- Pathmanathan R, Prasad U, Sadler R, Flynn K, Raab-Traub N. 1995. Clonal proliferations of cells infected with Epstein-Barr virus in preinvasive lesions related to nasopharyngeal carcinoma. *N Engl J Med* 333(11): 693-698.
- Piriou E, Asito AS, Sumba PO, Fiore N, Middeldorp JM, Moormann AM, Ploutz-Snyder R, Rochford R. 2012. Early age at time of primary Epstein-Barr virus infection results in poorly controlled viral infection in infants from Western Kenya: clues to the etiology of endemic Burkitt lymphoma. *J Infect Dis* 205(6): 906-913.
- Ponce RA, Gelzleichter T, Haggerty HG, Heidel S, Holdren MS, Lebec H, Mellon RD, Pallardy M. 2014. Immunomodulation and lymphoma in humans. *J Immunotoxicol* 11(1): 1-12.
- Raab-Traub N, Rajadurai P, Flynn K, Lanier AP. 1991. Epstein-Barr virus infection in carcinoma of the salivary gland. *J Virol* 65(12): 7032-7036.
- Raab-Traub N. 2002. Epstein-Barr virus in the pathogenesis of NPC. *Semin Cancer Biol* 12(6): 431-441.
- Rigaud S, Fondaneche MC, Lambert N, Pasquier B, Mateo V, Soulas P, *et al.* 2006. XIAP deficiency in humans causes an X-linked lymphoproliferative syndrome. *Nature* 444(7115): 110-114.
- Rowe M, Young LS, Crocker J, Stokes H, Henderson S, Rickinson AB. 1991. Epstein-Barr virus (EBV)-associated lymphoproliferative disease in the SCID mouse model: implications for the pathogenesis of EBV-positive lymphomas in man. *J Exp Med* 173(1): 147-158.

- Shinkura R, Yamamoto N, Koriyama C, Shinmura Y, Eizuru Y, Tokunaga M. 2000. Epstein-Barr virus-specific antibodies in Epstein-Barr virus-positive and -negative gastric carcinoma cases in Japan. *J Med Virol* 60(4): 411-416.
- Taylor AL, Marcus R, Bradley JA. 2005. Post-transplant lymphoproliferative disorders (PTLD) after solid organ transplantation. *Crit Rev Oncol Hematol* 56(1): 155-167.
- Thompson MP, Kurzrock R. 2004. Epstein-Barr virus and cancer. *Clin Cancer Res* 10(3): 803-821.
- Thorley-Lawson DA, Gross A. 2004. Persistence of the Epstein-Barr virus and the origins of associated lymphomas. *N Engl J Med* 350(13): 1328-1337.
- Tsai CC, Chen CL, Hsu HC. 1996. Expression of Epstein-Barr virus in carcinomas of major salivary glands: a strong association with lymphoepithelioma-like carcinoma. *Hum Pathol* 27(3): 258-262.
- Tsang CM, Deng W, Yip YL, Zeng MS, Lo KW, Tsao SW. 2014. Epstein-Barr virus infection and persistence in nasopharyngeal epithelial cells. *Chin J Cancer* 33(11): 549-555.
- Tsang WYW, Chan JKC. 2005. Lymphoepithelial carcinoma. In *World Health Organization Classification of Tumours – Pathology and Genetics of Head and Neck Tumours*. Barnes EL, Eveson JW, Reichart P, Sidransky D, eds. Lyon, France: International Agency for Research on Cancer. pp. 85-97. (as cited in IARC 2012)
- Tsao SW, Tsang CM, To KF, Lo KW. 2015. The role of Epstein-Barr virus in epithelial malignancies. *J Pathol* 235(2): 323-333.
- Vegso G, Hajdu M, Sebestyen A. 2011. Lymphoproliferative disorders after solid organ transplantation-classification, incidence, risk factors, early detection and treatment options. *Pathol Oncol Res* 17(3): 443-454.
- Wang CP, Chang YL, Ko JY, Lou PJ, Yeh CF, Sheen TS. 2004. Lymphoepithelial carcinoma versus large cell undifferentiated carcinoma of the major salivary glands. *Cancer* 101(9): 2020-2027.

This Page Intentionally Left Blank