

1 **NTP Technical Report on the**
2 **Toxicology and Carcinogenesis Studies of**
3 **Sodium Tungstate Dihydrate**
4 **(CASRN 10213-10-2) in Sprague Dawley**
5 **(Hsd:Sprague Dawley[®] SD[®]) Rats**
6 **and B6C3F1/N Mice**
7 **(Drinking Water Studies)**

8 Technical Report 599

9 February 2021

10 National Toxicology Program
11 Public Health Service
12 U.S. Department of Health and Human Services
13 ISSN: 2378-8925

14 Research Triangle Park, North Carolina, USA

NOTICE

This DRAFT Technical Report is distributed solely for the purpose of pre-dissemination peer review under the applicable information quality guidelines. It has not been formally disseminated by NTP. It does not represent and should not be construed to represent NTP determination or policy.

1

Foreword

2 The National Toxicology Program (NTP), established in 1978, is an interagency program within
3 the Public Health Service of the U.S. Department of Health and Human Services. Its activities
4 are executed through a partnership of the National Institute for Occupational Safety and Health
5 (part of the Centers for Disease Control and Prevention), the Food and Drug Administration
6 (primarily at the National Center for Toxicological Research), and the National Institute of
7 Environmental Health Sciences (part of the National Institutes of Health), where the program is
8 administratively located. NTP offers a unique venue for the testing, research, and analysis of
9 agents of concern to identify toxic and biological effects, provide information that strengthens
10 the science base, and inform decisions by health regulatory and research agencies to safeguard
11 public health. NTP also works to develop and apply new and improved methods and approaches
12 that advance toxicology and better assess health effects from environmental exposures.

13 The Technical Report series began in 1976 with carcinogenesis studies conducted by the
14 National Cancer Institute. In 1981, this bioassay program was transferred to NTP. The studies
15 described in the NTP Technical Report series are designed and conducted to characterize and
16 evaluate the toxicological potential, including carcinogenic activity, of selected substances in
17 laboratory animals (usually two species, rats and mice). Substances (e.g., chemicals, physical
18 agents, and mixtures) selected for NTP toxicity and carcinogenicity studies are chosen primarily
19 on the basis of human exposure, level of commercial production, and chemical structure. The
20 interpretive conclusions presented in NTP Technical Reports are derived solely from the results
21 of these NTP studies, and extrapolation of the results to other species, including characterization
22 of hazards and risks to humans, requires analyses beyond the intent of these reports. Selection for
23 study per se is not an indicator of a substance's carcinogenic potential.

24 NTP conducts its studies in compliance with its laboratory health and safety guidelines and the
25 Food and Drug Administration [Good Laboratory Practice Regulations](#) and meets or exceeds all
26 applicable federal, state, and local health and safety regulations. Animal care and use are in
27 accordance with the [Public Health Service Policy on Humane Care and Use of Laboratory](#)
28 [Animals](#). Studies are subjected to retrospective quality assurance audits before they are presented
29 for public review. Draft reports undergo external peer review before they are finalized and
30 published.

31 The NTP Technical Reports are available free of charge on the [NTP website](#) and cataloged in
32 [PubMed](#), a free resource developed and maintained by the National Library of Medicine (part of
33 the National Institutes of Health). Data for these studies are included in NTP's [Chemical Effects](#)
34 [in Biological Systems](#) database.

35 For questions about the reports and studies, please email NTP or call 984-287-3211.

1		
2	Foreword.....	ii
3	Tables.....	v
4	Figures.....	vii
5	About This Report.....	ix
6	Explanation of Levels of Evidence of Carcinogenic Activity	xiii
7	Peer Review	xv
8	Publication Details.....	xvi
9	Acknowledgments.....	xvi
10	Abstract.....	xvii
11	Perinatal and Three-month Study in Rats	xvii
12	Perinatal and Two-year Study in Rats.....	xviii
13	Three-month Study in Mice	xviii
14	Two-year Study in Mice.....	xix
15	Genetic Toxicology	xix
16	Conclusions	xx
17	Introduction.....	1
18	Chemical and Physical Properties.....	1
19	Production, Use, and Human Exposure	1
20	Regulatory Status	2
21	Absorption, Distribution, Metabolism, and Excretion.....	2
22	Experimental Animals	2
23	Humans	3
24	Toxicity	3
25	Experimental Animals	3
26	Humans	4
27	Reproductive and Developmental Toxicity	4
28	Experimental Animals	4
29	Humans	4
30	Carcinogenicity	4
31	Genetic Toxicity.....	4
32	Study Rationale.....	5
33	Materials and Methods.....	6
34	Procurement and Characterization of Sodium Tungstate Dihydrate.....	6
35	Preparation and Analysis of Dose Formulations.....	6
36	Animal Source.....	7
37	Animal Welfare.....	7
38	Three-month Studies	7
39	Initial Exposure Concentration Selection Rationale.....	7
40	Study Design for Rats	8
41	Study Design for Mice	9

1	Clinical Examinations and Pathology.....	9
2	Two-year Studies	11
3	Study Design for Rats	11
4	Study Design for Mice	12
5	Clinical Examinations, Tungsten Concentrations, and Pathology.....	13
6	Statistical Methods	20
7	Survival Analyses	20
8	Calculation of Incidence	20
9	Analysis of Neoplasm and Nonneoplastic Lesion Incidence.....	20
10	Analysis of Continuous Variables	22
11	Analysis of Gestational and Fertility Indices.....	22
12	Body Weight Adjustments.....	23
13	Historical Control Data.....	23
14	Quality Assurance Methods.....	23
15	Genetic Toxicology.....	23
16	Bacterial Mutagenicity.....	24
17	Peripheral Blood Micronucleus Test	24
18	Comet Assay.....	24
19	Results.....	25
20	Data Availability	25
21	Rats	25
22	Three-month Study (Perinatal Phase).....	25
23	Three-month Study (Postweaning Phase).....	31
24	Exposure Concentration Selection Rationale for the Perinatal and Two-year Study	
25	in Rats	40
26	Two-year Study (Perinatal Phase)	40
27	Two-year Study (Interim Evaluations – 3, 6, 12, and 18 Months).....	46
28	Two-year Study (Postweaning Phase).....	53
29	Mice	70
30	Three-month Study	70
31	Exposure Concentration Selection Rationale for Two-year Studies in Mice.....	77
32	Two-year Study (Interim Evaluations – 3, 6, 12, and 18 Months).....	78
33	Two-year Study.....	85
34	Genetic Toxicology.....	100
35	Discussion.....	102
36	Conclusions.....	106
37	References.....	107
38	Appendix A. Chemical Characterization and Dose Formulation Studies.....	A-1
39	Appendix B. Ingredients, Nutrient Composition, and Contaminant Levels in NIH-07 Rat	
40	Ration and NTP-2000 Rat and Mouse Ration	B-1
41	Appendix C. Sentinel Animal Program	C-1
42	Appendix D. Genetic Toxicology	D-1
43	Appendix E. Tungsten Concentration Determination.....	E-1

1	Appendix F. Peer-review Report	F-1
2	Appendix G. Supplemental Data	G-1

3 Tables

4	Summary of the Perinatal and Two-year Carcinogenesis and Genetic Toxicology Studies	
5	of Sodium Tungstate Dihydrate.....	xxi
6	Table 1. Experimental Design and Materials and Methods in the Three-month and Two-	
7	year Drinking Water Studies of Sodium Tungstate Dihydrate.....	15
8	Table 2. Summary of the Disposition of F ₀ Female Rats during Perinatal Exposure in the	
9	Perinatal and Three-month Drinking Water Study of Sodium Tungstate	
10	Dihydrate	25
11	Table 3. Summary of Mean Body Weights and Body Weight Gains of F ₀ Female Rats	
12	during Gestation and Lactation in the Perinatal and Three-month Drinking	
13	Water Study of Sodium Tungstate Dihydrate	26
14	Table 4. Summary of Water and Sodium Tungstate Dihydrate Consumption of F ₀ Female	
15	Rats during Gestation and Lactation in the Perinatal and Three-month Drinking	
16	Water Study	27
17	Table 5. Summary of Mean Litter Size and Survival Ratio of F ₁ Male and Female Rats	
18	during Lactation in the Perinatal and Three-month Drinking Water Study of	
19	Sodium Tungstate Dihydrate	29
20	Table 6. Summary of Prewaning F ₁ Male and Female Rat Pup Mean Body Weights	
21	Following Perinatal Exposure to Sodium Tungstate Dihydrate	30
22	Table 7. Summary of Survival and Mean Body Weights of Male Rats in the Perinatal and	
23	Three-month Drinking Water Study of Sodium Tungstate Dihydrate	32
24	Table 8. Summary of Survival and Mean Body Weights of Female Rats in the Perinatal	
25	and Three-month Drinking Water Study of Sodium Tungstate Dihydrate	33
26	Table 9. Summary of Water and Sodium Tungstate Dihydrate Consumption of Male and	
27	Female Rats in the Perinatal and Three-month Drinking Water Study	35
28	Table 10. Summary of Blood and Urine Tungsten Concentration Data for Male and	
29	Female Rats in the Perinatal and Three-month Drinking Water Study of	
30	Sodium Tungstate Dihydrate	36
31	Table 11. Summary of Select Clinical Pathology Data for Male and Female Rats in the	
32	Perinatal and Three-month Drinking Water Study of Sodium Tungstate	
33	Dihydrate	37
34	Table 12. Summary of Select Urinalysis Data for Male and Female Rats in the Perinatal	
35	and Three-month Drinking Water Study of Sodium Tungstate Dihydrate.....	38
36	Table 13. Summary of Renal Findings for Male and Female Rats in the Perinatal and	
37	Three-month Drinking Water Study of Sodium Tungstate Dihydrate	39
38	Table 14. Summary of the Disposition of F ₀ Female Rats during Perinatal Exposure in the	
39	Perinatal and Two-year Drinking Water Study of Sodium Tungstate Dihydrate.....	40
40	Table 15. Summary of Mean Body Weights and Body Weight Gains of F ₀ Female Rats	
41	during Gestation and Lactation in the Perinatal and Two-year Drinking Water	
42	Study of Sodium Tungstate Dihydrate.....	41

1	Table 16. Summary of Water and Sodium Tungstate Dihydrate Consumption by	
2	F ₀ Female Rats during Gestation and Lactation in the Perinatal and Two-year	
3	Drinking Water Study	42
4	Table 17. Summary of Mean Litter Size and Survival Ratio of F ₁ Male and Female Rats	
5	during Lactation in the Perinatal and Two-year Drinking Water Study of	
6	Sodium Tungstate Dihydrate	43
7	Table 18. Summary of Preweaning F ₁ Male and Female Rat Pup Mean Body Weights	
8	Following Perinatal Exposure to Sodium Tungstate Dihydrate	44
9	Table 19. Summary of Internal Dose Data for F ₀ Female Rats and Pups in the Perinatal	
10	and Two-year Drinking Water Study of Sodium Tungstate Dihydrate	45
11	Table 20. Summary of Select Organ Weights and Organ-Weight-to-Body-Weight Ratios	
12	for Male Rats Exposed to Sodium Tungstate Dihydrate in Drinking Water for	
13	3, 6, 12, and 18 Months	47
14	Table 21. Summary of Select Organ Weights and Organ-Weight-to-Body-Weight Ratios	
15	for Female Rats Exposed to Sodium Tungstate Dihydrate in Drinking Water for	
16	3, 6, 12, and 18 Months	48
17	Table 22. Summary of Plasma, Kidney, and Urine Tungsten Concentration Data for Male	
18	Rats Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12,	
19	and 18 Months	50
20	Table 23. Summary of Plasma, Kidney, and Urine Tungsten Concentration Data for	
21	Female Rats Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3,	
22	6, 12, and 18 Months	51
23	Table 24. Summary of Survival of Male and Female Rats in the Perinatal and Two-year	
24	Drinking Water Study of Sodium Tungstate Dihydrate	53
25	Table 25. Summary of Survival and Mean Body Weights of Male Rats in the Perinatal	
26	and Two-year Drinking Water Study of Sodium Tungstate Dihydrate	56
27	Table 26. Summary of Survival and Mean Body Weights of Female Rats in the Perinatal	
28	and Two-year Drinking Water Study of Sodium Tungstate Dihydrate	57
29	Table 27. Summary of Water and Sodium Tungstate Dihydrate Consumption of Male	
30	Rats in the Perinatal and Two-year Drinking Water Study	59
31	Table 28. Summary of Water and Sodium Tungstate Dihydrate Consumption of Female	
32	Rats in the Perinatal and Two-year Drinking Water Study of Sodium Tungstate	
33	Dihydrate	59
34	Table 29. Incidences of Neoplastic and Nonneoplastic Lesions of the Thyroid Gland in	
35	Female Rats in the Perinatal and Two-year Drinking Water Study of Sodium	
36	Tungstate Dihydrate.....	60
37	Table 30. Incidences of Nonneoplastic Lesions of the Kidney in Male and Female Rats in	
38	the Perinatal and Two-year Drinking Water Study of Sodium Tungstate	
39	Dihydrate	63
40	Table 31. Incidences of Neoplastic and Nonneoplastic Lesions of the Uterus in Female	
41	Rats in the Perinatal and Two-year Drinking Water Study of Sodium Tungstate	
42	Dihydrate	67
43	Table 32. Summary of Survival and Mean Body Weights of Male Mice in the Three-	
44	month Drinking Water Study of Sodium Tungstate Dihydrate	71
45	Table 33. Summary of Survival and Body Weights of Female Mice in the Three-month	
46	Drinking Water Study of Sodium Tungstate Dihydrate	72

1	Table 34. Summary of Water and Sodium Tungstate Dihydrate Consumption of Male and	
2	Female Mice in the Three-month Drinking Water Study	74
3	Table 35. Summary of Blood Tungsten Concentration Data for Male and Female Mice in	
4	the Three-month Drinking Water Study of Sodium Tungstate Dihydrate.....	75
5	Table 36. Summary of Select Hematology Data for Male Mice in the Three-month	
6	Drinking Water Study of Sodium Tungstate Dihydrate	76
7	Table 37. Summary of Renal Findings for Male and Female Mice in the Three-month	
8	Drinking Water Study of Sodium Tungstate Dihydrate	76
9	Table 38. Summary of Select Organ Weights and Organ-Weight-to-Body-Weight Ratios	
10	for Male Mice Exposed to Sodium Tungstate Dihydrate in Drinking Water for	
11	3, 6, 12, and 18 Months	79
12	Table 39. Summary of Select Organ Weights and Organ-Weight-to-Body-Weight Ratios	
13	for Female Mice Exposed to Sodium Tungstate Dihydrate in Drinking Water	
14	for 3, 6, 12, and 18 Months.....	80
15	Table 40. Summary of Plasma, Kidney, and Urine Tungsten Concentration Data for Male	
16	Mice Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12,	
17	and 18 Months	82
18	Table 41. Summary of Plasma, Kidney, and Urine Tungsten Concentration Data for	
19	Female Mice Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3,	
20	6, 12, and 18 Months	83
21	Table 42. Summary of Survival of Male and Female Mice in the Two-year Drinking	
22	Water Study of Sodium Tungstate Dihydrate.....	85
23	Table 43. Summary of Survival and Mean Body Weights of Male Mice in the Two-year	
24	Drinking Water Study of Sodium Tungstate Dihydrate	88
25	Table 44. Summary of Survival and Mean Body Weights of Female Mice in the Two-	
26	year Drinking Water Study of Sodium Tungstate Dihydrate	89
27	Table 45. Summary of Water and Sodium Tungstate Dihydrate Consumption of Male	
28	Mice in the Two-year Drinking Water Study	91
29	Table 46. Summary of Water and Sodium Tungstate Dihydrate Consumption of Female	
30	Mice in the Two-year Drinking Water Study	91
31	Table 47. Incidences of Neoplastic and Nonneoplastic Lesions of the Kidney in Male and	
32	Female Mice in the Two-year Drinking Water Study of Sodium Tungstate	
33	Dihydrate	93
34	Table 48. Incidences of Select Nonneoplastic Lesions in Male and Female Mice in the	
35	Two-year Drinking Water Study of Sodium Tungstate Dihydrate.....	98

36

Figures

37	Figure 1. Sodium Tungstate Dihydrate (CASRN 10213-10-2; Chemical Formula:	
38	$\text{Na}_2\text{WO}_4 \cdot 2\text{H}_2\text{O}$; Molecular Weight: 329.86).....	1
39	Figure 2. Growth Curves for Rats Exposed to Sodium Tungstate Dihydrate in Drinking	
40	Water for Three Months	34
41	Figure 3. Tungsten Concentrations in Plasma, Kidney, and Urine in Rats Exposed to	
42	Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months.....	52
43	Figure 4. Kaplan-Meier Survival Curves for Rats Exposed to Sodium Tungstate	
44	Dihydrate in Drinking Water for Two Years	54

1	Figure 5. Growth Curves for Rats Exposed to Sodium Tungstate Dihydrate in Drinking	
2	Water for Two Years	58
3	Figure 6. C-cell Adenoma in the Thyroid Gland of a Female Sprague Dawley Rat	
4	Exposed to 1,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E).....	61
5	Figure 7. C-cell Carcinoma in the Thyroid Gland of a Female Sprague Dawley Rat	
6	Exposed to 1,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E).....	62
7	Figure 8. Suppurative Inflammation in the Renal Tubules of the Kidney from a Male	
8	Sprague Dawley Rat Exposed to 1,000 mg/L Sodium Tungstate Dihydrate for	
9	Two Years (H&E)	64
10	Figure 9. Renal Tubule Regeneration in the Kidney of a Female Sprague Dawley Rat	
11	Exposed to 1,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E).....	65
12	Figure 10. Chronic Progressive Nephropathy in the Kidney of a Female Sprague Dawley	
13	Rat Exposed to 1,000 mg/L Sodium Tungstate Dihydrate for Two Years	
14	(H&E)	66
15	Figure 11. Atypical Hyperplasia in the Uterus of a Female Sprague Dawley Rat Exposed	
16	to 500 mg/L Sodium Tungstate Dihydrate for Two Years (H&E).....	68
17	Figure 12. Growth Curves for Mice Exposed to Sodium Tungstate Dihydrate in Drinking	
18	Water for Three Months	73
19	Figure 13. Tungsten Concentrations in Plasma, Kidney, and Urine in Mice Exposed to	
20	Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months	84
21	Figure 14. Kaplan-Meier Survival Curves for Mice Exposed to Sodium Tungstate	
22	Dihydrate in Drinking Water for Two Years.....	86
23	Figure 15. Growth Curves for Mice Exposed to Sodium Tungstate Dihydrate in Drinking	
24	Water for Two Years	90
25	Figure 16. Renal Tubule Adenoma in a Male B6C3F1/N Mouse Exposed to 1,000 mg/L	
26	Sodium Tungstate Dihydrate for Two Years (H&E).....	94
27	Figure 17. Renal Tubule Carcinoma in a Male B6C3F1/N Mouse Exposed to 2,000 mg/L	
28	Sodium Tungstate Dihydrate for Two Years (H&E).....	94
29	Figure 18. Normal Kidney from a Vehicle Control Female B6C3F1/N Mouse in the Two-	
30	year Study of Sodium Tungstate Dihydrate (H&E)	95
31	Figure 19. Moderate Renal Tubule Regeneration in a Female B6C3F1/N Mouse Exposed	
32	to 2,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E).....	96
33	Figure 20. Pigment in the Cecum of B6C3F1/N Mice Exposed to 0 or 2,000 mg/L	
34	Sodium Tungstate Dihydrate for Two Years (H&E).....	99
35	Figure 21. Germinal Epithelium Degeneration in the Testis from B6C3F1/N Mice	
36	Exposed to 2,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E)	100
37		

1

About This Report

2 National Toxicology Program¹3 ¹Division of the National Toxicology Program, National Institute of Environmental Health
4 Sciences, Research Triangle Park, North Carolina, USA

5

Collaborators

6 M. Behl, A.E. Brix, C.M. Briscoe, W.M. Black, C.R. Blystone, P. Brown, M.A. Buccellato,
7 B.L. Burbach, K.A. Carrico, K.Y. Cimon, D.A. Contos, M.C. Cora, T.A. Cristy, D.M. Fallacara,
8 L.M. Fomby, J.M. Fostel, H. Gong, S.W. Graves, M.R. Hejtmancik, M.J. Hooth,
9 A.P. King-Herbert, D.E. Malarkey, C. Martini, B.S. McIntyre, E. Mutlu, C. Myers, E.M. Quist,
10 J.S. Richey, G.K. Roberts, M.J. Ryan, N. Sayers, M. Shaw, K.A. Shipkowski, K.R. Shockley,
11 A.W. Singer, A.J. Skowronek, C.S. Sloan, S.L. Smith-Roe, N.L. South, B.R. Sparrow,
12 M.D. Stout, G.S. Travlos, R.W. Tyl, S. Waidyanatha, N.J. Walker, R. Whittlesey, K.L. Witt13 **Division of the National Toxicology Program, National Institute of Environmental Health**
14 **Sciences, Research Triangle Park, North Carolina, USA**15 *Designed studies, evaluated and interpreted results, and reported findings*

16 M. Behl, Ph.D., Study Scientist

17 C.R. Blystone, Ph.D.

18 M.C. Cora, D.V.M.

19 M.J. Hooth, Ph.D.

20 A.P. King-Herbert, D.V.M.

21 D.E. Malarkey, D.V.M., Ph.D.

22 B.S. McIntyre, Ph.D.

23 E. Mutlu, Ph.D.

24 G.K. Roberts, Ph.D.

25 K.A. Shipkowski, Ph.D.

26 K.R. Shockley, Ph.D.

27 S.L. Smith-Roe, Ph.D.

28 M.D. Stout, Ph.D.

29 G.S. Travlos, D.V.M.

30 S. Waidyanatha, Ph.D.

31 N.J. Walker, Ph.D.

32 K.L. Witt, M.S.

33 *Provided oversight for data management*

34 J.M. Fostel, Ph.D.

35 **Experimental Pathology Laboratories, Inc., Research Triangle Park, North Carolina, USA**36 *Evaluated and interpreted results and reported findings*

37 A.E. Brix, D.V.M., Ph.D., Study Pathologist

1 *Coordinated NTP Pathology Working Groups on perinatal and 2-year rats (April 3, 2018) and*
2 *on 2-year mice (May 3, 2018)*

3 K.Y. Cimon, D.V.M. (Rats)

4 E.M. Quist, D.V.M., Ph.D. (Mice)

5 **Battelle, Columbus, Ohio, USA**

6 *Conducted studies and evaluated pathology findings*

7 M.R. Hejtmancik, Ph.D., Principal Investigator

8 B.R. Sparrow, Ph.D., Principal Investigator

9 C.M. Briscoe, D.V.M.

10 M.A. Buccellato, D.V.M., Ph.D.

11 D.M. Fallacara, Ph.D.

12 L.M. Fomby, D.V.M., Ph.D.

13 M.J. Ryan, D.V.M., Ph.D.

14 A.W. Singer, D.V.M.

15 A.J. Skowronek, D.V.M., Ph.D.

16 *Conducted prestart chemistry activities, dose formulations, and biological sample chemistry*
17 *analyses*

18 S.W. Graves, B.S., Principal Investigator

19 B.L. Burbach, Ph.D., Principal Investigator

20 W.M. Black, B.S.

21 K.A. Carrico, B.A.

22 D.A. Contos, M.S.

23 T.A. Cristy, B.A.

24 J.S. Richey, B.S.

25 N.L. South, B.S.

26 **RTI International, Research Triangle Park, North Carolina, USA**

27 *Provided sperm count and vaginal cytology evaluation (SCVCE) analysis*

28 C.S. Sloan, M.S.

29 R.W. Tyl, Ph.D.

30 **ASRC Federal, Research Triangle Park, North Carolina, USA**

31 *Prepared data for report*

32 P. Brown, B.S.

33 H. Gong, M.S.

34 C. Martini, B.S.

35 C. Myers, M.S.

36 N. Sayers, B.S.

37 M. Shaw, B.S.

38 R. Whittlesey, M.S.

1 **Contributors**

2 **Division of the National Toxicology Program, National Institute of Environmental Health**
3 **Sciences, Research Triangle Park, North Carolina, USA**

4 *Provided oversight of external peer review*

5 E.A. Maull, Ph.D.

6 S.L. Scruggs, Ph.D.

7 M.S. Wolfe, Ph.D.

8 **NTP Pathology Working Group, National Institute of Environmental Health Sciences,**
9 **Research Triangle Park, North Carolina, USA**

10 *Participated in NTP Pathology Working Group on perinatal and 2-year rats (April 3, 2018)*

11 A.E. Brix, D.V.M., Ph.D., Experimental Pathology Laboratories, Inc.

12 D. Dixon, D.V.M., Ph.D., National Toxicology Program

13 S.A. Elmore, D.V.M., M.S., National Toxicology Program

14 G.P. Flake, M.D., National Toxicology Program

15 R.C. Kovi, M.V.Sc., Ph.D., Experimental Pathology Laboratories, Inc.

16 D.E. Malarkey, D.V.M., Ph.D., National Toxicology Program

17 K.S. Regan, D.V.M., Regan Path/Tox Services, Inc.

18 J.C. Seely, D.V.M., Experimental Pathology Laboratories, Inc.

19 A. Skowronek, D.V.M., Ph.D., Battelle (observer)

20 C.J. Willson, D.V.M., Ph.D., Integrated Laboratory Systems, LLC

21 *Participated in NTP Pathology Working Group on 2-year mice (May 3, 2018)*

22 A.E. Brix, D.V.M., Ph.D., Experimental Pathology Laboratories, Inc.

23 D. Dixon, D.V.M., Ph.D., National Toxicology Program

24 S.A. Elmore, D.V.M., M.S., National Toxicology Program

25 G.P. Flake, M.D., National Toxicology Program

26 K. Janardhan, Ph.D., Integrated Laboratory Systems, LLC

27 R.C. Kovi, M.V.Sc., Ph.D., Experimental Pathology Laboratories, Inc.

28 D.E. Malarkey, D.V.M., Ph.D., National Toxicology Program

29 K.S. Regan, D.V.M., Regan Path/Tox Services, Inc.

30 J.C. Seely, D.V.M., Experimental Pathology Laboratories, Inc.

31 A.W. Singer, D.V.M., Battelle (observer)

32 **Charles River Laboratories, Inc., Durham, North Carolina, USA**

33 *Performed pathology review on kidney step sections of 2-year mice (July 25, 2018)*

34 Q.D. Plumlee, D.V.M., Charles River Laboratories, Inc.

35 **Integrated Laboratory Systems, LLC, Research Triangle Park, North Carolina, USA**

36 *Conducted comet, micronucleus, and bacterial mutagenicity assays*

37 L. Recio, Ph.D., Principal Investigator

38 C.A. Hobbs, Ph.D.

39 C.D. Swartz, D.V.M., Ph.D.

40 C. Baldetti, A.A.

41 A.S. Green, B.S.

42 C.A. Lentz, B.S.

1 M.H. Prajapati, M.S.
2 K.G. Shepard, B.S.
3 J.W. Winters, B.S.

4 **CSS Corporation, Research Triangle Park, North Carolina, USA**

5 *Prepared quality assessment audits*
6 S. Brecher, Ph.D., Principal Investigator
7 S. Iyer, B.S.
8 V.S. Tharakan, D.V.M.

9 **Social & Scientific Systems, a DLH Company, Research Triangle Park, North Carolina,**
10 **USA**

11 *Provided statistical analyses*
12 S.J. McBride, Ph.D., Principal Investigator
13 L.J. Betz, M.S.
14 S.F. Harris, M.S.
15 J.D. Krause, Ph.D.

16 **ICF, Fairfax, Virginia, USA**

17 *Provided contract oversight*
18 D. Burch, M.E.M., Principal Investigator
19 J. Cleland, M.E.M.
20 J.A. Wignall, M.S.P.H.

21 *Prepared and edited report*

22 S.R. Gunnels, M.A.
23 T. Hamilton, M.S.
24 J. Luh, Ph.D.
25 M.E. McVey, Ph.D.
26 A. Murphy, B.S.
27 R. Shin, M.H.S.
28 K.A. Shipkowski, Ph.D.
29 S.J. Snow, Ph.D.

1 **Explanation of Levels of Evidence of Carcinogenic Activity**

2 The National Toxicology Program (NTP) describes the results of individual experiments on a
3 chemical agent and notes the strength of the evidence for conclusions regarding each study.
4 Negative results, in which the study animals do not have a greater incidence of neoplasia than
5 control animals, do not necessarily mean that a chemical is not a carcinogen, in as much as the
6 experiments are conducted under a limited set of conditions. Positive results demonstrate that a
7 chemical is carcinogenic for laboratory animals under the conditions of the study and indicate
8 that exposure to the chemical has the potential for hazard to humans. Other organizations, such
9 as the International Agency for Research on Cancer, assign a strength of evidence for
10 conclusions based on an examination of all available evidence, including animal studies such as
11 those conducted by NTP, epidemiologic studies, and estimates of exposure. Thus, the actual
12 determination of risk to humans from chemicals found to be carcinogenic in laboratory animals
13 requires a wider analysis that extends beyond the purview of these studies.

14 Five categories of evidence of carcinogenic activity are used in the Technical Report series to
15 summarize the strength of evidence observed in each experiment: two categories for positive
16 results (**clear evidence and some evidence**); one category for uncertain findings (**equivocal**
17 **evidence**); one category for no observable effects (**no evidence**); and one category for
18 experiments that cannot be evaluated because of major flaws (**inadequate study**). These
19 categories of interpretative conclusions were first adopted in June 1983 and then revised on
20 March 1986 for use in the Technical Report series to incorporate more specifically the concept of
21 actual weight of evidence of carcinogenic activity. For each separate experiment (male rats,
22 female rats, male mice, female mice), one of the following five categories is selected to describe
23 the findings. These categories refer to the strength of the experimental evidence and not to
24 potency or mechanism.

- 25 • **Clear evidence** of carcinogenic activity is demonstrated by studies that are
26 interpreted as showing a dose-related (i) increase of malignant neoplasms, (ii)
27 increase of a combination of malignant and benign neoplasms, or (iii) marked
28 increase of benign neoplasms if there is an indication from this or other studies of the
29 ability of such tumors to progress to malignancy.
- 30 • **Some evidence** of carcinogenic activity is demonstrated by studies that are
31 interpreted as showing a chemical-related increased incidence of neoplasms
32 (malignant, benign, or combined) in which the strength of the response is less than
33 that required for clear evidence.
- 34 • **Equivocal evidence** of carcinogenic activity is demonstrated by studies that are
35 interpreted as showing a marginal increase of neoplasms that may be chemical
36 related.
- 37 • **No evidence** of carcinogenic activity is demonstrated by studies that are interpreted
38 as showing no chemical-related increases in malignant or benign neoplasms.
- 39 • **Inadequate study** of carcinogenic activity is demonstrated by studies that, because of
40 major qualitative or quantitative limitations, cannot be interpreted as valid for
41 showing either the presence or absence of carcinogenic activity.

42 For studies showing multiple chemical-related neoplastic effects that if considered individually
43 would be assigned to different levels of evidence categories, the following convention has been

1 adopted to convey completely the study results. In a study with clear evidence of carcinogenic
2 activity at some tissue sites, other responses that alone might be deemed some evidence are
3 indicated as “were also related” to chemical exposure. In studies with clear or some evidence of
4 carcinogenic activity, other responses that alone might be termed equivocal evidence are
5 indicated as “may have been” related to chemical exposure.

6 When a conclusion statement for a particular experiment is selected, consideration must be given
7 to key factors that would extend the actual boundary of an individual category of evidence. Such
8 consideration should allow for incorporation of scientific experience and current understanding
9 of long-term carcinogenesis studies in laboratory animals, especially for those evaluations that
10 may be on the borderline between two adjacent levels. These considerations should include:

- 11 • adequacy of the experimental design and conduct;
- 12 • occurrence of common versus uncommon neoplasia;
- 13 • progression (or lack thereof) from benign to malignant neoplasia as well as from
14 preneoplastic to neoplastic lesions;
- 15 • some benign neoplasms have the capacity to regress but others (of the same
16 morphologic type) progress. At present, it is impossible to identify the difference.
17 Therefore, where progression is known to be a possibility, the most prudent course is
18 to assume that benign neoplasms of those types have the potential to become
19 malignant;
- 20 • combining benign and malignant tumor incidence known or thought to represent
21 stages of progression in the same organ or tissue;
- 22 • latency in tumor induction;
- 23 • multiplicity in site-specific neoplasia;
- 24 • metastases;
- 25 • supporting information from proliferative lesions (hyperplasia) in the same site of
26 neoplasia or other experiments (same lesion in another sex or species);
- 27 • presence or absence of dose relationships;
- 28 • statistical significance of the observed tumor increase;
- 29 • concurrent control tumor incidence as well as the historical control rate and
30 variability for a specific neoplasm;
- 31 • survival-adjusted analyses and false positive or false negative concerns;
- 32 • structure-activity correlations; and
- 33 • in some cases, genetic toxicology.

1

Peer Review

2 The National Toxicology Program (NTP) convened a virtual external ad hoc panel to peer review
3 the draft *NTP Technical Report on the Toxicology and Carcinogenesis Studies of Sodium*
4 *Tungstate Dihydrate (CASRN 10213-10-2) in Sprague Dawley (Hsd:Sprague Dawley® SD®)*
5 *Rats and B6C3F1/N Mice (Drinking Water Studies)* on April 2, 2021. NTP announced the peer-
6 review meeting in the Federal Register (X FR. XXXX. DATE). The public could view the
7 proceedings online, and opportunities were provided for submission of written and oral public
8 comments. The selection of panel members and conduct of the peer review were in accordance
9 with federal policies and regulations. The panel was charged to:

- 10 (1) Review and evaluate the scientific and technical elements of each study and its
11 presentation.
- 12 (2) Determine whether each study's experimental design, conduct, and findings support
13 the NTP's conclusions regarding the conditions of each study.

14 NTP carefully considered the panel's recommendations in finalizing the report. The peer-review
15 report is provided in Appendix F. Other meeting materials are available on the NTP website
16 (<https://ntp.niehs.nih.gov/go/meeting>).

17

Peer Reviewers

[to come]

1

Publication Details

2 Publisher: National Toxicology Program

3 Publishing Location: Research Triangle Park, NC

4 ISSN: 2378-8925

5 DOI:

6 Report Series: NTP Technical Report Series

7 Report Series Number: 599

8 *Official citation:* National Toxicology Program (NTP). 2021. NTP technical report on the
9 toxicology and carcinogenesis studies of sodium tungstate dihydrate (CASRN 10213-10-2) in
10 Sprague Dawley (Hsd:Sprague Dawley[®] SD[®]) rats and B6C3F1/N mice (drinking water studies).
11 Research Triangle Park, NC: National Toxicology Program. Technical Report 599.

Acknowledgments

13 This work was supported by the Intramural Research Program (ES103316, ES103318, and
14 ES103319) at the National Institute of Environmental Health Sciences, National Institutes of
15 Health and performed for the National Toxicology Program, Public Health Service, U.S.
16 Department of Health and Human Services under contracts HHSN273201800006C,
17 HHSN273201600011C, HHSN273201600020C, GS00Q14OADU417 (Order No.
18 HHSN273201600015U), HHSN273201500006C, HHSN273201500012C,
19 HHSN273201500014C, HHSN273201300009C, HHSN316201200054W,
20 HHSN273201000016C, N01-ES-45517, N01-ES-55541, N01-ES-05456, and N01-ES-65557.

Abstract

1
2 Sodium tungstate dihydrate (ST) is present naturally in the environment and can enter waterways
3 through the weathering of rocks and soils. ST also is a high-production volume compound that is
4 used in a variety of commercial applications including fire- and waterproofing fabrics, in the
5 preparation of complex compounds (e.g., phosphotungstate and silicotungstate), as a reagent for
6 biological products, and as a precipitant for alkaloids. Tungsten was nominated to the National
7 Toxicology Program (NTP) by the Centers for Disease Control and Prevention to evaluate its
8 potential to cause chronic toxicity and carcinogenicity because of concern about potential human
9 exposure via contaminated drinking water (e.g., in the form of salts like tungstate) and
10 inadequate data to assess human health implications of elevated exposures. ST was selected for
11 study because it is the most prevalent water-soluble form of tungsten. In these studies, Sprague
12 Dawley (Hsd:Sprague Dawley[®] SD[®]) rat dams were exposed to ST in drinking water from
13 gestation day (GD) 6 through lactation day (LD) 20. Their pups were exposed to the same
14 exposure concentrations in drinking water from postnatal day (PND) 12 through 3 months or
15 2 years. Adult male and female B6C3F1/N mice were exposed to ST in drinking water for
16 3 months or 2 years. Genetic toxicology studies were conducted in *Salmonella typhimurium*,
17 *Escherichia coli*, rat and mouse peripheral blood erythrocytes, and cells from liver, kidney, and
18 ileum; peripheral blood leukocytes from rats and mice also were assessed for DNA damage.

19 Perinatal and Three-month Study in Rats

20 Beginning on GD 6, groups of eight F₀ time-mated female rats were exposed to ST in drinking
21 water throughout gestation and lactation at one of five exposure concentrations (125, 250, 500,
22 1,000, or 2,000 mg/L) or were provided the vehicle control (deionized tap water). Groups of
23 10 F₁ rats per sex continued on in the study after weaning and were given drinking water
24 containing the same respective ST concentrations for 3 months. There were no significant effects
25 of ST exposure on pregnancy status, maternal survival, or littering parameters. By the end of
26 lactation, dams in the 1,000 and 2,000 mg/L groups showed significant decreases in group mean
27 body weight of approximately 10% and 18%, respectively, and water consumption was
28 significantly decreased for the 500, 1,000, and 2,000 mg/L groups relative to the vehicle control
29 group over the LD 17 to LD 21 interval. When adjusted for litter size, the mean body weight of
30 male and female pups in the 2,000 mg/L group on PND 21 was significantly decreased by
31 approximately 16% and 11%, respectively, compared to the corresponding vehicle control
32 groups.

33 There were no early deaths during the 3-month study. When compared to the vehicle control
34 group, final mean body weights were lower for the 1,000 and 2,000 mg/L males and 2,000 mg/L
35 females. Water consumption was lower for the 1,000 and 2,000 mg/L males and females. The
36 urine xanthine/creatinine ratios were significantly increased in all male and female exposed
37 groups. Serum insulin concentrations were significantly decreased in the 2,000 mg/L males
38 relative to the vehicle control males. Significantly decreased absolute weights were observed in
39 several organs but were considered secondary to body weights reductions. Exposure-related
40 histological lesions were limited to the kidneys and included increased incidences of renal tubule
41 regeneration in the 1,000 and 2,000 mg/L males and females; the increases in the 2,000 mg/L
42 groups were significant relative to the vehicle control group.

1 **Perinatal and Two-year Study in Rats**

2 Beginning on GD 6, F₀ time-mated females were exposed to ST in drinking water throughout
3 gestation and lactation at one of three exposure concentrations (250, 500, or 1,000 mg/L) or were
4 provided the vehicle control (deionized tap water). Groups of 50 F₁ rats/sex/group continued on
5 in the study after weaning and were provided drinking water containing the same respective ST
6 concentration as their dam for 2 years. An additional 40 F₁ rats/sex/exposure group were used for
7 interim evaluations and were provided dosed drinking water or the vehicle control for 3, 6, 12, or
8 18 months. There were no significant effects on reproductive performance, including the
9 percentage of mated females producing pups. During gestation and lactation, the mean body
10 weight of dams in the 1,000 mg/L group was lower than that of the vehicle control group. There
11 were no exposure-related differences between the vehicle control group and the ST-exposed
12 groups in the number of litters, litter size, mean litter weights, sex ratio, or the pup mean weights
13 of males and females.

14 Interim evaluations were performed on male and female rats at 3, 6, 12, or 18 months for organ
15 weights and tungsten concentrations in plasma, kidney, and urine. Although there was no
16 consistent pattern of changes in kidney weights across sex or over time, kidney tungsten
17 concentrations increased with exposure concentration, and the kidney/plasma ratios were higher
18 than 1 at all exposure concentrations and time points demonstrating retention of tungsten in the
19 kidney. This finding was consistent with the nephrotoxicity observed in the 2-year study.

20 Survival to study termination was significantly increased in all groups of exposed male rats
21 compared to the vehicle control males, with survival of the vehicle control males being lower
22 than that typically seen in groups of control male Sprague Dawley rats in previous 2-year NTP
23 studies. There were no significant differences in the survival of female groups. At study
24 termination, mean body weights of all groups of exposed males were within 10% of the vehicle
25 control group. In females, mean body weights of the 500 mg/L and 1,000 mg/L groups at study
26 termination were approximately 11% and 21% less than those of the vehicle control group,
27 respectively. Over the course of the 2-year study, mean water consumption for the 250, 500, and
28 1000 mg/L groups averaged 93%, 99%, and 84% of the vehicle control males and averaged 95%,
29 100%, and 91% of the vehicle control females.

30 The incidences of thyroid gland C-cell adenomas were higher in all exposed groups of female
31 rats, and the increase was significant in the 500 mg/L group relative to the vehicle control group.
32 Although not significant, the incidence of C-cell carcinomas was higher in the 1,000 mg/L
33 females. The incidences of C-cell adenoma or carcinoma (combined) exceeded the historical
34 control range in the 250 and 500 mg/L females.

35 In the kidney, the incidences of suppurative inflammation of the renal tubules were significantly
36 increased in the 1,000 mg/L males and females, and the incidence of renal tubule regeneration
37 was significantly increased in the 1,000 mg/L females, relative to the respective vehicle control
38 groups.

39 In the uterus, there was a significant increase in the incidence of atypical hyperplasia, relative to
40 the vehicle control group, in the 500 mg/L females.

41 **Three-month Study in Mice**

42 Groups of 10 male and 10 female mice were exposed to ST in drinking water for 3 months at one
43 of five exposure concentrations (125, 250, 500, 1,000, or 2,000 mg/L) or were provided the
44 vehicle control (deionized tap water). All mice survived to the end of the study. Over the course

1 of the study, mean body weights were below 90% of the vehicle control group for the 250, 1,000,
2 and 2,000 mg/L females and the 2,000 mg/L males. At study termination, the mean body weights
3 of all exposed groups of males and females were within 10% of the vehicle control groups.
4 Weekly mean water consumption values were <90% of the vehicle control groups in the 1,000
5 and 2,000 mg/L males and the 2,000 mg/L females. Lower absolute organ weights were
6 attributed to body weight reductions.

7 The only histological lesion associated with exposure was in the kidney. The incidences of renal
8 tubule regeneration were higher in the 1,000 and 2,000 mg/L male and female groups compared
9 to the respective vehicle control groups. The increases in the male groups were significant.

10 **Two-year Study in Mice**

11 Groups of 50 male and 50 female mice were exposed to ST in drinking water for 2 years at one
12 of three exposure concentrations (500, 1,000, or 2,000 mg/L) or were provided the vehicle
13 control (deionized tap water). An additional 40 mice/sex/exposure group were included for
14 interim evaluations at 3, 6, 12, and 18 months.

15 More males in the ST-exposed groups survived to study termination than did the vehicle control
16 males; however, the differences were not significant. Survival in females was similar across all
17 groups. At study termination, the mean body weight of the 2,000 mg/L males was 88% of the
18 vehicle control group, and water consumption was approximately 78% of the vehicle control
19 group; all other groups of exposed males and all groups of exposed females had mean body
20 weights within 10% of their respective vehicle control groups. Clinical observations included
21 more occurrences of thinness and ruffled fur in exposed males compared to vehicle control
22 males.

23 Kidney tungsten concentrations increased with exposure concentration and the kidney/plasma
24 ratios were higher than 1 at all exposure concentrations and time points, demonstrating retention
25 of tungsten in the kidney. Renal tubule neoplasms were only recorded in exposed males; one
26 renal tubule adenoma was observed in the 1,000 mg/L males, and two renal tubule carcinomas
27 were observed in the 2,000 mg/L males. Compared to the respective vehicle control groups, there
28 were significantly increased incidences of renal tubule regeneration in all exposed groups of
29 males and in the 1,000 and 2,000 mg/L groups of females.

30 In the large intestine, the incidences of pigment in the cecum were significantly increased in the
31 1,000 and 2,000 mg/L males and females.

32 In the testes, there was a significantly increased incidence of germinal epithelium degeneration in
33 the 500 mg/L group relative to the vehicle control group; the incidences were increased, but not
34 significant, in the 1,000 and 2,000 mg/L groups.

35 The incidence of hypercellularity of the bone marrow was significantly increased in the 500 and
36 1,000 mg/L males; the incidence of extramedullary hematopoiesis in the spleen was significantly
37 increased in the 500 and 1,000 mg/L females.

38 **Genetic Toxicology**

39 ST was not mutagenic in any of several bacterial tester strains, with or without exogenous
40 metabolic activation (S9 mix). No increases in micronucleated erythrocytes were seen in male
41 and female rats and mice administered ST in drinking water for 3 months. An exposure
42 concentration-related significant increase in the percent of circulating immature erythrocytes was

1 seen in male and female rats and in male mice, whereas there were no changes in this population
2 of cells in female mice. Significantly increased DNA damage, as measured by the comet assay
3 following administration of ST in drinking water for 3 months, was seen in liver cells of male
4 and female rats and male mice; it also was seen in cells from the ileum of male mice. No
5 increases in the levels of DNA damage were observed in blood leukocytes from either species or
6 in kidney cells from mice.

7 **Conclusions**

8 Under the conditions of these 2-year drinking water studies, there was *no evidence of*
9 *carcinogenic activity* of sodium tungstate dihydrate (ST) in male Hsd:Sprague Dawley® SD® rats
10 at exposure concentrations of 250, 500, or 1,000 mg/L. There was *equivocal evidence of*
11 *carcinogenic activity* of ST in female Hsd:Sprague Dawley® SD® rats based on increased
12 incidences of C-cell adenoma or carcinoma (combined) of the thyroid gland.

13 There was *equivocal evidence of carcinogenic activity* of ST in male B6C3F1/N mice based on
14 the occurrences of renal tubule adenoma or carcinoma (combined) in exposed animals. There
15 was *no evidence of carcinogenic activity* of ST in female B6C3F1/N mice at exposure
16 concentrations of 500, 1,000, or 2,000 mg/L.

17 Exposure to ST in drinking water caused increased incidences of nonneoplastic lesions in the
18 kidney of male and female rats and mice, in the uterus of female rats, in the testes and bone
19 marrow of male mice, and in the spleen of female mice.


20 **Synonyms:** Tungstic acid sodium salt dihydrate

1 **Summary of the Perinatal and Two-year Carcinogenesis and Genetic Toxicology Studies of Sodium**
 2 **Tungstate Dihydrate**

	Male Sprague Dawley Rats	Female Sprague Dawley Rats	Male B6C3F1/N Mice	Female B6C3F1/N Mice
Concentrations in Drinking Water	0, 250, 500, or 1,000 mg/L	0, 250, 500, or 1,000 mg/L	0, 500, 1,000, or 2,000 mg/L	0, 500, 1,000, or 2,000 mg/L
Survival Rates	12/50, 26/50, 24/50, 29/50	30/50, 33/50, 31/50, 31/50	26/50, 31/50, 35/50, 35/50	38/50, 41/50, 38/50, 40/50
Body Weights	Exposed groups similar to the vehicle control group	500 mg/L group 11% less than the vehicle control group; 1,000 mg/L group 22% less than the vehicle control group	2,000 mg/L group 12% less than the vehicle control group	Exposed groups similar to the vehicle control group
Nonneoplastic Effects	<u>Kidney</u> : renal tubule, inflammation, suppurative (25/50, 33/50, 35/50, 41/50)	<u>Kidney</u> : renal tubule, inflammation, suppurative (8/50, 9/50, 6/50, 19/50); renal tubule, regeneration (0/50, 0/50, 0/50, 18/50) <u>Uterus</u> : atypical hyperplasia (4/50, 7/50, 19/50, 8/50)	<u>Kidney</u> : renal tubule, regeneration (2/50, 21/50, 32/50, 38/50) <u>Large intestine</u> : cecum, pigment (3/50, 7/50, 17/50, 32/50) <u>Testis</u> : germinal epithelium, degeneration (11/50, 20/50, 20/50, 20/50) <u>Bone marrow</u> : hypercellularity (15/50, 35/50, 26/50, 19/50)	<u>Kidney</u> : renal tubule, regeneration (0/50, 1/50, 7/50, 7/50) <u>Large intestine</u> : cecum, pigment (0/50, 3/50, 7/50, 14/50) <u>Spleen</u> : extramedullary hematopoiesis (5/50, 18/50, 13/50, 8/50)
Neoplastic Effects	None	None	None	None
Equivocal Findings	None	<u>Thyroid gland</u> : C- cell adenoma (5/50, 13/50, 13/49, 8/50); C-cell carcinoma (2/50, 2/50, 2/49, 4/50); C-cell adenoma or carcinoma (combined) (7/50, 15/50, 14/49, 11/50)	<u>Kidney</u> : renal tubule adenoma (0/50, 0/50, 1/50, 0/50); renal tubule carcinoma (0/50, 0/50, 0/50, 2/50); renal tubule adenoma or carcinoma (combined) (0/50, 0/50, 1/50, 2/50)	None
Level of Evidence of Carcinogenic Activity	No evidence	Equivocal evidence	Equivocal evidence	No evidence

	Male Sprague Dawley Rats	Female Sprague Dawley Rats	Male B6C3F1/N Mice	Female B6C3F1/N Mice
<hr/> Genetic Toxicology				
Bacterial gene mutations: Negative in <i>Salmonella typhimurium</i> strains TA98 and TA100 and <i>Escherichia coli</i> strain WP2 <i>uvrA</i> pKM101 with and without S9				
Micronucleated erythrocytes (in vivo)				
Rat peripheral blood: Negative in males and females				
Mouse peripheral blood: Negative in males and females				
DNA damage				
Rat: Positive in liver (males and females); negative in leukocytes (males and females) and ileum (females); not reported in ileum (males) and kidney (males and females)				
Mouse: Positive in liver and ileum (males); negative in liver and ileum (females); negative in kidney and leukocytes (males and females)				

1 Introduction



2
3 **Figure 1. Sodium Tungstate Dihydrate (CASRN 10213-10-2; Chemical Formula: $\text{Na}_2\text{WO}_4 \cdot 2\text{H}_2\text{O}$;**
4 **Molecular Weight: 329.86)**

5 Synonyms: Tungstic acid sodium salt dihydrate.

6 Chemical and Physical Properties

7 Tungsten, also called wolfram, is a steel-gray to tin-white metal with a high melting point
8 and good electrical conductivity. Along with chromium and molybdenum (Mo), it is in
9 group VI of the periodic table. It can replace Mo in Mo-containing enzymes,^{1;2} such as aldehyde
10 oxidase, sulfite oxidase,³ and xanthine oxidase,⁴ and renders the enzymes inactive. Sodium
11 tungstate dihydrate (ST) is a chemical intermediate for tungsten and tungsten compounds.⁵ ST
12 has high solubility in water and is not volatile. It effloresces in dry air and loses its water at
13 100°C. As an aqueous solution, it is slightly alkaline (pH 8–9). When heated to
14 decomposition, it emits toxic fumes of sodium oxide.

15 Production, Use, and Human Exposure

16 Tungsten and its salts are present naturally in the environment and can enter waterways through
17 the weathering of rocks and soils.⁶ Atmospheric tungsten-containing particulates eventually
18 settle to the earth's surface by dry deposition or can be removed from the atmosphere by wet
19 deposition (i.e., precipitation). Upon reaching water and soil, tungsten will be in either soluble
20 (e.g., tungstate ion, WO_4^{2-}) or insoluble forms (e.g., tungsten trioxide) in sediment and soil.
21 Environmental exposure to ST by the general public occurs mainly through contaminated
22 drinking water. For example, tungsten has been detected in the municipal water of Fallon,
23 Nevada. However, the amount of tungsten in drinking water is generally not known.⁶

24 ST is a high-production volume compound and is produced for industrial purposes by the
25 reaction of a mixture of soft and hard tungsten carbide when combined with a mixture of sodium
26 nitrate and sodium hydroxide in a fusion process. ST is used in a variety of commercial
27 applications including fire- and waterproofing fabrics, in the preparation of complex compounds
28 (e.g., phosphotungstate and silicotungstate), as a reagent for biological products, and as a

1 precipitant for alkaloids. Occupational exposure can occur through inhalation of dusts and
2 dermal contact during the production or use of tungsten-containing compounds.^{6;7}

3 Clinically, ST is used as an antidiabetic agent to improve pancreatic function through a
4 combination of hyperglycemia-independent pathways and by its own direct and indirect effects^{8;}
5 ⁹; it is also used to treat infertility in people with diabetes.¹⁰

6 **Regulatory Status**

7 The American Conference of Governmental Industrial Hygienists limits for tungsten and its
8 soluble compounds include a time-weighted average (TWA) air concentration of 1 mg/m³ and a
9 short-term exposure limit (STEL) of 3 mg/m³. The National Institute for Occupational Safety and
10 Health (NIOSH) recommends a 10-hour TWA air concentration of 1 mg/m³ for tungsten and its
11 soluble compounds. When developing a final rule, the Occupational Safety and Health
12 Administration (OSHA) proposed an 8-hour TWA permissible exposure limit of 1 mg/m³ and a
13 15-minute STEL of 3 mg/m³ for tungsten. NIOSH concurred with OSHA's addition of the
14 STEL, and therefore the final rule established limits for tungsten and its soluble compounds of
15 1 mg/m³ as an 8-hour TWA and 3 mg/m³ as a 15-minute STEL, measured as tungsten.¹¹ No
16 federal drinking water standard or ambient water quality criterion have been established for
17 tungsten.¹²

18 **Absorption, Distribution, Metabolism, and Excretion**

19 **Experimental Animals**

20 The National Toxicology Program (NTP) previously evaluated the disposition of tungsten
21 (administered as ST) in female Sprague Dawley rats or C57BL/6N mice.^{13;14} Following a single
22 gavage administration of 1, 10, or 100 mg ST/kg body weight (mg/kg), animals were euthanized
23 1, 2, 4, and 24 hours postadministration.¹³ Plasma, liver, kidney, femur, and uterus
24 concentrations of tungsten increased with increasing dose in both species, with higher tissue
25 concentrations relative to plasma at each respective time point. Tissue concentrations were
26 generally higher in rats relative to mice, with some variation by tissue and time point. In general,
27 tungsten concentrations peaked approximately 4 hours postadministration in rats and 1–4 hours
28 postadministration in mice; concentrations then decreased over time with values approaching
29 background/endogenous concentrations detected in the study (0.04–0.15 µg/g, depending on the
30 matrix) by 24 hours postadministration. After a single intravenous administration of ST at
31 1 mg/kg in rats and mice, concentrations of tungsten peaked at approximately 1 hour
32 postadministration and steadily decreased through 24 hours postadministration.¹³

33 After 14 days of gavage administration of ST to rats and mice (10 mg/kg; animals necropsied
34 24 hours after the last administration), tungsten concentrations in plasma, liver, kidney, femur,
35 and uterus were either similar to or slightly higher than those in animals receiving a single
36 administration of a similar dose and necropsied 24 hours postadministration; this observation
37 suggests minimal accumulation of tungsten following repeat gavage dosing in rats and mice.¹⁴
38 After exposure via drinking water for 14 days (560 mg/L; dosed water offered until study
39 termination) in rats and mice, tungsten was detected in plasma and tissues; however, a direct
40 comparison between repeat gavage dosing and drinking water exposure cannot be established
41 due to the differences in the time of necropsy after the last exposure. In pregnant rats and mice,

1 following a similar drinking water exposure paradigm, tungsten was also detected in the fetus,
2 which demonstrated gestational transfer.¹⁴

3 **Humans**

4 Studies of human exposure to tungsten compounds are limited. The daily dietary intake of
5 tungsten is about 0.01 mg (0.05 μmol), whereas the median for daily urinary excretion is
6 0.007 mg (0.04 μmol).¹⁵ Tungsten (VI) is well absorbed, and approximately 75% of the amount
7 ingested is excreted in the urine.¹⁶ In a limited study with no specific exposure, four healthy
8 young adults eliminated trace quantities of tungsten in urine (2.0–13.0 μg [0.01–0.07 μmol]) and
9 feces (1.6–5.7 μg [8.7–31 nmol]) over 24-hour periods.¹⁷

10 **Toxicity**

11 **Experimental Animals**

12 Acute toxicity values (i.e., median lethal dose [LD₅₀]) for ST range from 240 to 1,904.1 mg/kg in
13 mice and from 1,904 to 1,928 mg/kg in rats.^{18; 19} Acute oral or intravenous administration of ST
14 in mice and rats decreased motor activity and muscle tone, and the animals exhibited ataxia,
15 palpebral ptosis, hunched back, pallor, prostration, and dyspnea.¹⁸

16 In a relatively recent study, the subchronic toxicity of an aqueous ST solution in male and female
17 Sprague Dawley rats was evaluated following daily administration via oral gavage to 0, 10, 75,
18 125, or 200 mg ST/kg body weight/day (mg/kg/day) for 90 days.²⁰ The kidney was noted as the
19 main target organ of toxicity in both male and female rats dosed at 125 or 200 mg/kg/day, with
20 mild to severe cortical tubule basophilia observed in those dose groups. In males, intraluminal
21 hypospermia with cell debris was observed in the epididymis after administration of
22 200 mg/kg/day. In both sexes, histopathological changes were observed in the glandular stomach
23 and included inflammation and metaplasia in the high-dose rats (125 and 200 mg/kg/day). The
24 histopathological effects seen in the kidneys indicate that the lowest-observed-adverse-effect
25 level from this study was 125 mg/kg/day and the no-observed-adverse-effect level was
26 75 mg/kg/day in both sexes of rats for oral subchronic toxicity. There was a significant decrease
27 in feed consumption and body weight gain in males at 200 mg/kg/day from days 77 to 90;
28 however, there was no effect on feed consumption and body weight in females. There were no
29 changes in the hematological or clinical parameters in this study. Histopathological changes were
30 seen in the kidney of male and female rats and in the epididymis of male rats.

31 In an older study, ST (equivalent to 2% tungsten) administered via the diet to young rats caused
32 the death of all animals within 10 days.^{17; 21} When dietary concentrations were reduced to an
33 equivalent of 0.5% tungsten, death occurred in 75% of rats by the end of the 70-day exposure
34 period. When given by gavage or in drinking water to young rats, ST (15–1,000 mg/kg/day
35 [0.051–3.403 mmol/kg/day]) for 4 or 13 weeks produced emesis, anorexia, cachexia, pallor, and
36 dyspnea.^{22; 23} At the highest dose, concentrations of urea, creatinine, and total cholesterol were
37 increased, whereas erythrocyte count, glucose, aspartate aminotransferase/alanine
38 aminotransferase, protein, hematocrit, and hemoglobin levels were decreased. All parameter
39 values returned to normal after a recovery period of 6 weeks. Another study in male rats noted
40 effects on spermatogenesis after inhalation exposure to ST (504 $\mu\text{g}/\text{m}^3$ [41.9 ppb]) for
41 24 hours/day for 17 weeks.²⁴

1 In a 28-day study of B6C3F1/N mice exposed to ST via drinking water at concentrations of
2 125–2,000 mg/L, NTP found limited effects on humoral and innate immunity, on developing
3 hematopoietic cells in the bone marrow, and on unstimulated splenocyte phenotypes. These data
4 indicated that, under conditions of co-exposure to an immune-stimulating agent, such as tumor
5 cells or genetically dissimilar leukocytes, ST may modulate the normal cell-mediated immune
6 response.²⁵

7 Chronic oral exposure to 5 ppm ST in drinking water has been shown to significantly reduce
8 longevity in male Long-Evans rats.²⁶ In male Wistar rats, daily gavage administration of less
9 than 150 mg/kg for up to 300 days produced no significant effects on body weights, organ
10 weights, or survival.²⁷

11 **Humans**

12 The data implicating ST as toxic, hazardous, or carcinogenic in humans are limited, although
13 tungsten poisoning has been reported following continuous occupational exposure to dusts and
14 vapors during the refining of tungsten metal.²⁸

15 **Reproductive and Developmental Toxicity**

16 **Experimental Animals**

17 Although some reproductive and teratological effects have been previously reported in rats and
18 mice exposed to tungsten, those studies are not well characterized. In mice, a single dose of ST
19 (concentration not specified) provided to dams at early fetal organogenesis was shown to
20 produce a high frequency of resorptions but did not induce any fetal malformations.²⁹ In
21 pregnant rats, doses that did not produce maternal toxicity increased embryo lethality and
22 inhibited bone ossification in fetuses.³⁰ Another study evaluated the reproductive and
23 neurobehavioral effects of ST in Sprague Dawley rats after 70 days of gavage administration
24 with 0, 5, 62.5, or 125 mg/kg/day starting prior to mating and continuing through gestation and
25 weaning (postnatal day 20). The perinatally exposed offspring showed subtle neurobehavioral
26 effects related to motor activity and emotionality.³¹

27 **Humans**

28 The literature contains no studies about reproductive or developmental toxicity in humans
29 following exposure to tungsten or tungsten compounds.

30 **Carcinogenicity**

31 The literature contains no carcinogenicity studies of ST in experimental animals or epidemiology
32 studies in humans.

33 **Genetic Toxicity**

34 Few reports have been published on the genotoxicity of ST, but the available data suggest that
35 the compound is not genotoxic. ST did not induce morphological transformation of Syrian
36 hamster embryo (SHE) cells exposed in culture or in a host-mediated in vivo/in vitro assay, in
37 which cells from embryos excised from Syrian golden hamster dams administered 2.5 or 5 mg

1 ST per 100 g maternal weight (intraperitoneal injection) were grown in a transformation assay.³²
2 Chromosomal aberrations and sister chromatid exchanges were not induced in human
3 lymphocytes or in SHE cells exposed to ST up to 10 µg/mL.³³

4 **Study Rationale**

5 Tungsten was nominated to NTP by the Centers for Disease Control and Prevention to evaluate
6 its potential to cause chronic toxicity and carcinogenicity because of concern about potential
7 human exposure via contaminated drinking water and inadequate data to assess human health
8 implications of elevated exposures. ST was selected for study because tungstate (WO_4^{2-}) is the
9 most naturally occurring form of soluble tungsten, and ST was the most water-soluble form of
10 tungstate.

11 In the studies described in this Technical Report, drinking water was used as the route of
12 exposure to mimic human exposure.

1 **Materials and Methods**

2 **Procurement and Characterization of Sodium Tungstate** 3 **Dihydrate**

4 Sodium tungstate dihydrate (ST) was procured from Sigma-Aldrich (St. Louis, MO) in two lots
5 (lot 12330JO and lot MKBG9975V). Lot 12330JO was obtained directly from Sigma-Aldrich
6 (St. Louis, MO), whereas lot MKBG9975V was produced by Sigma-Aldrich and obtained from
7 Government Scientific Source, Inc. (Reston, VA). Identity, purity, and stability analyses were
8 conducted by the analytical chemistry laboratory and study laboratory at Battelle (Columbus,
9 OH) (Appendix A). Reports on analyses performed in support of the ST studies are on file at the
10 National Institute of Environmental Health Sciences (NIEHS).

11 Lots 12330JO and MKBG9975V were white solids composed of fine crystals. The 3-month
12 studies used lot 12330JO. For the 2-year studies, the remainder of lot 12330JO was combined
13 with lot MKBG9975V to create lot 07072011.

14 The identities of the lots were confirmed using infrared spectroscopy. X-ray diffraction patterns
15 were in good agreement with reference standards, and proton-induced x-ray emission
16 spectroscopy yielded expected percent weights of tungsten and sodium. Magnesium (0.7–0.9 %) and
17 aluminum (approximately 0.3%) impurities were identified in both lots. The purities of
18 lots 12330JO and 07072011 were both determined to be approximately 99% using inductively
19 coupled plasma atomic emission spectrometry based on weight percentages of tungsten (55.2–
20 56.4%) and sodium (13.4–13.8%). Karl Fisher titration yielded a water content of 9.5% for
21 lot 12330JO and 10.0–10.3% for lot 07072011, slightly lower than the anticipated 10.9%.
22 Titration of tungstate ion with lead nitrate indicated a purity of 97.6% for lot 12330JO and 98.2%
23 for lot 07072011. Ion chromatography (IC) with a suppressed conductivity detector and liquid
24 chromatography with an inductively coupled plasma-mass spectrometer indicated a purity of
25 100% for both lots.

26 Accelerated stability studies were conducted on lot 12330JO and lot MKBG9975V using IC with
27 a suppressed conductivity detector. Stability was confirmed for at least 2 weeks when ST was
28 stored in sealed amber glass bottles at 25°C, 5°C, and –20°C. Therefore, bulk ST was stored in
29 sealed amber glass bottles at 25°C. Periodic analyses of the bulk chemical were conducted
30 during the 3-month and 2-year studies by the study laboratory, and no degradation of the bulk
31 chemical was detected.

32 **Preparation and Analysis of Dose Formulations**

33 The feed (NIH-07 and NTP-2000), tap water, and deionized water used in the 3-month and
34 2-year studies were analyzed for tungsten and molybdenum concentrations. NIH-07 feed
35 contained approximately 2 ppm tungsten and the concentration in NTP-2000 feed was at the
36 detection limit of the assay (0.80 ppm). Concentrations of tungsten in the tap water and
37 deionized water, and the concentration of molybdenum in all feed and water samples, were
38 below the limits of detection of the assay (0.20 to 0.80 ppm).

39 Stability studies conducted on the 20 mg/L formulation by the analytical chemistry laboratory
40 found that the formulation was stable when sealed and stored in Nalgene bottles for 42 days at

1 5°C and at room temperature (approximately 25°C). An animal room simulation was conducted
2 using the 20 mg/L formulation stored in a drinking water bottle with aliquots periodically
3 removed to simulate animal drinking. There was no significant loss in tungsten over 7 days at
4 room temperatures.

5 Dose formulations of ST were prepared monthly (Table A-2). Dose formulations were prepared
6 in deionized water, with the exception of the first two formulations used in the 3-month studies,
7 which used tap water instead. The 3-month study dose formulations were 0, 125, 250, 500,
8 1,000, and 2,000 mg/L for both mice and rats. The 2-year mouse study used 0, 500, 1,000 and
9 2,000 mg/L dose formulations, whereas the 2-year rat study used 0, 250, 500, and 1,000 mg/L
10 dose formulations. Dose formulations were stable for 42 days at room temperature
11 (Appendix A).

12 Preadministration and postadministration (animal room) analyses of dose formulations were
13 conducted monthly throughout the 3-month studies (Table A-3, Table A-4). During the 2-year
14 studies, preadministration dose formulations were analyzed every 1–3 months, whereas
15 postadministration dose formulations were analyzed every 6–8 months (Table A-5, Table A-6).
16 All preadministration formulations in the 3-month rat and mouse studies were within 10% of the
17 target concentration. In the 3-month mouse study, four postadministration samples were more
18 than 10% below the target concentration, with the largest difference being 12.8% below the
19 target. Three postadministration samples collected from carboys or bottles for the 125, 500, and
20 2,000 mg/L dose formulations in the 3-month rat study were 10.8% to 12.3% below the
21 corresponding target concentration. In the 2-year studies, all preadministration and
22 postadministration samples were within 10% of the target concentration.

23 **Animal Source**

24 Time-mated (F₀) female Sprague Dawley (Hsd:Sprague Dawley[®] SD[®]) rats were obtained from
25 Harlan Inc. (now Envigo, Indianapolis, IN) for the 3-month and 2-year studies. B6C3F1/N mice
26 were obtained from Taconic Biosciences, Inc. (Germantown, NY) for the 3-month and 2-year
27 studies.

28 **Animal Welfare**

29 Animal care and use were in accordance with the Public Health Service Policy on Humane Care
30 and Use of Animals. All animal studies were conducted in an animal facility accredited by
31 AAALAC International. Studies were approved by the Battelle (Columbus, OH) Animal Care
32 and Use Committee and conducted in accordance with all relevant National Institutes of Health
33 (NIH) and National Toxicology Program (NTP) animal care and use policies and applicable
34 federal, state, and local regulations and guidelines.

35 **Three-month Studies**

36 **Initial Exposure Concentration Selection Rationale**

37 For the perinatal and 3-month study in rats and the 3-month study in adult mice, selection of
38 exposure concentrations of 125, 250, 500, 1,000, and 2,000 mg/L ST was informed by a study
39 evaluating the antidiabetic effects of ST in rats following 8 months of exposure via drinking
40 water.³⁴

1 Study Design for Rats

2 F₀ female rats were 11 to 12 weeks old upon receipt. Gestation day (GD) 1 was defined as the
3 first day with evidence of mating. F₀ females were received on GD 1 and held for 5 days.

4 F₀ females were randomly assigned to one of six exposure groups on GD 5 (eight dams/group).
5 Randomization was stratified by body weight that produced similar group mean weights using
6 PATH/TOX SYSTEM software (Xybion Medical Systems Corporation, Cedar Knolls, NJ).

7 F₀ females were quarantined for 39 days after receipt. Ten nonmated females received in the
8 same shipment as the time-mated dams were designated for disease monitoring and were used
9 for gross necropsies 2 days after arrival; samples were collected for the presence of disease. The
10 health of the F₁ animals was monitored during the study according to the protocols of the NTP
11 Sentinel Animal Program (Appendix C). All test results were negative.

12 Beginning on GD 6, groups of eight F₀ time-mated females were provided ST in drinking water
13 throughout gestation and lactation at one of five exposure concentrations (125, 250, 500, 1,000,
14 or 2,000 mg/L) or the vehicle control (deionized tap water). Groups of 10 F₁ rats per sex
15 continued on in the study after weaning and were provided drinking water containing the same
16 respective ST concentration for 3 months.

17 F₀ female rats were housed individually during gestation and with their respective litters during
18 lactation. Feed and dosed water were available ad libitum. Dam body weights were recorded on
19 GDs 5, 6, 9, 12, 15, 18, and 21 and on lactation days (LDs) 1, 4, 7, 14, and 21. During gestation,
20 water consumption was measured over 3-day intervals from GD 6 through GD 21 (GDs 6–9, 9–
21 12, 12–15, 15–18, and 18–21). The day of parturition was considered to be postnatal (PND) 0.
22 On apparent GD 25, all time-mated females that failed to deliver were euthanized and the uteri
23 were examined and stained for evidence of implantation. Total litter weight and litter weights by
24 sex were collected on PND 1. Individual pup weights were recorded on PNDs 4, 7, 14, and 21.
25 On PND 1, clinical observations, including general appearance were recorded. Pup survival was
26 evaluated and recorded. During lactation, water consumption was measured over 3-day intervals
27 from PND 1 through PND 21 (PNDs 1–4, 4–7, 7–10, 10–14, 14–17, 17–21).

28 F₁ litters were standardized on PND 4 to eight pups/litter, with at least two pups of each sex and
29 a preference for four males and four females each. Litters that did not meet the minimum of eight
30 pups (or if they had fewer than two pups of either sex) were removed from the study. On the day
31 the last litter reached PND 20, pups were randomly assigned to the 3-month study. For all
32 exposure concentrations, except the 2,000 mg/L group, two pups per sex from five randomly
33 selected litters per exposure group were chosen. For the 2,000 mg/L group, a third male pup was
34 selected from two of the four available litters and a third female pup was selected from the other
35 two litters to obtain the complete number of animals needed for the study. After assignments to
36 the 3-month study were complete, five pups per sex from the remaining vehicle control pups
37 were randomly selected as the end-of-study sentinel animals. On the day the last litter reached
38 PND 21, dams were removed, and the pups were weaned. Weaning marked the beginning of the
39 3-month study.

40 After weaning, F₁ rats were housed five per cage. Feed and dosed water were available ad
41 libitum. Water consumption was measured weekly for 3 months. Cages were changed weekly
42 though PND 4, then changed twice weekly. Racks were changed and rotated at least every
43 2 weeks. Further details of animal maintenance are given in Table 1.

1 Two diets were used in the rat studies: (1) NIH-07 during the perinatal phase, and (2) NTP-2000
2 during the postweaning phase. The NIH-07 diet is a higher protein diet that supports
3 reproduction and lactation in rodents, whereas the NTP-2000 diet is a lower protein diet that
4 decreases the incidence of chronic nephropathy in adult rats. Information on feed composition
5 and contaminants for both diets is provided in Appendix B.

6 Because tungsten is capable of replacing molybdenum (Mo) in Mo-containing enzymes, NTP
7 evaluated the enzyme activity of xanthine oxidase and sulfite oxidase in the liver, kidney, and
8 intestine. Also, because tungsten has been shown to accumulate in femurs of rats and mice after
9 repeated oral gavage administration, urinary calcium and phosphorus concentrations were
10 measured. Other endpoints in the urine, including acetyl glucosaminidase, alkaline phosphatase,
11 and aspartate aminotransferase activities, were also measured.

12 Urine and blood were also analyzed for tungsten concentrations using validated analytical
13 methods as described in Appendix E.

14 **Study Design for Mice**

15 Male and female B6C3F1/N mice were 3 to 4 weeks old upon receipt and were quarantined for
16 11 days before study start. Mice were randomly assigned to one of six exposure groups
17 ($n = 10$ mice/sex/group). Randomization was stratified by body weight that produced similar
18 group mean weights using PATH/TOX SYSTEM software (Xybion Medical Systems
19 Corporation, Cedar Knolls, NJ). Mice were provided ST in drinking water for 3 months at one of
20 five exposure concentrations (125, 250, 500, 1,000, or 2,000 mg/L) or were provided the vehicle
21 control (deionized tap water).

22 Five male and five female mice were randomly selected for parasite evaluation and gross
23 observation of disease. The health of the mice was monitored during the study according to the
24 protocols of the NTP Sentinel Animal Program (Appendix C). All test results were negative.

25 Mice were housed individually (males) or five per cage (females). Feed and dosed water were
26 available ad libitum. Water consumption was measured weekly for 3 months. Cages were
27 changed at least once weekly (males) or twice weekly (females) and rotated every 2 weeks.
28 Racks were changed and rotated every 2 weeks. Further details of animal maintenance are in
29 Table 1. Information on feed composition and contaminants is given in Appendix B.

30 **Clinical Examinations and Pathology**

31 In the 3-month studies in rats and mice, animals were observed twice daily for signs of morbidity
32 and moribundity and were weighed before dosed water administration on day 1, weekly for
33 3 months, and at study termination. Clinical observations were recorded weekly and at study
34 termination. Water consumption was recorded weekly throughout the study.

35 At week 12, all F₁ rats were placed in metabolism cages and urine samples were collected during
36 a 16-hour overnight period for urinalysis. Rats were fasted during the collection period and had
37 access to untreated deionized water while in the metabolism cages. The parameters evaluated are
38 listed in Table 1. Once all urine parameters had been determined, 1 mL of urine from each rat
39 was designated for tungsten analysis, frozen at approximately -20°C , and shipped to Battelle

1 Toxicology Northwest (Richland, WA) for analysis. From the remaining urine, a minimum of
2 1 mL was frozen at approximately -20°C for xanthine/methionine analysis.

3 Blood was collected from the retroorbital plexus (rats) or sinus (mice) at the end of the 3-month
4 studies for hematology, clinical chemistry (rats only), erythrocyte micronuclei determination,
5 tungsten determination, serum retention for insulin determination (rats only), and for the comet
6 assay. Animals were anesthetized with a carbon dioxide/oxygen mixture and bled in a random
7 order. Blood was collected into tubes containing ethylenediaminetetraacetic acid (EDTA) (for
8 hematology, erythrocyte micronuclei, and tungsten determination/comet assay) or into serum
9 separator tubes (for clinical chemistry). Hematology parameters were analyzed using an Advia[®]
10 120 system (Bayer Diagnostics Division, Tarrytown, NY). Clinical chemistry parameters were
11 analyzed using the Roche cobas c501 Chemistry Analyzer (Roche Diagnostics, Indianapolis, IN).
12 The parameters measured are listed in Table 1. After evaluation of clinical chemistry parameters,
13 the remaining rat serum was frozen at approximately -80°C and shipped to the NTP Frozen
14 Tissue Bank (Durham, NC) for serum insulin analysis. Samples for erythrocyte micronuclei
15 determination were stored at 2°C – 8°C immediately after collection and shipped to Integrated
16 Laboratory Systems, LLC (Durham, NC) for analysis. Samples for the comet assay were
17 transferred to a cryogenic vial, frozen in liquid nitrogen, and stored at -80°C for at least 24 hours
18 before shipment to Integrated Laboratory Systems, LLC (Durham, NC) for analysis. Remaining
19 blood (not used for the comet assay) was designated for tungsten determination, frozen at
20 approximately -80°C , and shipped to Battelle Toxicology Northwest (Richland, WA).

21 At the end of the 3-month studies, samples were collected for sperm motility and vaginal
22 cytology evaluations from F₁ male and female rats and from male and female mice in the 0, 500,
23 1,000, and 2,000 mg/L groups. The parameters evaluated are listed in Table 1. Due to
24 inconsistent sample collection and slide staining, an assessment of estrous cyclicity could not be
25 made for F₁ female rats or female mice. Male animals were evaluated for sperm count and
26 motility. The left testis and left epididymis were isolated and weighed. The tail of the epididymis
27 (cauda epididymis) was then removed from the epididymal body (corpus epididymis) and
28 weighed. Test yolk (rats) or modified Tyrode's buffer (mice) was applied to slides, and a small
29 incision was made at the distal border of the cauda epididymis. The sperm effluxing from the
30 incision were dispersed in the buffer on the slides, and the numbers of motile and nonmotile
31 spermatozoa were counted for five fields per slide by two observers. After completion of sperm
32 motility estimates, each left cauda epididymis was placed in buffered saline solution. Caudae
33 were finely minced, and the tissue was incubated in the saline solution and then heat fixed at
34 65°C . Sperm density was determined microscopically with the aid of a hemocytometer. To
35 quantify spermatogenesis, the testicular spermatid head count was determined by removing the
36 tunica albuginea and homogenizing the left testis in phosphate-buffered saline containing 10%
37 dimethyl sulfoxide. Homogenization-resistant spermatid nuclei were counted with a
38 hemocytometer.

39 Necropsies were performed on all rats and mice at the end of the 3-month study. Organ weights
40 were recorded for the liver, thymus, right kidney, right testis, heart, and lungs. Tissue samples of
41 3–5 mm were collected from the left lateral liver lobe, ileum, and a longitudinal section of the
42 left kidney for the comet assay. These samples were transferred to cryogenic vials, frozen in
43 liquid nitrogen, and stored at -80°C for at least 24 hours before shipment to Integrated
44 Laboratory Systems, LLC (Durham, NC) for analysis. Tissues for microscopic examination were
45 fixed in 10% neutral buffered formalin (except eyes, which were first fixed in Davidson's

1 solution, and testes, vaginal tunics, and epididymides, which were first fixed in modified
2 Davidson's solution), processed and trimmed, embedded in paraffin, sectioned to a thickness of
3 4 to 6 μm , and stained with hematoxylin and eosin (H&E). Complete histopathological
4 examinations were performed by the study laboratory pathologist on all organs with gross lesions
5 and on all vehicle control and 2,000 mg/L rats and mice. The kidney was identified as a target
6 organ and examined to a no-effect level. Table 1 lists the tissues and organs routinely examined.

7 After a review of the laboratory reports and selected histopathology slides by a quality
8 assessment (QA) pathologist, the findings and reviewed slides were submitted to an NTP
9 Pathology Working Group (PWG) coordinator for a second independent review. Any
10 inconsistencies in the diagnoses made by the study laboratory and QA pathologists were resolved
11 by the NTP pathology peer-review process. Final diagnoses for reviewed lesions represent a
12 consensus of the PWG or a consensus between the study laboratory pathologist, NTP
13 pathologist, QA pathologist(s), and the PWG coordinator. Details of these review procedures
14 have been described, in part, by Maronpot and Boorman³⁵ and Boorman et al.³⁶

15 **Two-year Studies**

16 **Study Design for Rats**

17 F₀ female rats were 11 to 14 weeks old upon receipt. GD 1 was defined as the first day with
18 evidence of mating. F₀ females were received on GD 2 and held for 4 days. F₀ females were
19 randomly assigned to one of four exposure groups on GD 5. Forty-seven females were assigned
20 to the 0 mg/L group, whereas 41 females were assigned to each of the 250, 500, and 1,000 mg/L
21 groups. Randomization was stratified by body weight that produced similar group mean weights
22 using PATH/TOX SYSTEM software (Xybio Medical Systems Corporation, Cedar Knolls,
23 NJ).

24 F₀ females were quarantined for 23 days after receipt. Nonmated females designated for disease
25 monitoring and gross necropsies were not received, and therefore 10 undelivered dams were used
26 instead for disease monitoring and gross necropsies on GD 25; samples were collected for the
27 presence of disease or parasites. The health of the F₁ animals was monitored during the study
28 according to the protocols of the NTP Sentinel Animal Program (Appendix C). Pinworms
29 (*Syphacia* spp.) were diagnosed in sentinel animals during routine health monitoring evaluations.
30 Infected animals did not display clinical signs, and no pathological lesions were noted in relation
31 to the presence of the pinworms. In accordance with this finding, NTP, in coordination with the
32 testing laboratory, developed and implemented a successful plan of pinworm containment and
33 eradication. NTP required the testing laboratories to actively monitor animals to ensure the
34 continued exclusion of pinworms from all studies going forward. All other test results were
35 negative.

36 Beginning on GD 6, F₀ females were provided ST in drinking water throughout gestation and
37 lactation at one of four exposure concentrations (0, 250, 500, or 1,000 mg/L); deionized tap
38 water served as the vehicle control. Groups of 50 F₁ rats/sex/exposure group continued on in the
39 study after weaning and were provided drinking water containing the same respective ST
40 concentration as their dam for 2 years. An additional 40 F₁ rats/sex/exposure group were used for
41 interim evaluations and provided dosed drinking water for 3, 6, 12, and 18 months.

1 F₀ female rats were housed individually during gestation and with their respective litters during
2 lactation. Feed and dosed water were available ad libitum. Dam body weights were recorded on
3 GDs 2, 5, 6, 9, 12, 15, 18, and 21 and on LDs 1, 4, 7, 10, 14, 17, and 21. During gestation, water
4 consumption was continuously measured over 3-day intervals from GD 6 through GD 21 (GDs
5 6–9, 9–12, 12–15, 15–18, and 18–21). The day of parturition was considered to be PND 0. On
6 apparent GD 25, all time-mated females that failed to deliver were euthanized and the uteri were
7 examined and stained for evidence of implantation. Total litter weight and litter weights by sex
8 were collected on PND 1. Individual pup weights were recorded on PNDs 4, 7, 10, 14, and 21.
9 Clinical observations and survival were evaluated throughout lactation. During lactation, water
10 consumption was measured over 3-day intervals from PND 1 through PND 21 (PNDs 1–4, 4–7,
11 7–10, 10–14, 14–17, 17–21).

12 F₁ litters were standardized on PND 4 to eight pups/litter, with at least two pups of each sex and
13 a preference for four males and four females each. Litters that did not meet the minimum of eight
14 pups (or if they had fewer than two pups of either sex) were removed from the study. Before
15 weaning, pups (generally two/sex/litter) were randomly assigned to the 2-year study. After
16 assignments to the 2-year study were complete, five pups per sex from the remaining vehicle
17 control pups were randomly selected as the sentinel animals. On the day the last litter reached
18 PND 21, dams were removed, and the pups were weaned. Weaning marked the beginning of the
19 2-year study.

20 On the morning of the final PND 21, randomly selected dams (five/exposure group) and one
21 male and one female pup from each selected dam's litter were used for biological sample
22 collection. Blood was collected via cardiac puncture into tubes containing EDTA, the tubes were
23 centrifuged, and the resulting plasma was harvested and stored at –85°C to –60°C until
24 transferred for tungsten analysis.

25 F₁ rats were housed up to two (males) or five (females) per cage. Feed and dosed water were
26 available ad libitum. Water consumption was measured at the beginning of the study, weekly for
27 13 weeks, and then at 4-week intervals thereafter. Cages were changed weekly through PND 4,
28 then changed twice weekly. Racks were changed and rotated at least every 2 weeks. Further
29 details of animal maintenance are given in Table 1. Information on feed composition and
30 contaminants is provided in Appendix B.

31 **Study Design for Mice**

32 Male and female B6C3F1/N mice were 3 to 4 weeks old upon receipt and were quarantined for
33 11 days before study start. Mice were randomly assigned to one of four groups (n = 50
34 mice/sex/exposure group). Mice were provided ST in drinking water for 2 years at one of three
35 exposure concentrations (500, 1,000, or 2,000 mg/L) or were given the vehicle control
36 (deionized tap water). An additional 40 mice/sex/exposure group were included for interim
37 evaluations at 3, 6, 12, and 18 months. Randomization was stratified by body weight that
38 produced similar group mean weights using PATH/TOX SYSTEM software (Xybio Medical
39 Systems Corporation, Cedar Knolls, NJ).

40 Before study start, five male and five female mice were randomly selected for parasite evaluation
41 and gross observation of disease. The health of the mice was monitored during the study

1 according to the protocols of the NTP Sentinel Animal Program (Appendix C). All test results
2 were negative.

3 Mice were housed individually (males) or four (females) per cage. Feed and dosed water were
4 available ad libitum. Water consumption was measured at the beginning of the study, weekly for
5 13 weeks, and then at 4-week intervals thereafter. Cages were changed at least once weekly
6 (males) or twice weekly (females) and rotated every 2 weeks. Racks were changed and rotated
7 every 2 weeks. Further details of animal maintenance are given in Table 1. Information on feed
8 composition and contaminants is given in Appendix B.

9 **Clinical Examinations, Tungsten Concentrations, and Pathology**

10 In the 2-year studies in rats and mice, animals were observed twice daily for signs of morbidity
11 and moribundity and were weighed before dosed water administration on day 1, weekly for the
12 next 13 weeks, every 4 weeks thereafter, and at study termination. Clinical observations were
13 recorded every 4 weeks beginning on day 36 and at study termination. Water consumption was
14 recorded at the beginning of the study, weekly for 13 weeks, and at 4-week intervals thereafter.

15 At the 3-, 6-, 12-, and 18-month interim evaluations, urine, feces, blood, and tissues (liver,
16 kidneys, stomach, small intestine, and bone) were collected from up to 10 predesignated
17 F₁ rats/sex/exposure group and up to 10 predesignated mice/sex/exposure group for
18 determination of tungsten concentrations. Organ weights were recorded for selected tissues.
19 Early death animals were not replaced. On the morning of the day before scheduled blood
20 collection, animals were moved to metabolism cages (one animal per cage); while in the
21 metabolism cages, the animals had ad libitum access to feed and their assigned concentration of
22 dosed drinking water. Urine and feces were collected over a 24-hour period, and urine volume,
23 urine creatinine, and fecal weights were recorded. Blood was collected via cardiac puncture into
24 tubes containing K₃ EDTA, centrifuged, and the plasma harvested. Immediately after blood
25 collection, the animals were euthanized and the entire liver, both kidneys, stomach (separated
26 into glandular and non-glandular), small intestine, and both femurs were collected, weighed, and
27 maintained on dry ice until moved into storage. All samples were stored at -85°C to -60°C until
28 transferred for analysis.

29 Complete necropsies and microscopic examinations were performed on all F₁ rats and all mice at
30 the end of the 2-year studies. At necropsy, all organs and tissues were examined for grossly
31 visible lesions and all major tissues were fixed and preserved in 10% neutral buffered formalin
32 (except eyes, which were first fixed in Davidson's solution, and testes, vaginal tunics, and
33 epididymides, which were first fixed in modified Davidson's solution). Tissues were processed
34 and trimmed, embedded in paraffin, sectioned at a thickness of 4 to 6 µm, and stained with H&E
35 for microscopic examination. For all paired organs (e.g., adrenal gland, kidney, ovary), samples
36 from each organ were examined. In the original evaluation of the uterus, a transverse section
37 through each uterine horn, approximately 0.5 cm cranial to cervix, was collected for
38 histopathology evaluation. For the residual tissue evaluation of the uterus, all remaining uterine
39 tissue, including the cervix and vaginal tissue, was sectioned longitudinally, processed, and
40 examined histologically. Results from the residual uterine evaluation were combined with those
41 from the original, transverse section of uterus. Tissues examined microscopically are listed in
42 Table 1.

1 Microscopic evaluations were completed by the study laboratory pathologist, and the pathology
2 data were entered into the Toxicology Data Management System. The report, slides, paraffin
3 blocks, residual wet tissues, and pathology data were sent to the NTP Archives for inventory,
4 slide/block match, wet tissue audit, and storage. The slides, individual animal data records, and
5 pathology tables were evaluated by an independent QA laboratory. The individual animal
6 records and tables were compared for accuracy, the slide and tissue counts were verified, and the
7 histotechnique was evaluated. For the 2-year studies, a QA pathologist evaluated slides from all
8 neoplasms and all potential target organs, which included the kidney of rats and mice; the liver
9 of male rats and male mice; the uterus of female rats and female mice; the cecum of mice; the
10 adrenal cortex and mandibular lymph node of male rats; the testis and epididymis of male mice;
11 the ovary and nose of female rats; and the spleen and mesenteric lymph node of female mice.

12 The QA report and the reviewed slides were submitted to the NTP pathologist, who reviewed
13 and addressed any inconsistencies in the diagnoses made by the laboratory and QA pathologist.
14 The QA pathologist, who served as the coordinator of the Pathology Working Group (PWG)
15 presented representative histopathology slides containing examples of lesions related to test
16 agent administration, examples of disagreements in diagnoses between the laboratory and QA
17 pathologist, or lesions of general interest to the PWG for review. The PWG consisted of the NTP
18 pathologist and other pathologists experienced in rodent toxicological pathology. When the PWG
19 consensus differed from the opinion of the laboratory pathologist, the diagnosis was changed.
20 Final diagnoses for reviewed lesions represent a consensus between the laboratory pathologist,
21 QA pathologist, and the PWG. Details of these review procedures have been described, in part,
22 by Maronpot and Boorman³⁵ and Boorman et al.³⁷ or subsequent analyses of the pathology data;
23 the decision of whether to evaluate the diagnosed lesions for each tissue type separately or
24 combined was generally based on the guidelines of Brix et al.³⁸

25 In addition to the routine pathology review, a step section analysis of the kidneys from the
26 chronic mouse study was performed. Paraffin-embedded kidneys were sectioned at 1 mm
27 intervals to obtain three to four additional sections per kidney to allow for the observation of
28 additional renal neoplasms. The evaluation of these additional kidney slides was conducted by a
29 board-certified pathologist other than the QA pathologist. The step section pathologist shared the
30 evaluation findings with the NTP pathologist and the QA pathologist. Representative slides from
31 the kidney step section review were taken to the PWG for the chronic ST mouse study and were
32 examined by the members of that PWG. The final diagnosis of the kidney step section review
33 constitutes a consensus of the kidney step section pathologist, the NTP pathologist, and the PWG
34 participants.

1 **Table 1. Experimental Design and Materials and Methods in the Three-month and Two-year**
 2 **Drinking Water Studies of Sodium Tungstate Dihydrate**

Three-month Studies	Two-year Studies
Study Laboratory	
Rats: Battelle (Columbus, OH)	Same as 3-month studies
Mice: Battelle (Columbus, OH)	
Strain and Species	
Rats: Sprague Dawley (Hsd:Sprague Dawley® SD®)	Same as 3-month studies
Mice: B6C3F1/N	
Animal Source	
Rats: Harlan Laboratories, Inc. (now Envigo; Indianapolis, IN)	Same as 3-month studies
Mice: Taconic Biosciences, Inc. (Germantown, NY)	
Time Held Before Studies	
F ₀ female rats: 5 days	F ₀ female rats: 4 days
Mice: 11 days	Mice: 11 days
Average Age When Studies Began	
F ₀ female rats: 11 to 12 weeks	F ₀ female rats: 11 to 14 weeks
Mice: 5 to 6 weeks	Mice: 5 to 6 weeks
Date of First Exposure	
F ₀ female rats: May 23, 2009	F ₀ female rats: December 23, 2011
F ₁ rats: June 30 (males) or July 1 (females), 2009	F ₁ rats: January 30 (males) or 31 (females), 2011
Mice: June 1 (females) or 2 (males), 2009	Mice: January 16 (females) or 17 (males), 2012
Duration of Exposure	
F ₀ female rats: GD 6 to LD 21	F ₀ female rats: GD 6 to LD 21
F ₁ rats: 3 months	F ₁ rats (interim evaluations): 3, 6, 12, and 18 months F ₁ rats (2-year study): 105 weeks
Mice: 3 months	Mice (interim evaluations): 3, 6, 12, and 18 months Mice (2-year study): 105 weeks
Date of Last Exposure	
F ₀ female rats: June 30, 2009	F ₀ female rats: January 30, 2012
F ₁ rats: September 28 (males) or 29 (females), 2009	F ₁ rats (3-month interim): May 1 (males) or 2 (females), 2012 F ₁ rats (6-month interim): August 1 (males) or 2 (females), 2012 F ₁ rats (12-month interim): January 30 (males) or 31 (females), 2013 F ₁ rats (18-month interim): July 31 (males) or August 1 (females), 2013 F ₁ rats (2-year study): January 28 (males) or 31 (females), 2014

Three-month Studies	Two-year Studies
Mice: August 31 (females) or September 1 (males), 2009	Mice (3-month interim): April 17 (females) or 18 (males), 2012 Mice (6-month interim): July 17 (females) or 18 (males), 2012 Mice (12-month interim): January 15 (females) or 16 (males), 2013 Mice (18-month interim): July 16 (females) or 17 (males), 2013 Mice (2-year study): January 15 (females) or 17 (males), 2014
Necropsy Dates	
F ₁ rats: September 28 (males) or 29 (females), 2009	F ₁ rats (2-year study): January 27 and 28 (males) or 29 to 31 (females), 2014
Mice: August 31 (females) or September 1 (males), 2009	Mice (2-year study): January 13 to 15 (females) or 15 to 17 (males), 2014
Average Age at Necropsy	
F ₁ rats: 15 to 16 weeks	F ₁ rats (2-year study): 108 weeks
Mice: 18 to 19 weeks	Mice (2-year study): 109 to 110 weeks
Size of Study Groups	
F ₀ female rats: 8 F ₁ rats: 10/sex	F ₀ female rats: 47 (0 mg/L) or 41 (250, 500, and 1,000 mg/L) F ₁ rats (interim evaluations): 40/sex F ₁ rats (2-year study): 50/sex
Mice: 10/sex	Mice (interim evaluations): 40/sex Mice (2-year study): 50/sex
Method of Distribution	
Rats: Dams were distributed randomly into groups of approximately equal initial mean body weights. Pups were standardized on each litter's respective PND 4 to a maximum of eight pups per litter. Weaned pups were randomized on PND 20.	Rats: Dams were distributed randomly into groups of approximately equal initial mean body weights. Pups were standardized on each litter's respective PND 4 to a maximum of eight pups per litter. Weaned pups were randomized on PND 19.
Mice: Animals were distributed randomly into groups of approximately equal initial mean body weights.	Mice: Same as 3-month study
Animals per Cage	
F ₀ female rats: 1 (with litter)	F ₀ female rats: 1 (with litter)
F ₁ rats: 5 (males) or 5 (females)	F ₁ rats: up to 2 (males) or up to 5 (females)
Mice: 1 (male) or 5 (females)	Mice: 1 (male) or up to 4 (females)
Method of Animal Identification	
F ₀ female rats: Cage card and tail marking with permanent pen	Same as 3-month studies
F ₁ rats: Cage card and tail tattoo	
Mice: Cage card and tail tattoo	

Three-month Studies	Two-year Studies
Diet	
Irradiated NIH-07 wafer feed (rats; perinatal phase) or irradiated NTP-2000 wafer feed (rats and mice; 3-month studies) (Zeigler Brothers Inc., Gardners, PA), available ad libitum, changed weekly	Same as 3-month studies
Water	
Tap water (Columbus municipal supply), deionized, either untreated or containing a formulation of ST via glass bottles (Wheaton Science Products, Millville, NJ [rats and female mice] or Supelco, Bellefonte, PA [male mice]), available ad libitum, changed twice weekly	Tap water (Columbus municipal supply), deionized, either untreated or containing a formulation of ST via glass bottles (Fisher Scientific, Pittsburgh, PA [rats], Qorpak, Bridgeville, PA [female mice], or VWR, West Chester, PA [male mice]), available ad libitum, changed twice weekly
Cages	
Solid polycarbonate (Lab Products, Inc., Seaford, DE) Rats: changed weekly through PND 4, then twice weekly, rotated every 2 weeks Mice: Changed weekly (males) or twice weekly (females), rotated every 2 weeks	Same as 3-month studies
Bedding	
Irradiated Sani-Chips® (P.J. Murphy Forest Products Corporation, Montville, NJ), changed with cage changes	Same as 3-month studies
Rack Filters	
Spun-bonded polyester (Snow Filtration Company, Cincinnati, OH), changed every 2 weeks	Same as 3-month studies
Racks	
Stainless steel (Lab Products, Inc., Seaford, DE), changed and rotated every 2 weeks	Same as 3-month studies
Animal Room Environment	
Temperature: 72°F ± 3°F Relative humidity: 50% ± 15% Room fluorescent light: 12 hours/day Room air changes: 10/hour	Same as 3-month studies
Exposure Concentrations	
0, 125, 250, 500, 1,000, or 2,000 mg/L	Rats: 0, 250, 500, or 1,000 mg/L Mice: 0, 500, 1,000, or 2,000 mg/L

Three-month Studies	Two-year Studies
Type and Frequency of Observation	
<p>F₀ female rats: Observed twice daily. Weighed on GDs 5, 6, 9, 12, 15, 18, and 21 and on LDs 1, 4, 7, 14, and 21. Water consumption was measured over 3-day intervals from GD 6 through LD 21.</p>	<p>F₀ female rats: Observed twice daily. Weighed on GDs 2, 5, 6, 9, 12, 15, 18, and 21 and on LDs 1, 4, 7, 10, 14, 17, and 21. Water consumption was measured over 3-day intervals from GD 6 through LD 21.</p>
<p>F₁ rats: Observed twice daily. Litter data (total litter weight, litter weights by sex, and litter observations) were recorded on PND 1. Pup survival was evaluated and recorded. Individual pups were weighed on PNDs 4, 7, 14, and 21, weekly for 3 months, and at the end of the study. Clinical observations were recorded weekly and at the end of the study. Water consumption was recorded weekly for the duration of the study.</p>	<p>F₁ rats: Observed twice daily. Litter data (total litter weight, litter weights by sex, and litter observations) were recorded on PND 1. Pups per litter were recorded on PNDs 2 and 3. Pups were weighed on PNDs 4, 7, 10, 14, and 21, weekly for 3 months, then every 4 weeks, and at the end of the study. Clinical observations were recorded every 4 weeks beginning at week 6 and at the end of the study. Water consumption was recorded initially, weekly for 3 months, and then at 4-week intervals thereafter.</p>
<p>Mice: Observed twice daily. Weighed initially, weekly for 3 months, and at the end of the study. Clinical observations were recorded weekly and at the end of the study. Water consumption was recorded weekly for the duration of the study.</p>	<p>Mice: Observed twice daily. Weighed initially, weekly for 3 months, then every 4 weeks, and at the end of the study. Clinical observations were recorded at week 6 then every 4 weeks and at the end of the study. Water consumption was measured initially, weekly for 3 months, and then at 4-week intervals thereafter.</p>
Method of Euthanasia	
Carbon dioxide	Same as 3-month studies
Necropsy	
<p>Necropsies were performed on all animals. Organs weighed at the end of the study were: liver, thymus, right kidney, right testis, heart, and lungs.</p>	<p>Necropsies were performed on all core animals. Organs collected and weighed at the 3-, 6-, 12-, and 18-month interim evaluations were: liver, left and right kidneys, stomach, small intestine, and bone (femur).</p>
Clinical Pathology	
<p>At the end of the studies, blood was collected from the retroorbital plexus (rats) or sinus (mice) for clinical chemistry (rats only), hematology, and insulin determination (rats only).</p>	None
<p><i>Hematology</i>: erythrocyte count, mean corpuscular volume, hemoglobin, hematocrit, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration, leukocyte count and differentials, reticulocyte count, and platelet count</p>	
<p><i>Clinical chemistry (rats)</i>: alanine aminotransferase, albumin, alkaline phosphatase, bile acids, cholesterol, creatinine, creatine kinase, glucose, sorbitol dehydrogenase, total protein, triglycerides, and urea nitrogen</p>	

Three-month Studies	Two-year Studies
<p>Histopathology</p> <p>Complete histopathology was performed on all F₁ rats and all mice in the vehicle control and 2,000 mg/L groups. The kidney was identified as a target organ and examined to a no-effect level. In addition to gross lesions and tissue masses, the following tissues were examined: adrenal gland, bone with marrow, brain, clitoral gland, esophagus, eyes, gallbladder (mice), Harderian gland, heart, kidney, large intestine (cecum, colon, rectum), small intestine (duodenum, jejunum, ileum), larynx, liver, lung, lymph nodes (mandibular and mesenteric), mammary gland, nose, ovary, pancreas, parathyroid gland, pharynx, pituitary gland, preputial gland, prostate gland, salivary gland, seminal vesicle, skin, spinal cord, spleen, sternum, stomach (forestomach and glandular), testis with epididymis, thymus, thyroid gland, tongue, trachea, urinary bladder, uterus, vagina, and Zymbal's gland.</p>	<p>Complete histopathology was performed on all core F₁ rats and all core mice. In addition to gross lesions and tissue masses, the following tissues were examined: adrenal gland, bone with marrow, brain, clitoral gland, esophagus, eyes, gallbladder (mice), Harderian gland, heart, kidney, large intestine (cecum, colon, and rectum), small intestine (duodenum, jejunum, and ileum), liver, lung, lymph nodes (mandibular and mesenteric), mammary gland, nose, ovary, pancreas, parathyroid gland, pituitary gland, preputial gland, prostate gland, salivary gland, seminal vesicle, spleen, stomach (forestomach and glandular), testis with epididymis, thymus, thyroid gland, trachea, urinary bladder, uterus (including cervix), and vagina.</p>
<p>Sperm Motility</p> <p>At the end of the studies, sperm samples were collected from F₁ male rats and male mice in the vehicle control, 500, 1,000, and 2,000 mg/L groups for sperm evaluations. The following parameters were evaluated: spermatid heads per testis and per gram testis, and epididymal spermatozoal motility and concentration. The left cauda, left epididymis, and left testis were weighed.</p>	<p>None</p>
<p>Urinalysis</p> <p>During week 12, urine samples were collected from F₁ rats in metabolism cages for urinalysis.</p> <p><i>Parameters Evaluated (rats):</i> specific gravity, volume, sediment exam, protein, glucose, creatinine, calcium, phosphorous, N-acetyl-β-glucosaminidase, alkaline phosphatase, and aspartate aminotransferase</p>	<p>At the 3-, 6-, 12-, and 18-month interim evaluations, urine was collected from up to 10 predesignated rats/sex/exposure group in metabolism cages for urinalysis.</p> <p><i>Parameters Evaluated (rats):</i> volume and creatinine</p>
<p>Xanthine and Methionine Analysis</p> <p>At the end of the studies, urine and blood samples were collected from all F₁ rats for xanthine/methionine determination.</p>	<p>None</p>

Three-month Studies	Two-year Studies
Internal Dose Assessment	
At the end of the studies, urine (F ₁ rats) and blood (F ₁ rats, mice) samples were collected for tungsten determinations.	At the 3-, 6-, 12-, and 18-month interim evaluations, urine, plasma, and kidney were collected from up to 10 predesignated animals/sex/exposure group for tungsten determination. Tungsten concentrations were determined using validated analytical methods (Appendix E). On PND 21, tungsten concentrations were determined in plasma from dams (five/exposure group) and their pups (one male and one female from each selected dam's litter).

1 GD = gestation day; LD = lactation day; PND = postnatal day.

2 **Statistical Methods**

3 **Survival Analyses**

4 The probability of survival was estimated by the product-limit procedure of Kaplan and Meier³⁹
 5 and is presented graphically. Animals surviving to the end of the observation period are treated
 6 as censored observations, as are animals dying from unnatural causes within the observation
 7 period. Animals dying from natural causes are included in analyses and are treated as uncensored
 8 observations. For the 2-year mouse study, exposure concentration-related trends are identified
 9 with Tarone's life-table test,⁴⁰ and pairwise exposure concentration-related effects are assessed
 10 using Cox's method.⁴¹ For the rat perinatal study, exposure concentration-related trends and
 11 pairwise exposure concentration-related effects on survival are assessed using a Cox proportional
 12 hazards model⁴¹ with a random litter effect. All reported p values for the survival analyses are
 13 two-sided.

14 **Calculation of Incidence**

15 The incidences of neoplasms or nonneoplastic lesions are presented as the numbers of animals
 16 bearing such lesions at a specific anatomical site. For calculation of incidence rates, the
 17 denominator for most neoplasms and all nonneoplastic lesions is the number of animals where
 18 the site was examined microscopically. However, when macroscopic examination was required
 19 to detect neoplasms in certain tissues (e.g., mesentery, pleura, peripheral nerve, skeletal muscle,
 20 tongue, tooth, and Zymbal's gland) before microscopic evaluation, the denominator consists of
 21 the number of animals that had a gross abnormality. When neoplasms had multiple potential sites
 22 of occurrence (e.g., leukemia or lymphoma), the denominator consists of the number of animals
 23 on which a necropsy was performed. Additional study data also give the survival-adjusted
 24 neoplasm rate for each group and each site-specific neoplasm. This survival-adjusted rate (based
 25 on the Poly-3 method described below) accounts for differential mortality by assigning a reduced
 26 risk of neoplasm, proportional to the third power of the fraction of time on study, only to site-
 27 specific, lesion-free animals that do not reach terminal euthanasia.

28 **Analysis of Neoplasm and Nonneoplastic Lesion Incidence**

29 Statistical analyses of neoplasm and nonneoplastic lesion incidence considered two features of
 30 the data. Some animals did not survive the entire 2 years of the study, so survival differences
 31 between groups had to be considered. Also, for the rat study, up to two animals per sex were
 32 randomly selected from each rat litter to participate in the study. The statistical analysis of lesion

1 incidence used the Poly-3 test to account for survival differences, with a Rao-Scott adjustment
2 for litter effects as needed. This analysis is described below.

3 The Poly-k test⁴²⁻⁴⁴ was used to assess neoplasm and nonneoplastic lesion prevalence. This test is
4 a survival-adjusted quantal-response procedure that modifies the Cochran-Armitage linear trend
5 test to account for survival differences. More specifically, this method modifies the denominator
6 in the quantal estimate of lesion incidence to approximate more closely the total number of
7 animal years at risk. For analysis of a given site, each animal is assigned a risk weight. This
8 value is 1 if the animal had a lesion at that site or if it survived until terminal euthanasia; if the
9 animal died before terminal euthanasia and did not have a lesion at that site, its risk weight is the
10 fraction of the entire study time that it survived, raised to the kth power.

11 This method yields a lesion prevalence rate that depends only on the choice of a shape parameter
12 for a Weibull hazard function describing cumulative lesion incidence over time.⁴² Unless
13 otherwise specified, a value of $k = 3$ was used in the analysis of site-specific lesions. This value
14 was recommended by Bailer and Portier⁴² following an evaluation of neoplasm onset time
15 distributions for a variety of site-specific neoplasms in control Fischer 344 rats and
16 B6C3F1 mice.⁴⁵ Bailer and Portier⁴² showed that the Poly-3 test gave valid results if the true
17 value of k was anywhere in the range from 1 to 5. A further advantage of the Poly-3 method is
18 that it does not require lesion lethality assumptions. Variation introduced by the use of risk
19 weights, which reflect differential mortality, was accommodated by adjusting the variance of the
20 Poly-3 statistic as recommended by Bieler and Williams.⁴⁶ Poly-3 tests used the continuity
21 correction described by Nam.⁴⁷

22 Littermates tend to be more like each other than fetuses/pups in other litters. Failure to account
23 for correlation within litters leads to underestimates of variance in statistical tests, resulting in
24 higher probabilities of Type I errors (“false positives”). Because up to two pups/sex/litter were
25 present in the core rat study, the Poly-3 test was modified to accommodate litter effects using the
26 Rao-Scott approach.⁴⁸ The Rao-Scott approach accounts for litter effects by estimating the ratio
27 of the variance in the presence of litter effects to the variance in the absence of litter effects. This
28 ratio is then used to adjust the sample size downward to yield the estimated variance in the
29 presence of litter effects. The Rao-Scott approach was implemented in the Poly-3 test as
30 recommended by Fung et al.,⁴⁹ formula \bar{T}_{RS2} .

31 Tests of significance included pairwise comparisons of each exposure group with control groups
32 and a test for an overall exposure concentration-related trend. Continuity-corrected Rao-Scott-
33 adjusted Poly-3 tests were used in the analysis of lesion incidence and reported p values are one-
34 sided. The significance of a lower incidence or negative trend in lesions is approximated as $1-p$
35 with the letter N added (e.g., $p = 0.99$ is presented as $p = 0.01N$). For neoplasms and
36 nonneoplastic lesions observed without litter structure (e.g., for the mouse studies), Poly-3 tests
37 that included the continuity correction, but without adjustment for potential litter effects, were
38 used for trend and pairwise comparisons to the control group.

39 To evaluate incidence rates by litter, the proportions of litters affected by each lesion type were
40 tested among groups. Cochran-Armitage trend tests and Fisher’s exact tests⁵⁰ were used to test
41 for trends and pairwise differences from the control group, respectively.

1 **Analysis of Continuous Variables**

2 Before statistical analysis, extreme values identified by the outlier test of Dixon and Massey⁵¹ for
3 small samples ($n < 20$) and Tukey's outer fences method⁵² for large samples ($n \geq 20$) were
4 examined by NTP personnel, and implausible values were eliminated from the analysis.

5 For the perinatal and 2-year study in rats and the 3-month and 2-year studies in mice, litter
6 effects were not considered in the analysis of the continuous data. Organ and body weight
7 measurements, which historically have approximately normal distributions, were analyzed with
8 the parametric multiple comparison procedures of Dunnett⁵³ and Williams.^{54; 55} Dam gestational
9 and lactational feed consumption, litter sizes, pup survival, implantations, number of resorptions,
10 and proportions of male pups per litter for all rat studies were analyzed using the nonparametric
11 multiple comparison methods of Shirley⁵⁶ (as modified by Williams⁵⁷ and Dunn⁵⁸) given that
12 these endpoints typically have skewed distributions. For all quantitative endpoints unaffected by
13 litter structure, the Jonckheere test⁵⁹ was used to assess the significance of the exposure
14 concentration-related trends and to determine, at the 0.01 level of significance, whether a trend-
15 sensitive test (the Williams or Shirley test) was more appropriate for pairwise comparisons than a
16 test that does not assume a monotonic exposure concentration-related trend (the Dunnett or Dunn
17 test).

18 For the perinatal and 3-month study in rats, there were two or more littermates in each exposure
19 group analyzed. Consequently, organ and body weight endpoints were analyzed using linear
20 mixed models, with litter as a random effect. To adjust for multiple comparisons, a Dunnett-Hsu
21 adjustment was used.⁶⁰ For other endpoints, for which normality was not assumed, the trend
22 across exposure groups was analyzed by a permutation test based on the Jonckheere trend test
23 implemented by randomly permuting whole litters across exposure groups and bootstrapping
24 within the litters (see, for example, Davison and Hinkley⁶¹). Pairwise comparisons were made
25 by using a modified Wilcoxon test that incorporated litter effects.⁶² The Hommel procedure was
26 used to adjust for multiple comparisons.⁶³

27 Postweaning body weights were measured on two pups/sex/litter in the perinatal and 2-year and
28 perinatal and 3-month rat studies; more than two pups/sex/litter were possible in preweaning
29 body weight measurements. The analyses of pup body weights and body weights adjusted for
30 litter size (described below) took litter effects into account using a mixed model with litter as
31 random effect. To adjust for multiple comparisons, a Dunnett-Hsu adjustment was used.⁶⁰ Dam
32 body weights during gestation and lactation were analyzed with the parametric multiple
33 comparison procedures of Dunnett⁵³ or Williams,^{54; 55} depending on whether the Jonckheere test
34 indicated the use of a trend-sensitive test. P values for these analyses are two-sided.

35 **Analysis of Gestational and Fertility Indices**

36 Cochran-Armitage trend tests were used to test the significance of trends in gestational and
37 fertility data across exposure groups. Fisher's exact test was used to conduct pairwise
38 comparisons of each exposure group with the control group. P values for these analyses are
39 two-sided.

1 **Body Weight Adjustments**

2 To adjust preweaning pup body weights for live litter size, a linear model was fit to body weights
3 as a function of exposure concentration and litter size. The estimated coefficient of litter size was
4 then used to adjust each pup body weight based on the difference between its litter size and the
5 mean litter size. Prestandardization PND 4 body weights were adjusted for PND 1 litter size, and
6 body weights measured between PND 4 poststandardization and PND 21 were adjusted for
7 PND 4 poststandardization litter size. After adjustment, body weights were analyzed with a
8 linear mixed model with a random litter effect.

9 **Historical Control Data**

10 The concurrent control group is the most valid comparison to the exposed groups and is the only
11 control group analyzed statistically in NTP bioassays. Historical control data are often helpful in
12 interpreting potential exposure-related effects, however, particularly for uncommon or rare
13 neoplasm types. For meaningful comparisons, the conditions for studies in the historical control
14 data must be generally similar. Significant factors affecting the background incidence of
15 neoplasms at a variety of sites are diet, sex, strain/stock, and route of exposure. The NTP
16 historical control database contains all 2-year studies for each species, sex, and strain/stock with
17 histopathological findings in control animals completed within the most recent 5-year period⁶⁴⁻⁶⁶
18 including the concurrent control for comparison across multiple technical reports. In general, the
19 historical control data for a given study includes studies using the same route of administration,
20 and the overall incidence of neoplasms in control groups for all routes of administration are
21 included for comparison, including the current study.

22 **Quality Assurance Methods**

23 The 3-month and 2-year studies were conducted in compliance with U.S. Food and Drug
24 Administration Good Laboratory Practice Regulations.⁶⁷ In addition, the 3-month and 2-year
25 study reports were audited retrospectively by an independent QA contractor against study
26 records submitted to the NTP Archives. Separate audits covered completeness and accuracy of
27 the pathology data, pathology specimens, final pathology tables, and a draft of this NTP
28 Technical Report. Audit procedures and findings are presented in the reports and are on file at
29 NIEHS. The audit findings were reviewed and assessed by NTP staff, and all comments were
30 resolved or otherwise addressed during the preparation of this Technical Report.

31 **Genetic Toxicology**

32 The genetic toxicity of ST was assessed by testing the ability of the chemical to induce mutations
33 in various strains of *Salmonella typhimurium* and *Escherichia coli*, to increase the frequency of
34 micronucleated erythrocytes in rat and mouse peripheral blood, and to increase DNA damage in
35 cells from liver, kidney, ileum, and peripheral blood from rats and mice. The protocols and
36 results for these studies are given in Appendix D.

37 The genetic toxicity studies are an outcome of an earlier effort by NTP to develop a
38 comprehensive database permitting a critical anticipation of a chemical's carcinogenicity in
39 experimental animals based on the results from several in vitro and in vivo short-term tests
40 measuring functionally distinct genotoxicity endpoints. The short-term tests were originally
41 developed to clarify proposed mechanisms of chemical-induced DNA damage on the basis of the

1 relationship between electrophilicity and mutagenicity⁶⁸ and the somatic mutation theory of
2 cancer.^{69; 70} It should be noted, however, that not all cancers arise through genotoxic
3 mechanisms.

4 **Bacterial Mutagenicity**

5 DNA reactivity combined with *Salmonella* mutagenicity is highly correlated with induction of
6 carcinogenicity in multiple species/sexes of rodents and at multiple tissue sites.⁷¹ A positive
7 response in the *Salmonella* test was shown to be the most predictive in vitro indicator for rodent
8 carcinogenicity (89% of the *Salmonella* mutagens are rodent carcinogens).^{72; 73} Additionally, no
9 battery of tests that included the *Salmonella* test improved the predictivity of the *Salmonella* test
10 alone; however, these other tests can provide useful information on the types of DNA and
11 chromosomal damage induced by the chemical under investigation. The protocol for these
12 studies and the results are given in Appendix D.

13 **Peripheral Blood Micronucleus Test**

14 Micronuclei are biomarkers of induced structural or numerical chromosomal alterations and are
15 formed when acentric fragments or whole chromosomes fail to incorporate into either of two
16 daughter nuclei during cell division.^{74; 75} The predictivity for carcinogenicity of a positive
17 response in acute in vivo bone marrow chromosome aberration or micronucleus tests appears to
18 be less than that of the *Salmonella* test.^{76; 77} However, clearly positive results in long-term
19 peripheral blood micronucleus tests have high predictivity for rodent carcinogenicity; a weak
20 response in one sex only or negative results in both sexes in this assay do not correlate well with
21 either negative or positive results in rodent carcinogenicity studies.⁷⁸ Because of the theoretical
22 and observed associations between induced genetic damage and adverse effects in somatic and
23 germ cells, the determination of in vivo genetic effects is important to the overall understanding
24 of the risks associated with exposure to a particular chemical. The protocol for these studies and
25 the results are given in Appendix D.

26 **Comet Assay**

27 The alkaline (pH > 13) comet assay⁷⁹ (also known as the single cell gel electrophoresis assay)
28 detects DNA damage in any of a variety of eukaryotic cell types⁸⁰⁻⁸³; cell division is not required.
29 The type of DNA damage detected includes nicks, adducts, strand breaks, and abasic sites that
30 are converted to DNA strand breaks after treatment of cells in an alkaline (pH > 13) solution.
31 Transient DNA strand breaks generated by the process of DNA excision repair might also be
32 detected. DNA damage caused by crosslinking agents has been detected as a reduction of DNA
33 migration.^{84; 85} The fate of the DNA damage detected by the comet assay is varied; most of the
34 damage is rapidly repaired and results in no sustained effect on the tissue, but some might result in
35 cell death or be incorrectly processed by repair proteins and lead to a fixed mutation or
36 chromosomal alteration. The protocol for these studies and the results are given in Appendix D.

37 In the rat study, ileum (male rats) and kidney (male and female rats) samples were tested in the
38 comet assay, but due to inconsistencies of the results, the assay was deemed an invalid test for
39 these samples.

1 Results

2 Data Availability

3 The National Toxicology Program (NTP) evaluated all study data. Data relevant for evaluating
4 toxicological findings are presented here. All study data are available in the NTP Chemical
5 Effects in Biological Systems (CEBS) database: <https://doi.org/10.22427/NTP-DATA-TR-599>.⁸⁶

6 Rats

7 Three-month Study (Perinatal Phase)

8 No significant effects related to sodium tungstate dihydrate (ST) exposure were observed on
9 pregnancy status, maternal survival, or the number of dams that littered (Table 2). There were no
10 clinical observations in dams (Appendix G). One dam in the 250 mg/L group was euthanized on
11 gestation day (GD) 25 due to moribundity associated with incomplete labor, and one 2,000 mg/L
12 dam (and her pups) was euthanized moribund on lactation day (LD) 6.

13 No significant effects on dam mean body weight during gestation were observed, but mean body
14 weights were significantly decreased in the 1,000 and 2,000 mg/L groups starting at LD 14. By
15 the end of lactation (LD 21), the 1,000 and 2,000 mg/L dam groups showed significant decreases
16 in group mean body weight of approximately 10% and 18%, respectively, when compared to the
17 vehicle control group (Table 3).

18 **Table 2. Summary of the Disposition of F₀ Female Rats during Perinatal Exposure in the Perinatal**
19 **and Three-month Drinking Water Study of Sodium Tungstate Dihydrate**

	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Reproductive Performance						
Time-mated Females (GD 6)	8	8	8	8	8	8
Females Pregnant (%) ^a	7 (87.5)	8 (100.0)	8 (100.0)	6 (75.0)	6 (75.0)	6 (75.0)
Females Not Pregnant (%)	1 (12.5)	0	0	2 (25.0)	2 (25.0)	2 (25.0)
Dams Not Delivering with Evidence of Pregnancy (%)	0	0	1 (12.5) ^b	0	0	0
Dams with Litters on LD 0 (%)	7 (100)	8 (100.0)	7 (100.0)	6 (100.0)	6 (100.0)	6 (100.0)
Litters Poststandardization (LD 4) ^c	5	7	7	6	5	5

20 GD = gestation day; LD = lactation day.

21 ^aStatistical analysis performed by the Cochran-Armitage (trend) and Fisher's exact (pairwise) tests.

22 ^bDam died before littering.

23 ^cStandardization to eight pups/litter (four pups/sex). Only litters with at least two pups/sex and at least eight pups total/litter were
24 retained to continue on study.

1 **Table 3. Summary of Mean Body Weights and Body Weight Gains of F₀ Female Rats during**
 2 **Gestation and Lactation in the Perinatal and Three-month Drinking Water Study of Sodium**
 3 **Tungstate Dihydrate**

Parameter ^{a,b}	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Gestation Day						
6	229.8 ± 3.1 (7)	220.5 ± 8.8 (8)	224.5 ± 6.2 (8)	228.1 ± 5.7 (6)	225.5 ± 4.6 (6)	228.9 ± 4.6 (6)
9	243.3 ± 2.7 (7)	241.4 ± 5.3 (8)	245.6 ± 4.3 (8)	239.1 ± 5.4 (6)	237.2 ± 5.1 (6)	242.1 ± 5.8 (6)
12	259.1 ± 3.6 (7)	257.4 ± 4.9 (8)	261.4 ± 4.4 (8)	255.5 ± 5.8 (6)	254.5 ± 4.8 (6)	258.3 ± 6.6 (6)
15	275.5 ± 6.0 (7)	279.9 ± 4.9 (8)	281.5 ± 5.1 (8)	276.2 ± 5.0 (6)	269.3 ± 4.3 (6)	277.9 ± 8.6 (6)
18	304.9 ± 10.9 (7)	317.6 ± 5.6 (8)	313.4 ± 7.8 (8)	309.3 ± 5.8 (6)	301.2 ± 4.5 (6)	306.1 ± 16.0 (6)
21	345.0 ± 19.2 (7)	367.4 ± 7.2 (8)	353.9 ± 11.9 (8)	352.0 ± 6.0 (6)	338.7 ± 8.4 (6)	339.3 ± 26.0 (6)
Gestation Weight Change						
Gestation Day Interval						
6–9	13.5 ± 0.9 (7)	20.8 ± 4.3 (8)	21.1 ± 5.5 (8)	11.0 ± 0.7 (6)	11.7 ± 1.2 (6)	13.2 ± 1.6 (6)
9–12	15.8 ± 1.2 (7)	16.1 ± 0.9 (8)	15.8 ± 0.8 (8)	16.3 ± 1.0 (6)	17.3 ± 0.9 (6)	16.1 ± 1.5 (6)
12–15	16.4 ± 3.3 (7)	22.5 ± 1.3 (8)	20.0 ± 1.7 (8)	20.8 ± 1.2 (6)	14.8 ± 1.0 (6)	19.6 ± 2.9 (6)
15–18	29.3 ± 5.6 (7)	37.6 ± 1.4 (8)	32.0 ± 3.5 (8)	33.0 ± 1.5 (6)	31.8 ± 4.9 (6)	28.2 ± 8.7 (6)
18–21	40.1 ± 8.4 (7)	49.9 ± 3.0 (8)	40.5 ± 4.4 (8)	42.7 ± 1.8 (6)	37.5 ± 5.1 (6)	33.1 ± 11.5 (6)
6–21	115.2 ± 18.0 (7)	146.9 ± 6.4 (8)	129.4 ± 12.7 (8)	123.9 ± 2.2 (6)	113.3 ± 10.6 (6)	110.3 ± 22.9 (6)
Lactation Day						
1	275.0 ± 4.7 (6)	270.6 ± 3.7 (8)	277.2 ± 7.4 (7)	268.3 ± 4.7(6)	264.3 ± 5.3 (6)	257.2 ± 10.9 (6)
4	290.1 ± 5.4 (5)	287.2 ± 5.1 (7)	285.8 ± 7.2 (7)	279.9 ± 2.9(6)	271.1 ± 3.9 (5)	280.3 ± 8.2 (5)
7	300.2 ± 5.3* (5)	289.4 ± 4.5 (7)	299.4 ± 6.2 (7)	289.0 ± 2.7(6)	279.9 ± 5.1 (5)	281.9 ± 6.0 (4) ^c
14	325.4 ± 5.7** (5)	331.8 ± 6.7 (7)	326.1 ± 6.6 (7)	313.3 ± 6.3(6)	303.0 ± 3.7* (5)	288.6 ± 6.0** (4)
21	313.4 ± 5.5** (5)	315.2 ± 8.1 (7)	308.3 ± 8.3 (7)	306.4 ± 5.7(6)	283.4 ± 3.4* (5)	256.1 ± 8.7** (4)
Lactation Weight Change						
Lactation Day Interval						
1–4	12.7 ± 3.8 (5)	15.8 ± 3.2 (7)	8.5 ± 3.2 (7)	11.7 ± 2.1 (6)	11.3 ± 1.7 (5)	13.2 ± 3.4 (5)
4–7	10.1 ± 2.6 (5)	2.3 ± 2.0 (7)	13.6 ± 2.1 (7)	9.1 ± 2.4 (6)	8.8 ± 4.4 (5)	2.8 ± 6.0 (4) ^c
7–14	25.1 ± 2.7** (5)	42.4 ± 3.3 (7)	26.7 ± 4.3 (7)	24.3 ± 4.5 (6)	23.1 ± 4.8 (5)	6.7 ± 3.4** (4)
14–21	-12.0 ± 8.7 (5)	-16.6 ± 3.7 (7)	-17.8 ± 6.4 (7)	-7.0 ± 5.2 (6)	-19.6 ± 2.5 (5)	-32.5 ± 7.3 (4)
1–21	36.0 ± 8.6** (5)	43.8 ± 4.7 (7)	31.0 ± 9.7 (7)	38.1 ± 5.6 (6)	23.6 ± 4.5 (5)	-9.6 ± 6.0** (4)

4 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

5 Statistical significance for the vehicle control group indicates a significant trend test.

6 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

7 ^aEach exposure group was compared to the vehicle control group with the Williams test when a trend was present ($p \leq 0.01$ from the Jonckheere trend test) or with the Dunnett test when no trend was present.

8 ^bData presented as mean ± standard error (number of dams). Body weight data are presented in grams.

9 ^cOne dam and her pups in the 2,000 mg/L group were euthanized on PND 6 due to moribundity.

10

1 There were no significant changes in water consumption of exposed dams compared to vehicle
 2 control dams during gestation; however, during most of the lactation period, water consumption
 3 was significantly decreased for the 2,000 mg/L dams, with the largest difference (approximately
 4 27%) occurring during the end of lactation (LD 17–21) (Table 4). Water consumption by the
 5 500 mg/L group was up to 19% lower (LD 4–7) than that of the vehicle control group; however,
 6 consumption by the 1,000 mg/L group was no more than 8% lower (LD 17–21) than that of the
 7 vehicle control group (Table 4). Daily ST consumption for the 125, 250, 500, 1,000, and
 8 2,000 mg/L groups averaged approximately 17, 33, 58, 132, and 247 mg ST/kg body weight/day
 9 (mg/kg/day), respectively, during GD 6–21 and approximately 25, 56, 96, 220, and
 10 374 mg/kg/day, respectively, during LD 1–14 (Table 4).

11 **Table 4. Summary of Water and Sodium Tungstate Dihydrate Consumption of F₀ Female Rats**
 12 **during Gestation and Lactation in the Perinatal and Three-month Drinking Water Study**

Parameter ^a	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Gestation Day Interval^{b,c}						
6–9	28.9 ± 1.7 (7)	30.5 ± 1.2 (8)	30.9 ± 1.6 (8)	26.4 ± 2.0 (6)	28.1 ± 1.5 (6)	27.5 ± 1.7 (6)
9–12	30.7 ± 1.8 (7)	30.6 ± 1.7 (8)	30.7 ± 1.0 (8)	27.9 ± 1.7 (6)	36.3 ± 7.1 (6)	27.7 ± 1.2 (6)
12–15	33.1 ± 2.3 (7)	36.0 ± 1.2 (8)	36.2 ± 1.3 (8)	31.4 ± 0.8 (6)	33.5 ± 2.1 (6)	33.9 ± 1.6 (6)
15–18	40.6 ± 3.4 (7)	45.8 ± 2.9 (8)	40.4 ± 1.7 (8)	37.5 ± 1.2 (6)	39.7 ± 3.5 (6)	42.0 ± 1.8 (6)
18–21	41.4 ± 3.7* (7)	44.4 ± 2.9 (8)	42.1 ± 1.2 (8)	37.3 ± 1.3 (6)	39.6 ± 4.1 (6)	37.8 ± 3.0 (6)
6–21	34.9 ± 2.4* (7)	37.4 ± 1.5 (8)	36.1 ± 1.2 (8)	32.1 ± 1.2 (6)	35.5 ± 3.1 (6)	33.8 ± 1.0 (6)
Lactation Day Interval^{b,c}						
1–4	49.25 ± 2.66 (6)	51.10 ± 2.24 (8)	51.37 ± 2.67 (7)	44.20 ± 1.07 (6)	57.63 ± 7.59 (6)	45.00 ± 1.99 (6)
4–7	53.62 ± 2.43 (5)	45.77 ± 1.25 (7)	61.27 ± 5.07 (7)	43.50 ± 1.85* (6)	50.56 ± 2.06 (5)	45.08 ± 1.94 (4) ^d
7–10	68.78 ± 3.43** (5)	64.20 ± 1.51 (7)	71.41 ± 5.17 (7)	58.77 ± 2.59 (6)	64.94 ± 1.46 (5)	50.20 ± 1.25** (4)
10–14	85.46 ± 3.83** (5)	78.13 ± 2.49 (7)	84.93 ± 2.65 (7)	74.73 ± 1.33 (6)	78.56 ± 3.16 (5)	66.05 ± 2.66** (4)
14–17	90.48 ± 4.39* (5)	85.83 ± 4.03 (7)	89.60 ± 2.63 (7)	82.78 ± 2.94 (6)	87.48 ± 2.65 (5)	64.08 ± 0.54* (4)
17–21	108.6 ± 2.86** (5)	104.5 ± 2.59 (7)	107.3 ± 2.20 (6)	96.67 ± 3.50* (6)	99.96 ± 0.71* (5)	79.68 ± 2.02** (4)
1–14	66.42 ± 2.70** (5)	61.25 ± 1.70 (7)	68.61 ± 2.21 (7)	56.79 ± 1.14 (6)	62.48 ± 1.61 (5)	52.79 ± 1.31** (4)
Chemical Intake (mg/kg/day)^{e,f}						
GD 6–21	0.00 ± 0.00 (7)	16.80 ± 0.50 (8)	32.50 ± 1.11 (8)	58.33 ± 1.35 (6)	131.9 ± 11.20 (6)	246.6 ± 8.10 (6)
LD 1–14	0.00 ± 0.00 (5)	25.35 ± 0.69 (7)	56.46 ± 1.76 (7)	96.43 ± 1.61 (6)	219.5 ± 4.80 (5)	374.3 ± 6.22 (4)

13 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

14 Statistical significance for the vehicle control group indicates a significant trend test.

15 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

16 GD = gestation day; LD = lactation day.

17 ^aData presented as mean ± standard error (number of dams).

18 ^bWater consumption data are presented as grams/animal/day.

19 ^cEach exposure group was compared to the vehicle control group with the Shirley test when a trend was present ($p \leq 0.01$ from the Jonckheere trend test) or with the Dunn test when no trend was present.

20 ^dOne dam and her pups in the 2,000 mg/L group were euthanized on PND 6 due to moribundity.

21 ^eChemical intake calculated as: ([exposure concentration × water consumption]/[average body weight of day range]).

22 ^fNo statistical analysis was performed on the chemical intake data.

23

1 There were no exposure-related differences between the vehicle control group and the
2 ST-exposed groups in the number of litters, total litter weight, or litter size of males and females
3 on PND 1 or PND 4 (Table 5). The litter size did not change appreciably between PND 1 and
4 litter standardization on PND 4.

5 Two pups in the 125 mg/L group and one pup in the 1,000 mg/L group were found dead on
6 PND 1; one pup in the 1,000 mg/L group was missing on PND 2; and one pup in the 2,000 mg/L
7 group was found dead on PND 3. There was no pup mortality after PND 6. There were no
8 exposure-related clinical observations in dams or pups (Appendix G).

9 When adjusted for litter size, the mean body weight of male and female pups in the 2,000 mg/L
10 group on PND 21 was significantly decreased by approximately 16% and 11%, respectively,
11 compared to the corresponding vehicle control groups. For male and female pups in the 2,000
12 mg/L group combined, the mean body weight on PND 21 was significantly decreased by
13 approximately 14% relative to the vehicle control male and female pups combined (Table 6).

1 **Table 5. Summary of Mean Litter Size and Survival Ratio of F₁ Male and Female Rats during**
 2 **Lactation in the Perinatal and Three-month Drinking Water Study of Sodium Tungstate Dihydrate**

Parameter	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
PND 1^{a,b}						
Total	11.0 ± 1.2 (6)	13.0 ± 0.7 (8)	12.0 ± 0.7 (7)	12.0 ± 0.7 (6)	10.8 ± 2.1 (6)	12.2 ± 1.7 (6)
Live	11.0 ± 1.2 (6)	12.8 ± 0.7 (8)	12.0 ± 0.7 (7)	12.0 ± 0.7 (6)	10.7 ± 2.0 (6)	12.2 ± 1.7 (6)
% Male per Litter ^c	42.3 ± 8.2 (4)	44.3 ± 6.1 (5)	48.4 ± 4.9 (5)	53.7 ± 9.2 (4)	39.2 ± 10.7 (5)	54.7 ± 6.1 (5)
% Male ^{c,d,e}	39 (38)	44 (63)	49 (61)	52 (48)	48 (50)	54 (56)
Male^{a,b}						
PND 1 ^c	3.8 ± 0.3 (4)	5.6 ± 1.0 (5)	6.0 ± 0.9 (5)	6.3 ± 0.8 (4)	4.8 ± 1.3 (5)	6.0 ± 1.0 (5)
PND 4 Prestandardization ^c	2.8 ± 0.9 (4)	5.0 ± 1.4 (5)	6.0 ± 0.9 (5)	6.3 ± 0.8 (4)	4.8 ± 1.3 (5)	5.4 ± 1.5 (5)
PND 4 Poststandardization	3.8 ± 0.2* (5)	4.0 ± 0.0 (7)	4.0 ± 0.0 (7)	4.7 ± 0.4 (6)	4.0 ± 0.0 (5)	4.4 ± 0.4 (5)
Female^{a,b}						
PND 1 ^c	5.8 ± 1.3 (4)	7.0 ± 0.9 (5)	6.2 ± 0.5 (5)	5.8 ± 1.3 (4)	5.2 ± 1.2 (5)	5.2 ± 1.2 (5)
PND 4 Prestandardization ^c	5.3 ± 1.8 (4)	5.0 ± 1.3 (5)	6.2 ± 0.5 (5)	5.8 ± 1.3 (4)	5.0 ± 1.4 (5)	4.6 ± 1.5 (5)
PND 4 Poststandardization	4.2 ± 0.2* (5)	4.0 ± 0.0 (7)	4.0 ± 0.0 (7)	3.3 ± 0.4 (6)	4.0 ± 0.0 (5)	3.6 ± 0.4 (5)
Male and Female^{a,b}						
PND 4 Prestandardization	11.0 ± 1.2 (6)	12.8 ± 0.7 (8)	12.0 ± 0.7 (7)	12.0 ± 0.7 (6)	10.5 ± 1.9 (6)	12.0 ± 1.7 (6)
PND 4 Poststandardization	8.0 ± 0.0 (5)	8.0 ± 0.0 (7)	8.0 ± 0.0 (7)	8.0 ± 0.0 (6)	8.0 ± 0.0 (5)	8.0 ± 0.0 (5)
PND 21	8.0 ± 0.0 (5)	8.0 ± 0.0 (7)	8.0 ± 0.0 (7)	8.0 ± 0.0 (6)	8.0 ± 0.0 (5)	6.4 ± 1.6 (5)
Survival per Litter						
Total Dead: PND 1–4 ^{e,f}	0 (6)	2 (8) ^g	0 (7)	0 (6)	2 (6) ^g	1 (6)
Total Dead: PND 4–21 ^{e,f}	0 (5)	0 (7)	0 (7)	0 (6)	0 (5)	8 (5) ^h
Dead: PND 1–4 ^{a,b,i}	0.0 ± 0.0 (6)	0.3 ± 0.2 (8)	0.0 ± 0.0 (7)	0.0 ± 0.0 (6)	0.3 ± 0.3 (6)	0.2 ± 0.2 (6)
Dead: PND 4–21 ^{a,b,i}	0.0 ± 0.0 (5)	0.0 ± 0.0 (7)	0.0 ± 0.0 (7)	0.0 ± 0.0 (6)	0.0 ± 0.0 (5)	1.6 ± 1.6 (5)
Survival Ratio: PND 1–4 ^{a,b,j}	1.00 ± 0.00 (6)	1.00 ± 0.00 (8)	1.00 ± 0.00 (7)	1.00 ± 0.00 (6)	0.99 ± 0.01 (6)	0.99 ± 0.01 (6)
Survival Ratio: PND 4–21 ^{a,b,k}	1.00 ± 0.00 (5)	1.00 ± 0.00 (7)	1.00 ± 0.00 (7)	1.00 ± 0.00 (6)	1.00 ± 0.00 (5)	0.80 ± 0.20 (5)

3 Statistical significance for the vehicle control group indicates a significant trend test.

4 *Statistically significant at $p \leq 0.05$.

5 PND = postnatal day.

6 ^aEach exposure group was compared to the vehicle control group with the Shirley test when a trend was present ($p \leq 0.01$ from the Jonckheere trend test) or with the Dunn test when no trend was present.

7 ^bData presented as mean ± standard error (number of dams).

8 ^cLitters where the male/female pup counts were inconsistent between PND 1 and PND 4 were excluded from the male/female-specific endpoints.

9 ^d $100 \times [\text{number of live males in exposure group}]/[\text{number of live males and females in dietary exposure group}]$ (number of pups).

10 ^eNo statistics done on this endpoint.

11 ^fTotal dead in exposure group (number of dams).

12 ^gTwo pups in the 125 mg/L group and one pup in the 1,000 mg/L group were found dead on PND 1.

13 ^hOne dam and her pups in the 2,000 mg/L group were euthanized on PND 6 due to moribundity.

14 ⁱNumber dead/litter.

15 ^jSurvival per litter: number of pups prestandardization on PND 4/total live pups on PND 1.

16 ^kSurvival per litter: number of live pups on PND 21/number of live pups poststandardization on PND 4.

1 **Table 6. Summary of Preweaning F₁ Male and Female Rat Pup Mean Body Weights Following**
 2 **Perinatal Exposure to Sodium Tungstate Dihydrate**

Parameter	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Male (g)						
PND 1 ^{a,b,c}	7.8 ± 0.4** (4)	7.8 ± 0.2 (5)	7.8 ± 0.4 (5)	7.3 ± 0.3 (4)	7.2 ± 0.2 (4)	6.7 ± 0.3* (5)
PND 4 ^{d,e,f}	10.84 ± 0.46* (19/5)	10.49 ± 0.28 (28/7)	10.88 ± 0.52 (28/7)	9.90 ± 0.34 (28/6)	10.49 ± 0.39 (20/5)	9.34 ± 0.48 (22/5)
PND 7 ^{d,e,g}	16.94 ± 0.73* (19/5)	15.96 ± 0.41 (28/7)	16.67 ± 0.80 (28/7)	14.65 ± 0.43 (28/6)	15.61 ± 0.43 (20/5)	14.70 ± 0.81 (18/4)
PND 14 ^{d,e,g}	32.98 ± 0.87** (19/5)	32.16 ± 0.56 (28/7)	33.13 ± 1.15 (28/7)	30.40 ± 1.04 (28/6)	31.41 ± 0.73 (20/5)	29.22 ± 1.09 (18/4)
PND 21 ^{d,e,g}	55.56 ± 1.59** (19/5)	54.74 ± 0.58 (28/7)	55.05 ± 1.75 (28/7)	50.77 ± 1.92 (28/6)	51.89 ± 0.89 (20/5)	46.53 ± 1.44** (18/4)
Female (g)						
PND 1 ^{a,b,c}	7.1 ± 0.2 (4)	7.2 ± 0.1 (5)	7.6 ± 0.3 (5)	7.0 ± 0.3 (4)	7.1 ± 0.3 (5)	6.6 ± 0.5 (5)
PND 4 ^{d,e,f}	10.16 ± 0.49 (21/5)	9.69 ± 0.22 (28/7)	10.43 ± 0.47 (28/7)	9.19 ± 0.50 (20/6)	9.95 ± 0.25 (20/5)	9.19 ± 0.60 (18/5)
PND 7 ^{d,e,g}	15.91 ± 0.76 (21/5)	14.95 ± 0.27 (28/7)	16.29 ± 0.71 (28/7)	13.64 ± 0.74 (20/6)	15.00 ± 0.24 (20/5)	14.45 ± 1.16 (14/4)
PND 14 ^{d,e,g}	31.80 ± 0.99 (21/5)	30.48 ± 0.52 (28/7)	32.44 ± 1.22 (28/7)	28.04 ± 1.74 (20/6)	30.44 ± 0.53 (20/5)	28.63 ± 1.33 (14/4)
PND 21 ^{d,e,g}	51.71 ± 1.31** (21/5)	51.21 ± 0.39 (28/7)	53.09 ± 1.58 (28/7)	46.06 ± 2.53 (20/6)	49.40 ± 0.88 (20/5)	45.79 ± 2.04* (14/4)
Male and Female (g)						
PND 1 ^{a,b,c}	7.4 ± 0.2** (6)	7.4 ± 0.1 (8)	7.5 ± 0.2 (7)	7.2 ± 0.2 (6)	7.1 ± 0.3 (6)	6.6 ± 0.3* (6)
PND 4 ^{d,e,f}	10.49 ± 0.45* (40/5)	10.09 ± 0.23 (56/7)	10.65 ± 0.49 (56/7)	9.69 ± 0.32 (48/6)	10.22 ± 0.32 (40/5)	9.24 ± 0.51 (40/5)
PND 7 ^{d,e,g}	16.41 ± 0.72 (40/5)	15.46 ± 0.31 (56/7)	16.48 ± 0.75 (56/7)	14.37 ± 0.36 (48/6)	15.31 ± 0.32 (40/5)	14.52 ± 0.93 (32/4)
PND 14 ^{d,e,g}	32.37 ± 0.88** (40/5)	31.32 ± 0.44 (56/7)	32.79 ± 1.19 (56/7)	29.76 ± 0.84 (48/6)	30.92 ± 0.56 (40/5)	28.91 ± 1.19 (32/4)
PND 21 ^{d,e,g}	53.53 ± 1.29** (40/5)	52.98 ± 0.38 (56/7)	54.07 ± 1.65 (56/7)	49.33 ± 1.55 (48/6)	50.65 ± 0.76 (40/5)	46.08 ± 1.65** (32/4)

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

6 PND = postnatal day.

7 ^aData presented as mean ± standard error (number of dams).

8 ^bEach exposure group was compared to the vehicle control group with the Williams test when a trend was present ($p \leq 0.01$ from the Jonckheere trend test) or with the Dunnett test when no trend was present.

9 ^cLitter weights of live pups at PND 1 were divided by live litter size on PND 1 to obtain a pup mean weight/litter. Mean values were then adjusted using live litter size on PND 1 as a covariate.

10 ^dStatistical analysis was performed using mixed models with random litter effect for both trend and pairwise tests, using the Dunnett-Hsu adjustment for multiple comparisons.

11 ^eData presented as mean of litter means ± standard error (number of pups/number of dams).

12 ^fPND 4 post-standardization.

13 ^gIndividual pup weights first adjusted for live litter size on PND 4 poststandardization.

1 **Three-month Study (Postweaning Phase)**

2 There were no early deaths during the 3-month study; all F₁ rats survived until study termination
3 (Table 7, Table 8). There were no clinical observations related to exposure, and all exposed
4 animals were similar in overt behavior and general appearance to the vehicle control animals
5 (Appendix G). Initial mean body weights were 9% and 16% below those of the vehicle control
6 group for the 1,000 and 2,000 mg/L males, respectively; and 14%, 11%, and 13% below those of
7 the vehicle control group for the 500, 1,000, and 2,000 mg/L females, respectively (Table 7,
8 Table 8; Figure 2). Final mean body weights were lower for the 1,000 and 2,000 mg/L males and
9 females, with the 2,000 mg/L males weighing approximately 29% less than the vehicle control
10 group and the 2,000 mg/L females weighing approximately 18% less than the vehicle control
11 group.

12 Water consumption was lower for the 1,000 and 2,000 mg/L males and females, with overall
13 reductions of 27% and 42% for males and females, respectively, in the 2,000 mg/L groups
14 compared to the respective vehicle control groups (Table 9; Appendix G). Drinking water
15 concentrations of 125, 250, 500, 1,000, and 2,000 mg/L resulted in average daily ST doses of
16 approximately 11.8, 24.3, 48.9, 91.8, and 157.2 mg/kg/day for males and 14.0, 26.1, 54.4, 101.4,
17 and 160.5 mg/kg/day for females.

1 **Table 7. Summary of Survival and Mean Body Weights of Male Rats in the Perinatal and Three-month Drinking Water Study of Sodium**
 2 **Tungstate Dihydrate**

Study Day ^a	0 mg/L		125 mg/L		250 mg/L		500 mg/L		1,000 mg/L		2,000 mg/L						
	Av. Wt. (g) ^b	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters			
1	57.4	5	57.8	100.7	5	57.3	99.9	5	55.2	96.2	5	52.2	91.1	5	48.4	84.3	4
7	82.1	5	81.4	99.2	5	81.5	99.2	5	76.4	93.1	5	71.4	87.0	5	60.0	73.2	4
14	108.1	5	119.0	110.1	5	112.0	103.6	5	109.1	100.9	5	98.3	90.9	5	74.7	69.1	4
21	171.9	5	175.5	102.1	5	172.8	100.6	5	159.8	93.0	5	136.4	79.3	5	91.2	53.1	4
28	191.1	5	193.0	101.0	5	198.1	103.6	5	181.4	94.9	5	168.2	88.0	5	112.4	58.8	4
35	261.7	5	257.7	98.4	5	249.2	95.2	5	245.1	93.6	5	200.0	76.4	5	132.9	50.8	4
42	300.6	5	294.1	97.9	5	298.5	99.3	5	284.8	94.7	5	243.0	80.9	5	160.7	53.5	4
49	319.6	5	322.7	101.0	5	324.6	101.6	5	300.8	94.1	5	267.4	83.7	5	188.6	59.0	4
56	352.5	5	355.1	100.8	5	352.3	99.9	5	339.4	96.3	5	301.5	85.5	5	212.6	60.3	4
63	345.9	5	345.4	99.9	5	341.5	98.7	5	335.2	96.9	5	310.5	89.8	5	222.4	64.3	4
70	384.2	5	383.9	99.9	5	379.5	98.8	5	368.0	95.8	5	339.7	88.4	5	250.1	65.1	4
77	398.1	5	398.3	100.1	5	389.5	97.8	5	385.5	96.8	5	350.4	88.0	5	270.6	68.0	4
84	414.0	5	412.3	99.6	5	407.9	98.5	5	400.0	96.6	5	365.2	88.2	5	285.0	68.8	4
EOS	422.7	5	425.7	100.7	5	419.7	99.3	5	412.9	97.7	5	379.3	89.7	5	300.6	71.1	4

3 EOS = end of study; No. of litters = number of litters represented in weight average.

4 ^aStudy day 1 is the day animals were placed on study after pups were weaned.

5 ^bAverage weights shown are means of litter means.


1 **Table 8. Summary of Survival and Mean Body Weights of Female Rats in the Perinatal and Three-month Drinking Water Study of**
 2 **Sodium Tungstate Dihydrate**

Study Day ^a	0 mg/L		125 mg/L		250 mg/L		500 mg/L		1,000 mg/L		2,000 mg/L						
	Av. Wt. (g) ^b	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters			
1	54.4	5	57.5	105.8	5	56.1	103.1	5	46.7	85.9	5	48.5	89.2	5	47.5	87.4	4
7	79.7	5	81.1	101.8	5	80.7	101.3	5	69.2	86.8	5	70.8	88.8	5	60.5	76.0	4
14	104.1	5	116.0	111.4	5	111.1	106.7	5	100.8	96.8	5	98.4	94.5	5	74.5	71.6	4
21	142.4	5	147.0	103.3	5	150.5	105.7	5	129.6	91.0	5	121.7	85.5	5	86.9	61.1	4
28	165.1	5	173.5	105.1	5	167.2	101.3	5	153.6	93.1	5	141.9	85.9	5	102.2	61.9	4
35	179.2	5	188.9	105.4	5	191.1	106.6	5	175.2	97.8	5	163.7	91.3	5	114.7	64.0	4
42	198.3	5	207.0	104.4	5	208.9	105.4	5	191.0	96.3	5	179.1	90.3	5	132.7	66.9	4
49	210.8	5	223.4	106.0	5	216.3	102.6	5	202.4	96.0	5	191.8	91.0	5	150.7	71.5	4
56	215.5	5	225.4	104.6	5	230.3	106.9	5	217.9	101.1	5	203.8	94.6	5	166.0	77.1	4
63	230.9	5	239.3	103.6	5	243.6	105.5	5	230.0	99.6	5	214.6	92.9	5	177.3	76.8	4
70	238.6	5	255.5	107.1	5	252.8	105.9	5	241.1	101.0	5	222.2	93.1	5	190.0	79.6	4
77	243.4	5	255.8	105.1	5	254.1	104.4	5	244.4	100.4	5	228.6	93.9	5	194.8	80.0	4
84	250.4	5	260.4	104.0	5	260.8	104.2	5	251.7	100.5	5	236.0	94.3	5	201.3	80.4	4
EOS	259.5	5	268.6	103.5	5	271.2	104.5	5	257.9	99.4	5	242.8	93.6	5	213.1	82.1	4


3 EOS = end of study; No. of litters = number of litters represented in weight average.

4 ^aStudy day 1 is the day animals were placed on study after pups were weaned.

5 ^bAverage weights shown are means of litter means.



1



2

3

4

Figure 2. Growth Curves for Rats Exposed to Sodium Tungstate Dihydrate in Drinking Water for Three Months

1 **Table 9. Summary of Water and Sodium Tungstate Dihydrate Consumption of Male and Female Rats in the Perinatal and Three-month**
 2 **Drinking Water Study**

Week	0 mg/L		125 mg/L		250 mg/L		500 mg/L		1,000 mg/L		2,000 mg/L	
	Water (g/day) ^a	Water (g/day)	Dose (mg/kg/day) ^b	Water (g/day)	Dose (mg/kg/day)	Water (g/day)	Dose (mg/kg/day)	Water (g/day)	Dose (mg/kg/day)	Water (g/day)	Dose (mg/kg/day)	
Male												
1	11.5	10.2	18.3	11.1	40.0	9.5	72.2	8.8	142.4	6.6	241.3	
4	23.4	22.3	14.4	23.8	30.0	22.9	63.1	19.9	118.3	10.0	176.3	
12	17.2	17.0	5.2	17.3	10.6	16.6	20.8	14.2	38.9	10.8	75.3	
Female												
1	11.1	11.0	19.8	10.9	39.8	10.0	86.3	9.6	160.9	6.5	241.6	
4	20.9	21.3	15.3	19.2	28.7	19.2	62.5	16.6	117.0	8.8	171.3	
12	20.8	22.2	10.7	19.9	19.1	18.4	36.5	16.9	71.6	12.5	124.1	

3 ^aGrams of water consumed/animal/day.

4 ^bMilligrams of sodium tungstate dihydrate consumed/kilogram body weight/day.

1 Blood was collected from up to 10 animals per group (originating from 4 or 5 litters) on the
 2 morning of day 91. Urine was collected overnight (for approximately 16 hours) during week 12
 3 of the study from up to five animals per group; during collection, animals had access to untreated
 4 water. Total blood and urine tungsten concentrations were determined using validated analytical
 5 methods (Appendix E) and corresponding data are presented in Table 10. In both males and
 6 females, the total tungsten concentration in blood increased proportionally to the exposure
 7 concentration with no observed sex difference. The blood tungsten concentration in vehicle
 8 control animals was below the limit of detection (LOD; 0.0016 $\mu\text{g/g}$) of the assay. The urine
 9 tungsten concentration is presented as both $\mu\text{g/g}$ of urine and after correcting for urinary
 10 creatinine concentrations ($\mu\text{g/mg}$ creatinine) (Table 10). Low concentrations of tungsten were
 11 detected in urine from vehicle control male and female groups. The concentrations of
 12 creatinine-corrected tungsten in urine increased proportionally to the exposure concentration in
 13 both males and females and were significantly increased in all exposed groups compared to the
 14 corresponding vehicle control groups. As with blood, there was no observed sex difference in
 15 urinary tungsten concentrations.

16 **Table 10. Summary of Blood and Urine Tungsten Concentration Data for Male and Female Rats in**
 17 **the Perinatal and Three-month Drinking Water Study of Sodium Tungstate Dihydrate^a**

	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
n	5	5	5	5	5	4
Male						
Blood ($\mu\text{g/g}$)	BD	0.49 \pm 0.07	0.99 \pm 0.10	1.93 \pm 0.21	4.67 \pm 0.45	10.66 \pm 0.93
Urine ^{b,c}						
Urine ($\mu\text{g/g}$ urine)	0.04 \pm 0.00**	9.78 \pm 3.02**	24.21 \pm 5.08**	67.11 \pm 22.66**	61.91 \pm 11.79**	184.98 \pm 30.14**
Urine ($\mu\text{g/mg}$ creatinine)	0.06 \pm 0.01**	11.84 \pm 1.43**	33.68 \pm 4.97**	42.40 \pm 4.64**	86.68 \pm 5.67**	291.52 \pm 25.65**
Female						
Blood ($\mu\text{g/g}$)	BD	0.59 \pm 0.06	1.19 \pm 0.13	2.83 \pm 0.29	5.67 \pm 0.35	11.54 \pm 1.03
Urine ^{b,c}						
Urine ($\mu\text{g/g}$ urine)	0.03 \pm 0.00**	9.10 \pm 1.54**	26.40 \pm 8.95**	27.65 \pm 6.43**	98.98 \pm 14.40**	182.81 \pm 33.44**
Urine ($\mu\text{g/mg}$ creatinine)	0.07 \pm 0.01**	18.88 \pm 1.40**	33.55 \pm 3.82**	45.59 \pm 4.34**	142.92 \pm 21.99**	280.85 \pm 46.16**

18 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

19 Statistical significance for the vehicle control group indicates a significant trend test.

20 **Statistically significant at $p \leq 0.01$.

21 BD = below detection; group did not have more than 20% of its values above the limit of detection (LOD).

22 ^aData presented as mean \pm standard error of the litter means, where n = the number of litters.

23 ^bValues below the LOD (0.0054 $\mu\text{g/g}$) were substituted with 1/2 the LOD value. If 80% or more of the values in the vehicle
 24 control group were below the LOD, no mean or standard error were calculated, and no statistical analysis was performed.

25 ^cStatistical analysis performed using a bootstrapped Jonckheere test for trend and a Datta-Satten modified Wilcoxon test with
 26 Hommel adjustment for pairwise comparisons.

27 In female rats, there was a mild (<10%) significant decrease in the erythron characterized by a
 28 significant decrease in the hemoglobin concentration in the 2,000 mg/L group and a significant
 29 negative trend in the hematocrit concentration, hemoglobin concentration, and erythrocyte count
 30 with increasing exposure (Table 11). Although there were no significant pairwise changes
 31 observed in the male erythron, there were significant negative trends in hematocrit concentration,
 32 hemoglobin concentration, and erythrocyte count with increasing exposure concentration. The

1 reticulocyte count was unchanged in both males and females. These mild erythron changes were
 2 most likely due to the stress of exposure,⁸⁷ which is supported by the lower mean body weights
 3 observed in the 2,000 mg/L groups.

4 In male rats, blood urea nitrogen (BUN) was significantly increased, and the total protein,
 5 globulin concentrations, and insulin concentrations were significantly decreased in the
 6 2,000 mg/L group (Table 11). The BUN was likely increased due to the lower water
 7 consumption values in that exposure group. The toxicological relevance of the observed
 8 decreases in the total protein and globulins is uncertain; these changes could be a secondary
 9 effect of exposure.

10 The urine xanthine/creatinine ratios were significantly increased in all male and female exposed
 11 groups relative to the vehicle control groups (Table 12).

12 **Table 11. Summary of Select Clinical Pathology Data for Male and Female Rats in the Perinatal**
 13 **and Three-month Drinking Water Study of Sodium Tungstate Dihydrate^{a,b}**

	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
n	5	5	5	5	5	4
Male						
Hematocrit (%)	50.0 ± 1.2**	48.2 ± 0.9	49.3 ± 0.5	48.5 ± 0.6	46.8 ± 0.8	45.7 ± 1.1
Hemoglobin (g/dL)	15.2 ± 0.3**	14.9 ± 0.2	15.3 ± 0.1	15.2 ± 0.1	14.6 ± 0.3	14.3 ± 0.3
Erythrocytes (10 ⁶ /μL)	8.69 ± 0.18*	8.45 ± 0.10	8.72 ± 0.10	8.54 ± 0.13	8.23 ± 0.15	8.13 ± 0.22
Urea Nitrogen (mg/dL)	18.1 ± 0.5**	14.8 ± 0.5*	17.3 ± 0.3	17.6 ± 0.7	19.2 ± 0.4	24.4 ± 1.8*
Total Protein (g/dL)	6.64 ± 0.04	6.67 ± 0.06	6.76 ± 0.09	6.75 ± 0.10	6.73 ± 0.07	6.04 ± 0.07*
Globulin (g/dL)	2.46 ± 0.05*	2.37 ± 0.03	2.48 ± 0.07	2.48 ± 0.05	2.37 ± 0.05	1.86 ± 0.04*
Insulin (ng/mL)	4.36 ± 0.29**	3.36 ± 0.33	3.09 ± 0.40	3.36 ± 0.42	2.52 ± 0.33	1.94 ± 0.34*
Female						
Hematocrit (%)	44.7 ± 0.6*	45.5 ± 0.9	44.4 ± 0.6	44.6 ± 0.7	44.5 ± 0.8	42.3 ± 0.2
Hemoglobin (g/dL)	14.2 ± 0.1*	14.3 ± 0.3	14.1 ± 0.2	14.2 ± 0.1	14.0 ± 0.2	13.4 ± 0.1*
Erythrocytes (10 ⁶ /uL)	7.80 ± 0.07*	7.87 ± 0.14	7.78 ± 0.05	7.70 ± 0.08	7.73 ± 0.07	7.38 ± 0.12

14 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

15 Statistical significance for the vehicle control group indicates a significant trend test.

16 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

17 ^aData presented as mean ± standard error of the litter means, where n = the number of litters.

18 ^bStatistical analysis performed using a bootstrapped Jonckheere test for trend and a Datta-Satten modified Wilcoxon test with
 19 Hommel adjustment for pairwise comparisons.

1 **Table 12. Summary of Select Urinalysis Data for Male and Female Rats in the Perinatal and**
 2 **Three-month Drinking Water Study of Sodium Tungstate Dihydrate^{a,b}**

	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
n	5	5	5	5	5	4
Male						
Creatinine (mg/dL)	77.9 ± 10.2	80.5 ± 22.6	76.2 ± 11.3	154.5 ± 44.5	73.9 ± 15.1	73.5 ± 11.8
Xanthine (µg/mL)	1.20 ± 0.16**	1.51 ± 0.44	1.58 ± 0.24	4.48 ± 1.75	3.20 ± 0.52*	5.45 ± 0.88*
Xanthine/Creatinine (µg/mg)	1.55 ± 0.09**	1.83 ± 0.04*	2.13 ± 0.08*	2.79 ± 0.37*	4.63 ± 0.27*	7.65 ± 0.26*
Female						
Creatinine (mg/dL)	43.7 ± 6.1*	47.8 ± 4.6	77.1 ± 22.9	58.9 ± 10.2	66.7 ± 4.1	82.3 ± 22.3
Xanthine (µg/mL)	0.43 ± 0.07**	0.74 ± 0.12	1.77 ± 0.54*	2.00 ± 0.35*	3.30 ± 0.21*	3.72 ± 0.75
Xanthine/Creatinine (µg/mg)	1.02 ± 0.09**	1.51 ± 0.14*	2.27 ± 0.27**	3.39 ± 0.18**	5.21 ± 0.38**	7.28 ± 0.86*

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

6 ^aData presented as mean ± standard error of the litter means, where n = the number of litters.

7 ^bStatistical analysis performed using a bootstrapped Jonckheere test for trend and a Datta-Satten modified Wilcoxon test with
 8 Hommel adjustment for pairwise comparisons.

9 Absolute kidney weights were reduced in males in all exposed groups, relative to the vehicle
 10 control group, with a significant decrease observed in the 2,000 mg/L group (approximately
 11 21%). Relative kidney weights were higher in 1,000 mg/L females and significantly increased in
 12 the 2,000 mg/L males and females, relative to the vehicle control group (Table 13). Although the
 13 kidney was a target tissue, it is unlikely that the lesions observed were responsible for the
 14 differences in kidney weights; it is more likely that these organ weight differences are an effect
 15 of body weight differences.

16 When compared to vehicle control groups, significant differences were also observed in other
 17 organ weights, including decreased absolute heart and lung weights in males and females;
 18 decreased absolute liver weights in males and increased relative liver weights in females;
 19 decreased absolute thymus weights in males and increased relative testis weights (Appendix G).
 20 These changes were considered secondary to body weight reductions. Rats administered
 21 2,000 mg/L ST exhibited significantly decreased left cauda epididymis (14%) and epididymis
 22 (13%) weights, and lower testis weights (8%) compared to the vehicle control group
 23 (Appendix G). Although these were significant (cauda and epididymis) and/or displayed a
 24 significant negative trend with increasing exposure concentration (right testis), rats in the 2,000
 25 mg/L group displayed mean body weights that were 28% lower than the vehicle control group.
 26 There were no changes in reproductive parameters or alterations in contralateral testis and
 27 epididymis or in histopathology (Appendix G). Given the magnitude of the body weight effect
 28 and the absence of changes in other endpoints, the lower reproductive organ weights are likely
 29 secondary to effects on body weight.

30 Although the weights of the left epididymis and the left cauda were significantly decreased in the
 31 2,000 mg/L males, there were no corresponding changes in sperm parameters, including number

1 of sperm/mg cauda epididymis, total number of sperm/cauda, sperm motility, number of
2 homogenization-resistant spermatids/mg testis, or total number of spermatids (Appendix G).

3 No exposure-related gross lesions were recorded. Exposure-related histological lesions were
4 found in the kidneys (Table 13). Renal tubule regeneration was increased in the male and female
5 1,000 and 2,000 mg/L groups; the increases in the 2,000 mg/L groups were significant relative to
6 the vehicle control groups (Table 13). The lesion was characterized by hyperplasia of proximal
7 convoluted tubular epithelial cells that manifested as cytoplasmic basophilia, nuclear crowding,
8 and occasional mitotic figures. Renal tubule regeneration differed from chronic progressive
9 nephropathy (CPN) by the lack of thickened basement membranes, associated inflammatory
10 cells, proteinaceous casts, and cytoplasmic pigment—all features typically seen with CPN. The
11 incidences and severities of CPN were not increased in exposed groups of animals
12 (Appendix G).

13 **Table 13. Summary of Renal Findings for Male and Female Rats in the Perinatal and Three-month**
14 **Drinking Water Study of Sodium Tungstate Dihydrate**

	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
n	5	5	5	5	5	4
Male						
Necropsy Body Wt. (g) ^a	422.7 ± 5.2**	425.7 ± 3.9	419.7 ± 9.0	412.9 ± 10.2	379.3 ± 6.7**	300.6 ± 17.9**
R. Kidney Weight ^a						
Absolute (g)	1.35 ± 0.04**	1.32 ± 0.03	1.25 ± 0.05	1.26 ± 0.04	1.22 ± 0.02	1.06 ± 0.06**
Relative (mg/g) ^b	3.20 ± 0.08**	3.11 ± 0.06	2.99 ± 0.10	3.06 ± 0.03	3.21 ± 0.01	3.56 ± 0.09**
Histological Findings						
Kidney ^c	10	10	10	10	10	10
Renal tubule, regeneration ^d	0**	0	0	0	3 (1.0) ^e	10** (2.0)
Female						
Necropsy Body Wt. (g)	259.5 ± 3.8**	268.6 ± 9.1	271.2 ± 7.0	257.9 ± 6.3	242.8 ± 0.8	213.1 ± 4.1**
R. Kidney Weight						
Absolute (g)	0.84 ± 0.02	0.87 ± 0.03	0.84 ± 0.02	0.81 ± 0.02	0.84 ± 0.02	0.85 ± 0.03
Relative (mg/g)	3.25 ± 0.04**	3.25 ± 0.03	3.08 ± 0.05	3.13 ± 0.05	3.47 ± 0.07	3.98 ± 0.15**
Histological Findings						
Kidney	10	10	10	10	10	10
Renal tubule, regeneration	0**	0	0	0	3 (1.0)	10** (2.0)

15 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

16 Statistical significance for the vehicle control group indicates a significant trend test.

17 Statistical analysis for organ weight data performed using mixed models, with litter as a random effect and a Dunnett-Hsu
18 adjustment for multiple comparisons. Statistical analysis for histological findings performed by the Rao-Scott test.

19 **Statistically significant at $p \leq 0.01$.

20 ^aData presented as mean ± standard error of the litter means.

21 ^bRelative organ weights (organ-weight-to-body-weight ratios) are given as mg organ weight/g body weight.

22 ^cNumber of animals examined microscopically.

23 ^dNumber of animals with lesion.

24 ^eAverage severity grade of observed lesion in affected animals: 1 = minimal; 2 = mild; 3 = moderate; 4 = marked.

1 **Exposure Concentration Selection Rationale for the Perinatal and Two-year** 2 **Study in Rats**

3 A >10% reduction in final mean body weight in males and females (Table 7, Table 8) and up to a
4 65% reduction in water consumption for males (at week 3), and a 60% reduction in water
5 consumption for females (at week 3) (Table 9) in the 3-month studies coupled with lesions noted
6 in the kidney (Table 13), led to the assessment that 2,000 mg/L was too high for the 2-year
7 studies. The 1,000 mg/L concentration was considered an adequate top exposure concentration to
8 challenge the animals based on an approximate 10% reduction in final mean body weight in
9 males, an approximate 20% reduction in water consumption in females, and minimal to mild
10 renal tubule regeneration in the 3-month studies. Because there was no overlap in tungstate blood
11 concentrations between the 1,000 mg/L and 2,000 mg/L groups, and no significant effects were
12 noted at 500 mg/L or lower, the exposure concentrations were spaced by half. Hence, exposure
13 concentrations selected for the chronic studies were 0, 250, 500 and 1,000 mg/L. Additionally,
14 tissue tungsten concentrations were evaluated in the kidney, plasma, and urine at 3, 6, 12, and 18
15 months in an additional group of animals to determine systemic exposure and to help identify
16 species differences given the findings from the 3-month studies showing the kidney as a target
17 organ of toxicity.

18 **Two-year Study (Perinatal Phase)**

19 No significant clinical observations were noted in dams and no significant effects were noted on
20 reproductive performance, including the percentage of mated females producing pups (Table 14).
21 Gestational mean body weights of dams in the 1,000 mg/L group were less than that of the
22 vehicle control group by up to 5% by GD 21 (Table 15). During lactation, mean body weights of
23 dams in the 1,000 mg/L group were significantly decreased by 3%, 3%, and 4% relative to the
24 vehicle control group at LD 1, LD 17, and LD 21, respectively (Table 15).

25 **Table 14. Summary of the Disposition of F₀ Female Rats during Perinatal Exposure in the Perinatal**
26 **and Two-year Drinking Water Study of Sodium Tungstate Dihydrate**

	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
Reproductive Performance				
Time-mated Females (GD 6)	47	41	41	41
Females Pregnant (%) ^a	41 (87.2)	34 (82.9)	36 (87.8)	36 (87.8)
Females Not Pregnant (%)	6 (12.8)	7 (17.1)	5 (12.2)	5 (12.2)
Dams Not Delivering with Evidence of Pregnancy (%)	0	0	1 (2.8) ^b	2 (5.6)
Dams with Litters on LD 0 (%)	41 (100.0)	34 (100.0)	35 (100.0)	34 (94.4)
Litters Poststandardization (LD 4) ^c	37	33	33	32

27 GD = gestation day; LD = lactation day.

28 ^aStatistical analysis performed by the Cochran-Armitage (trend) and Fisher's exact (pairwise) tests.

29 ^bDam died before littering.

30 ^cStandardization to eight pups/litter (four pups/sex). Only litters with at least two pups/sex and at least eight pups total/litter were
31 retained to continue on study.

1 **Table 15. Summary of Mean Body Weights and Body Weight Gains of F₀ Female Rats during**
 2 **Gestation and Lactation in the Perinatal and Two-year Drinking Water Study of Sodium Tungstate**
 3 **Dihydrate**

Parameter ^{a,b}	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
Gestation Day				
6	233.4 ± 1.87 (41)	236.6 ± 2.01 (34)	236.2 ± 1.71 (36)	233.4 ± 1.82 (36)
9	251.4 ± 2.07 (41)	252.9 ± 2.10 (34)	250.6 ± 1.93 (36)	248.2 ± 1.61 (36)
12	265.7 ± 1.81 (41)	267.1 ± 2.36 (34)	264.5 ± 1.90 (36)	260.7 ± 1.73 (36)
15	288.2 ± 2.21** (41)	289.0 ± 2.45 (34)	286.3 ± 2.05 (36)	279.2 ± 1.90** (36)
18	325.7 ± 2.75** (41)	326.9 ± 2.85 (34)	323.8 ± 2.56 (36)	312.6 ± 2.96** (36)
21	373.7 ± 4.07** (41)	377.0 ± 3.44 (34)	371.0 ± 3.45 (36)	354.3 ± 4.76** (36)
Gestation Weight Change				
Gestation Day Interval				
6–9	17.9 ± 0.8** (41)	16.3 ± 0.5 (34)	14.3 ± 0.6** (36)	14.8 ± 0.9** (36)
9–12	14.3 ± 1.0 (41)	14.2 ± 0.7 (34)	14.0 ± 0.5 (36)	12.5 ± 0.6 (36)
12–15	22.5 ± 0.7** (41)	21.9 ± 0.6 (34)	21.8 ± 0.6 (36)	18.5 ± 0.8** (36)
15–18	37.5 ± 1.2 (41)	37.9 ± 0.9 (34)	37.5 ± 0.9 (36)	33.4 ± 1.6* (36)
18–21	48.0 ± 1.6** (41)	50.1 ± 1.2 (34)	47.2 ± 1.3 (36)	41.7 ± 2.1** (36)
6–21	140.3 ± 3.6** (41)	140.4 ± 2.1 (34)	134.8 ± 2.6 (36)	120.9 ± 4.8** (36)
Lactation Day				
1	282.4 ± 2.04** (41)	282.7 ± 2.49 (34)	276.9 ± 2.01 (35)	273.8 ± 2.16** (34)
4	295.1 ± 2.15* (41)	295.7 ± 2.58 (34)	291.8 ± 2.42 (35)	288.7 ± 2.48 (34)
7	302.6 ± 2.04 (37)	303.8 ± 2.65 (33)	297.0 ± 2.74 (33)	298.4 ± 2.07 (32)
10	312.8 ± 2.4* (37)	310.4 ± 3.0 (32)	308.6 ± 2.7 (32)	305.0 ± 2.4 (32)
14	321.2 ± 2.45 (37)	321.5 ± 2.66 (32)	320.0 ± 2.66 (32)	316.6 ± 2.48 (32)
17	320.5 ± 2.35* (37)	321.0 ± 3.12 (32)	321.9 ± 3.20 (32)	309.9 ± 2.13* (32)
21	301.6 ± 2.70* (37)	306.4 ± 3.45 (32)	299.8 ± 3.84 (32)	289.2 ± 4.05* (32)
Lactation Weight Change				
Lactation Day Interval				
1–4	12.7 ± 1.2 (41)	12.9 ± 1.5 (34)	15.0 ± 1.2 (35)	15.0 ± 2.2 (34)
4–7	7.5 ± 1.2 (37)	8.6 ± 1.1 (33)	4.8 ± 1.5 (33)	8.7 ± 1.9 (32)
7–10	10.2 ± 1.4 (37)	6.4 ± 1.4 (32)	10.9 ± 1.2 (32)	6.5 ± 1.6 (32)
10–14	8.4 ± 1.6 (37)	11.0 ± 2.0 (32)	11.4 ± 1.6 (32)	11.6 ± 2.0 (32)
14–17	-0.7 ± 2.1 (37)	-0.5 ± 1.8 (32)	1.8 ± 1.7 (32)	-6.7 ± 2.0 (32)
17–21	-18.9 ± 2.6 (37)	-14.5 ± 2.8 (32)	-22.1 ± 3.3 (32)	-20.7 ± 3.5 (32)
1–21	19.5 ± 2.9 (37)	24.2 ± 2.5 (32)	22.5 ± 3.4 (32)	15.0 ± 4.1 (32)

4 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

5 Statistical significance for the vehicle control group indicates a significant trend test.

6 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

7 ^aEach exposure group was compared to the vehicle control group with the Williams test when a trend was present ($p \leq 0.01$ from the Jonckheere trend test) or with the Dunnett test when no trend was present.

9 ^bData presented as mean ± standard error (number of dams). Body weight data are presented in grams.

1 Summaries of water and ST consumption are presented in Table 16. Water consumption trends
 2 were similar during gestation and lactation. During gestation, water consumption of all
 3 ST-exposed groups was within 9% of the vehicle control group (ranging from 99% to 109%).
 4 During lactation, water consumption of all ST-exposed groups was within 8% of the vehicle
 5 control group (ranging from 92% to 100%) (Table 16). Daily ST consumption for the 250, 500,
 6 and 1,000 mg/L groups averaged approximately 32, 65, and 143 mg/kg/day, respectively, during
 7 GD 6–21, and approximately 48, 98, and 197 mg/kg/day, respectively, during LD 1–14
 8 (Table 16).

9 **Table 16. Summary of Water and Sodium Tungstate Dihydrate Consumption by F₀ Female Rats**
 10 **during Gestation and Lactation in the Perinatal and Two-year Drinking Water Study**

Parameter ^a	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
Gestation Day Interval^{b,c}				
6–9	29.72 ± 0.83* (39)	29.89 ± 0.77 (34)	30.21 ± 0.91 (35)	33.29 ± 1.03 (36)
9–12	31.55 ± 0.96 (41)	32.01 ± 0.93 (34)	31.92 ± 0.82 (36)	34.46 ± 1.04 (36)
12–15	36.94 ± 0.79 (41)	36.47 ± 0.89 (34)	35.33 ± 0.91 (36)	38.93 ± 1.06 (36)
15–18	43.39 ± 1.00* (41)	44.50 ± 1.23 (34)	43.54 ± 1.09 (36)	47.06 ± 1.38* (36)
18–21	45.27 ± 1.16 (41)	45.61 ± 1.25 (34)	44.43 ± 1.13 (36)	46.40 ± 1.54 (36)
6–21	37.31 ± 0.83* (39)	37.70 ± 0.91 (34)	37.19 ± 0.89 (35)	40.03 ± 1.08 (36)
Lactation Day Interval^{b,c}				
1–4	47.08 ± 1.08 (41)	46.17 ± 1.07 (34)	46.03 ± 0.93 (35)	46.36 ± 1.22 (34)
4–7	50.35 ± 1.05 (37)	49.85 ± 1.09 (33)	47.90 ± 0.92 (33)	48.68 ± 0.81 (32)
7–10	63.61 ± 1.51 (37)	60.39 ± 1.27 (32)	60.49 ± 1.20 (32)	60.80 ± 1.36 (32)
10–14	74.39 ± 1.27 (35)	74.75 ± 1.18 (31)	76.51 ± 1.43 (32)	74.59 ± 1.49 (32)
14–17	86.82 ± 1.69* (37)	82.54 ± 1.50 (32)	85.16 ± 1.93 (32)	80.28 ± 1.85 (32)
17–21	96.58 ± 1.57 (37)	91.38 ± 1.92 (32)	90.17 ± 2.46 (32)	90.51 ± 2.68 (31)
1–21	59.71 ± 1.04 (35)	58.86 ± 0.92 (31)	59.17 ± 0.92 (32)	59.10 ± 1.04 (32)
Chemical Intake (mg/kg/day)^{d,e}				
GD 6–21	0.00 ± 0.00 (39)	32.49 ± 0.68 (34)	64.77 ± 1.52 (35)	143.2 ± 3.70 (36)
LD 1–14	0.00 ± 0.00 (35)	48.14 ± 0.72 (31)	97.92 ± 1.53 (32)	197.3 ± 3.44 (32)

11 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

12 Statistical significance for the vehicle control group indicates a significant trend test.

13 *Statistically significant at $p \leq 0.05$.

14 GD = gestation day; LD = lactation day.

15 ^aData presented as mean ± standard error (number of dams).

16 ^bWater consumption data are presented as grams/animal/day.

17 ^cEach exposure group was compared to the vehicle control group with the Shirley test when a trend was present ($p \leq 0.01$ from the Jonckheere trend test) or with the Dunn test when no trend was present.

18 ^dChemical intake calculated as: ([exposure concentration × water consumption]/[average body weight of day range]).

19 ^eNo statistical analysis was performed on the chemical intake data.

21 No significant clinical observations were noted in pups (Appendix G). No exposure-related
 22 differences were noted between the vehicle control groups and the ST-exposed groups in the
 23 number of litters, litter size, mean litter weights, sex ratio, or the pup mean weights of males and

1 females on PND 1 or PND 4 (Table 17, Table 18). Litter size did not change appreciably
 2 between PND 1 and litter standardization on PND 4. Pups were weaned on PND 21, and this was
 3 considered day 1 of the 2-year exposure period.

4 **Table 17. Summary of Mean Litter Size and Survival Ratio of F₁ Male and Female Rats during**
 5 **Lactation in the Perinatal and Two-year Drinking Water Study of Sodium Tungstate Dihydrate**

Parameter	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
PND 1^{a,b}				
Total	12.83 ± 0.45 (41)	13.15 ± 0.31 (34)	13.03 ± 0.36 (35)	11.82 ± 0.49 (34)
Live	12.51 ± 0.49 (41)	12.82 ± 0.33 (34)	12.69 ± 0.37 (35)	11.68 ± 0.51 (34)
% Male per Litter	51.83 ± 2.36 (41)	49.27 ± 3.10 (34)	47.95 ± 2.53 (35)	51.42 ± 2.04 (34)
% Male ^{c,d}	52 (513)	50 (436)	48 (441)	52 (397)
Male^{a,b}				
PND 1	6.54 ± 0.42 (41)	6.35 ± 0.44 (34)	6.11 ± 0.38 (35)	6.03 ± 0.37 (34)
PND 4 Prestandardization	6.51 ± 0.42 (41)	6.29 ± 0.44 (34)	6.11 ± 0.38 (35)	6.03 ± 0.37 (34)
PND 4 Poststandardization	4.05 ± 0.12 (37)	4.00 ± 0.13 (33)	3.97 ± 0.08 (33)	3.97 ± 0.07 (32)
Female^{a,b}				
PND 1	5.98 ± 0.33 (41)	6.47 ± 0.37 (34)	6.57 ± 0.33 (35)	5.56 ± 0.34 (34)
PND 4 Prestandardization	5.85 ± 0.34 (41)	6.38 ± 0.39 (34)	6.49 ± 0.34 (35)	5.56 ± 0.33 (34)
PND 4 Poststandardization	3.95 ± 0.12 (37)	4.00 ± 0.13 (33)	4.03 ± 0.08 (33)	4.03 ± 0.07 (32)
Male and Female^{a,b}				
PND 4 Prestandardization	12.37 ± 0.49 (41)	12.68 ± 0.34 (34)	12.60 ± 0.39 (35)	11.59 ± 0.51 (34)
PND 4 Poststandardization	8.00 ± 0.00 (37)	8.00 ± 0.00 (33)	8.00 ± 0.00 (33)	8.00 ± 0.00 (32)
PND 21	7.92 ± 0.05 (37)	7.94 ± 0.04 (32)	7.94 ± 0.04 (32)	7.81 ± 0.13 (32)
Survival per Litter				
Total Dead: PND 1–4 ^{d,e}	19 (41)	16 (34)	15 (35)	8 (34)
Total Dead: PND 5–21 ^{d,e}	3 (37)	2 (32)	2 (32)	6 (32)
Dead: PND 1–4 ^{a,b,f}	0.463 ± 0.140 (41)	0.471 ± 0.185 (34)	0.429 ± 0.170 (35)	0.235 ± 0.085 (34)
Dead: PND 4–21 ^{a,b,f}	0.081 ± 0.045 (37)	0.063 ± 0.043 (32)	0.063 ± 0.043 (32)	0.188 ± 0.130 (32)
Survival Ratio: PND 1–4 ^{a,b,g}	0.988 ± 0.007 (41)	0.988 ± 0.007 (34)	0.991 ± 0.005 (35)	0.993 ± 0.004 (34)
Survival Ratio: PND 4–21 ^{a,b,h}	0.990 ± 0.006 (37)	0.992 ± 0.005 (32)	0.992 ± 0.005 (32)	0.977 ± 0.016 (32)

6 PND = postnatal day.

7 ^aEach exposure group was compared to the vehicle control group with the Shirley test when a trend was present ($p \leq 0.01$ from
 8 the Jonckheere trend test) or with the Dunn test when no trend was present.

9 ^bData presented as mean ± standard error (number of dams).

10 ^c $100 \times [\text{number of live males in exposure group}] / [\text{number of live males and females in exposure group}]$.

11 ^dNo statistics done on this endpoint.

12 ^eTotal dead in exposure group (number of dams).

13 ^fNumber dead/litter.

14 ^gSurvival per litter: number of pups prestandardization on PND 4/total live pups on PND 1.

15 ^hSurvival per litter: number of live pups on PND 21/number of live pups poststandardization on PND 4.

1 **Table 18. Summary of Prewaning F₁ Male and Female Rat Pup Mean Body Weights Following**
 2 **Perinatal Exposure to Sodium Tungstate Dihydrate**

Parameter	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
Male (g)				
PND 1 ^{a,b,c}	6.82 ± 0.07 (41)	6.80 ± 0.08 (34)	6.74 ± 0.07 (35)	6.69 ± 0.11 (34)
PND 4 ^{d,e,f}	9.80 ± 0.15 (267/41)	9.30 ± 0.22 (214/34)	9.75 ± 0.15 (214/35)	9.18 ± 0.21* (205/34)
PND 7 ^{d,e,g}	14.97 ± 0.35 (150/37)	14.19 ± 0.37 (131/33)	14.70 ± 0.35 (131/33)	14.35 ± 0.34 (125/32)
PND 14 ^{d,e,g}	30.97 ± 0.65 (148/37)	30.12 ± 0.48 (126/32)	30.26 ± 0.55 (126/32)	29.84 ± 0.55 (125/32)
PND 21 ^{d,e,g}	51.32 ± 1.05 (148/37)	49.48 ± 0.86 (126/32)	49.77 ± 0.94 (126/32)	48.96 ± 1.08 (125/32)
Female (g)				
PND 1 ^{a,b,c}	6.43 ± 0.08 (41)	6.50 ± 0.09 (34)	6.46 ± 0.07 (35)	6.34 ± 0.09 (34)
PND 4 ^{d,e,f}	9.31 ± 0.17 (240/41)	8.98 ± 0.22 (217/34)	9.43 ± 0.15 (226/35)	8.86 ± 0.18 (189/34)
PND 7 ^{d,e,g}	14.19 ± 0.35 (146/37)	13.67 ± 0.41 (130/33)	14.21 ± 0.36 (131/33)	13.56 ± 0.34 (128/32)
PND 14 ^{d,e,g}	29.71 ± 0.65 (145/37)	29.17 ± 0.52 (128/32)	29.55 ± 0.63 (128/32)	28.39 ± 0.69 (125/32)
PND 21 ^{d,e,g}	48.27 ± 1.03 (145/37)	47.00 ± 0.87 (128/32)	47.59 ± 0.92 (128/32)	46.26 ± 1.10 (125/32)
Male and Female (g)				
PND 1 ^{a,b,c}	6.63 ± 0.07 (41)	6.65 ± 0.09 (34)	6.60 ± 0.07 (35)	6.52 ± 0.10 (34)
PND 4 ^{d,e,f}	9.58 ± 0.16 (507/41)	9.14 ± 0.22 (431/34)	9.59 ± 0.15 (440/35)	9.05 ± 0.19 (394/34)
PND 7 ^{d,e,g}	14.60 ± 0.34 (296/37)	13.94 ± 0.37 (261/33)	14.47 ± 0.36 (262/33)	13.95 ± 0.34 (253/32)
PND 14 ^{d,e,g}	30.37 ± 0.63 (293/37)	29.64 ± 0.49 (254/32)	29.91 ± 0.58 (254/32)	29.18 ± 0.54 (250/32)
PND 21 ^{d,e,g}	49.84 ± 1.01 (293/37)	38.23 ± 0.85 (254/32)	48.71 ± 0.91 (254/32)	47.64 ± 1.01 (250/32)

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 *Statistically significant at p ≤ 0.05.

5 PND = postnatal day.

6 ^aData presented as mean ± standard error (number of dams).

7 ^bEach exposure group was compared to the vehicle control group with the Williams test when a trend was present (p ≤ 0.01 from the Jonckheere trend test) or with the Dunnett test when no trend was present.

8 ^cLitter weights of live pups at PND 1 were divided by live litter size on PND 1 to obtain a pup mean weight/litter. Mean values were then adjusted using live litter size on PND 1 as a covariate.

9 ^dStatistical analysis was performed using mixed models with random litter effect for both trend and pairwise tests, using the Dunnett-Hsu adjustment for multiple comparisons.

10 ^eData presented as the mean of litter means ± standard error (number of pups/number of dams).

11 ^fPND 4 prestandardization.

12 ^gIndividual pup weights first adjusted for live litter size on PND 4 poststandardization.

1 Plasma from up to five dams per exposure group along with plasma from one pup/sex/dam were
 2 collected on PND 21. Total tungsten concentrations were determined using a validated analytical
 3 method (Appendix E). In dams, tungsten concentrations increased with the exposure
 4 concentration (Table 19). Tungsten was also detected in male and female pups with
 5 concentrations slightly lower than in dams, suggesting significant exposure of pups to tungsten
 6 via lactation and/or from direct consumption of dosed drinking water. There was no apparent sex
 7 difference in tungsten concentrations in the pups.

8 **Table 19. Summary of Internal Dose Data for F₀ Female Rats and Pups in the Perinatal and**
 9 **Two-year Drinking Water Study of Sodium Tungstate Dihydrate^a**

	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
n	5	5	5	5
Tungsten Concentration (µg/mL)				
Postnatal Day 21				
Dam plasma	BD	1.54 ± 0.17	2.35 ± 0.10	3.82 ± 0.84
Male pup plasma	BD	0.84 ± 0.17	1.48 ± 0.30	3.68 ± 0.89
Female pup plasma	BD	0.76 ± 0.26	1.35 ± 0.18	2.58 ± 0.56 ^b

10 BD = below detection; group did not have over 20% of its values above the limit of detection (LOD; 0.013 µg/mL).

11 ^aData are presented as mean ± standard error.

12 ^bn = 4.

1 **Two-year Study (Interim Evaluations – 3, 6, 12, and 18 Months)**

2 Ten predesignated rats/sex from each exposure group were evaluated at 3, 6, 12, or 18 months
3 for organ weights and tungsten concentrations. Mean body weights of the 500 mg/L males were
4 significantly decreased by approximately 12% relative to the vehicle control group at the
5 12-month time point (Table 20). Mean body weights of all other groups, both males and females,
6 were within 10% of their respective vehicle control group at all interim evaluations (Table 20,
7 Table 21).

8 The mean relative kidney weights were significantly increased at 3 months in the 1,000 mg/L
9 males relative to the vehicle control group (Table 20); in females, there were no significant
10 differences in kidney weights between exposed groups and the vehicle control group (Table 21).
11 At 6 months, there were no significant differences in the kidney weights of males (Table 20), but
12 in females, the mean absolute kidney weights were significantly increased in the 500 mg/L and
13 1,000 mg/L groups by approximately 6% and 7%, respectively, compared to the vehicle control
14 group, and the mean relative kidney weight was significantly increased in the 1,000 mg/L group
15 (Table 21). By 12 months, the mean absolute kidney weight was significantly decreased by
16 approximately 13% in the 500 mg/L males compared to the vehicle control males (Table 20), and
17 in females, the mean relative kidney weights were significantly increased in the 500 and
18 1,000 mg/L groups, relative to the vehicle control group (Table 21). At 18 months, mean
19 absolute kidney weights were significantly decreased in the 500 and 1,000 mg/L males, by
20 approximately 16% and 28%, respectively, relative to the vehicle control group (Table 20),
21 whereas in females, there were no significant differences in kidney weights (Table 21).

22 At 3 months, the mean absolute liver weights were significantly decreased relative to the vehicle
23 control group by approximately 12% in the 250 mg/L males. By 12 months, the mean absolute
24 liver weights were significantly decreased relative to the vehicle control group in all male
25 exposed groups by approximately 11%, 16%, and 17% in the 250, 500 and 1,000 mg/L males,
26 respectively (Table 20). At 18 months, the mean absolute liver weight was significantly
27 decreased relative to the vehicle control group by approximately 18% in the 1,000 mg/L males
28 (Table 20). In females, the trend was significant for decreased absolute liver weight with
29 increasing exposure concentration at 18 months (Table 21). The biological importance of these
30 changes in liver weights is unknown; no liver effects were observed histologically in the 2-year
31 study.

32 Other sporadic differences in organ weights were considered isolated changes of no toxicological
33 significance (Appendix G).

1 **Table 20. Summary of Select Organ Weights and Organ-Weight-to-Body-Weight Ratios for Male**
 2 **Rats Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months^{a,b}**

	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
Three Months^c	10	10	10	10
Necropsy Body Wt. (g)	420.7 ± 8.3*	387.7 ± 8.9*	407.8 ± 11.0*	384.0 ± 6.3*
Kidneys				
Absolute (g)	2.82 ± 0.07	2.72 ± 0.07	2.81 ± 0.11	2.76 ± 0.08
Relative (mg/g) ^d	6.71 ± 0.10*	7.02 ± 0.12	6.87 ± 0.14	7.17 ± 0.14*
Liver				
Absolute (g)	16.95 ± 0.55	14.84 ± 0.51*	16.30 ± 0.74	15.48 ± 0.55
Relative (mg/g)	40.23 ± 0.84	38.21 ± 0.65	39.85 ± 1.09	40.24 ± 0.96
Six Months	10	10	10	10
Necropsy Body Wt. (g)	481.4 ± 14.1	473.9 ± 9.9	467.3 ± 5.7	452.6 ± 12.0
Kidneys				
Absolute (g)	2.96 ± 0.12	2.87 ± 0.08	2.84 ± 0.07	2.75 ± 0.05
Relative (mg/g)	6.14 ± 0.11	6.05 ± 0.10	6.08 ± 0.14	6.09 ± 0.11
Liver				
Absolute (g)	15.55 ± 0.75	14.96 ± 0.59	15.03 ± 0.41	14.02 ± 0.50
Relative (mg/g)	32.16 ± 0.79	31.48 ± 0.72	32.19 ± 0.85	30.95 ± 0.63
Twelve Months	8	10	10	10
Necropsy Body Wt. (g)	586.4 ± 18.8*	571.9 ± 16.3	515.3 ± 15.8*	524.8 ± 18.5
Kidneys				
Absolute (g)	3.57 ± 0.09	3.45 ± 0.10	3.10 ± 0.11*	3.31 ± 0.14
Relative (mg/g)	6.11 ± 0.19	6.04 ± 0.15	6.03 ± 0.19	6.32 ± 0.20
Liver				
Absolute (g)	21.84 ± 0.93**	19.46 ± 0.44*	18.26 ± 0.73**	18.09 ± 0.91**
Relative (mg/g)	37.25 ± 1.08	34.20 ± 0.95	35.40 ± 0.81	34.38 ± 0.79
Eighteen Months	8	9	8	9
Necropsy Body Wt. (g)	604.9 ± 24.1*	609.5 ± 14.2	594.2 ± 15.6	500.0 ± 33.6*
Kidneys				
Absolute (g)	4.57 ± 0.36**	4.04 ± 0.19	3.84 ± 0.11*	3.28 ± 0.18**
Relative (mg/g)	7.72 ± 0.81	6.64 ± 0.34	6.48 ± 0.23	6.67 ± 0.32
Liver				
Absolute (g)	21.01 ± 0.64**	21.98 ± 0.78	19.91 ± 1.10	17.28 ± 1.31*
Relative (mg/g)	34.90 ± 0.96	36.03 ± 0.75	33.46 ± 1.46	34.47 ± 0.82

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

6 ^aData presented as mean ± standard error.

7 ^bStatistical analysis performed by the Jonckheere (trend) and the Williams or Dunnett (pairwise) tests.

8 ^cNumber of animals examined at each time point.

9 ^dRelative organ weights (organ-weight-to-body-weight ratios) are given as mg organ weight/g body weight.

1 **Table 21. Summary of Select Organ Weights and Organ-Weight-to-Body-Weight Ratios for Female**
 2 **Rats Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months^{a,b}**

	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
Three Months^c	10	10	10	10
Necropsy Body Wt. (g)	251.6 ± 3.9	257.2 ± 6.9	245.6 ± 5.9	236.9 ± 6.5
Kidneys				
Absolute (g)	1.67 ± 0.03	1.66 ± 0.05	1.66 ± 0.06	1.67 ± 0.05
Relative (mg/g) ^d	6.64 ± 0.09	6.45 ± 0.14	6.74 ± 0.20	7.05 ± 0.13
Liver				
Absolute (g)	9.01 ± 0.14	9.25 ± 0.46	9.09 ± 0.31	8.59 ± 0.31
Relative (mg/g)	35.82 ± 0.53	35.83 ± 1.04	36.94 ± 0.55	36.23 ± 0.74
Six Months	10	10	10	10
Necropsy Body Wt. (g)	278.8 ± 4.0	277.3 ± 6.7	287.5 ± 6.4	275.6 ± 7.4
Kidneys				
Absolute (g)	1.71 ± 0.03**	1.67 ± 0.04	1.82 ± 0.05*	1.83 ± 0.03*
Relative (mg/g)	6.12 ± 0.04**	6.04 ± 0.08	6.32 ± 0.11	6.67 ± 0.17**
Liver				
Absolute (g)	8.66 ± 0.37	8.69 ± 0.35	9.09 ± 0.34	8.45 ± 0.27
Relative (mg/g)	31.01 ± 1.14	31.23 ± 0.64	31.59 ± 0.91	30.73 ± 0.92
Twelve Months	9	10	10	10
Necropsy Body Wt. (g)	313.9 ± 16.0	336.9 ± 15.7	310.0 ± 9.6	290.9 ± 10.5
Kidneys				
Absolute (g)	1.87 ± 0.07	1.95 ± 0.05	2.06 ± 0.06	1.93 ± 0.06
Relative (mg/g)	6.02 ± 0.21**	5.84 ± 0.16	6.68 ± 0.22*	6.67 ± 0.23*
Liver				
Absolute (g)	9.61 ± 0.44	10.81 ± 0.53	10.03 ± 0.49	8.73 ± 0.36
Relative (mg/g)	30.74 ± 0.84	32.40 ± 1.64	32.33 ± 1.12	30.06 ± 0.99
Eighteen Months	10	8	10	8
Necropsy Body Wt. (g)	359.5 ± 11.5*	372.9 ± 25.5	353.8 ± 11.5	313.6 ± 16.4
Kidneys				
Absolute (g)	2.20 ± 0.06	2.12 ± 0.05	2.12 ± 0.08	2.05 ± 0.06
Relative (mg/g)	6.16 ± 0.16	5.91 ± 0.49	6.02 ± 0.26	6.63 ± 0.28
Liver				
Absolute (g)	10.97 ± 0.52*	11.58 ± 0.64	10.54 ± 0.58	9.49 ± 0.50
Relative (mg/g)	30.67 ± 1.52	31.98 ± 2.75	29.83 ± 1.47	30.29 ± 0.62

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

6 ^aData presented as mean ± standard error.

7 ^bStatistical analysis performed by the Jonckheere (trend) and the Williams or Dunnett (pairwise) tests.

8 ^cNumber of animals examined at each time point.

9 ^dRelative organ weights (organ-weight-to-body-weight ratios) are given as mg organ weight/g body weight.

1 Plasma, kidney, and urine from up to 10 animals per exposure group were collected from interim
2 animals at 3, 6, 12, and 18 months. Total tungsten concentrations in all matrices were determined
3 using validated analytical methods (Appendix E). In male rats, plasma tungsten concentrations
4 increased proportionally with exposure concentration at all time points (except at 18 months for
5 the 500 mg/L group) with no observed differences between time points (Table 22; Figure 3A). In
6 female rats, plasma tungsten concentrations increased proportionally with exposure
7 concentration up to 500 mg/L at all time points (except at 18 months for the 250 mg/L group);
8 however, at 1,000 mg/L, the trend was toward a more-than-proportional increase in tungsten
9 concentrations with increasing exposure concentration (Table 23; Figure 3A). Low tungsten
10 concentrations were observed in some vehicle control groups; however, tungsten concentrations
11 in exposed groups were significantly higher than those in corresponding vehicle control groups.
12 There was no observed sex difference in plasma tungsten concentrations in rats (Table 22,
13 Table 23; Figure 3A).

14 In male rats, at 3 months, tungsten concentrations in the kidney increased proportionally with
15 exposure concentration (Table 22; Figure 3B); however, the trend was toward a less-than-
16 proportional increase in tungsten concentrations with increasing exposure concentration at 6, 12,
17 and 18 months in males (except at 18 months for the 1,000 mg/L group) and at all time points in
18 females (Table 22, Table 23; Figure 3B). Kidney tungsten concentrations increased with
19 increasing exposure duration and concentration in both males and females, with the
20 kidney/plasma ratio >1 (Table 22, Table 23). Taken collectively, these data demonstrate that
21 tungsten is retained in the kidney, and the retention increases with the exposure duration
22 (Figure 3B). Low tungsten concentrations were observed in some vehicle control groups;
23 however, tungsten concentrations in exposed groups were significantly higher than those in
24 corresponding vehicle control groups. There were no observed sex differences in kidney tungsten
25 concentrations in rats (Table 22, Table 23; Figure 3B).

26 Tungsten concentrations in urine are presented as both $\mu\text{g/mL}$ of urine and $\mu\text{g/mg}$ creatinine.
27 Creatinine-corrected tungsten concentrations in urine increased with exposure concentration in
28 both males and females (Figure 3C). The trend was toward a less-than-proportional increase in
29 tungsten concentration with increasing exposure concentration and an increase in tungsten
30 concentration with exposure duration, both of which were more evident in females than in males.
31 Depending on the exposure concentration and duration, females excreted 1.2- to 4-fold more
32 tungsten in urine compared to males (Table 22, Table 23). Low tungsten concentrations were
33 observed in some vehicle control groups; however, the tungsten concentrations in exposed
34 groups were significantly higher than those in the corresponding vehicle control groups.

1 **Table 22. Summary of Plasma, Kidney, and Urine Tungsten Concentration Data for Male Rats**
 2 **Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months^a**

	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
Three Months				
Kidney (µg/g)	BD	2.79 ± 0.24 (10)	5.30 ± 0.50 (10)	10.50 ± 2.94 (10)
Plasma (µg/mL) ^{b,c}	0.01 ± 0.00** (10)	1.34 ± 0.20** (10)	2.91 ± 0.27** (10)	5.87 ± 0.54** (10)
Kidney/Plasma Ratio ^d	BD	2.51 ± 0.42 (10)	1.87 ± 0.16 (10)	1.83 ± 0.51 (10)
Urine ^{b,c}				
Urine (µg/mL urine)	0.04 ± 0.01** (10)	135.36 ± 16.14** (9)	321.33 ± 21.53** (9)	862.80 ± 58.27** (10)
Urine (µg/mg creatinine)	0.02 ± 0.00** (10)	101.00 ± 12.63** (9)	268.52 ± 30.82** (9)	663.17 ± 41.98** (10)
Six Months				
Kidney (µg/g)	BD	6.77 ± 2.13 (10)	13.40 ± 1.86 (10)	14.11 ± 1.05 (10)
Plasma (µg/mL) ^{b,c}	0.02 ± 0.00** (10)	1.37 ± 0.14** (10)	4.89 ± 1.02** (10)	6.48 ± 0.78** (10)
Kidney/Plasma Ratio	BD	5.55 ± 2.04 (10)	3.39 ± 0.64 (10)	2.45 ± 0.33 (10)
Urine ^{b,c}				
Urine (µg/mL urine)	0.04 ± 0.00** (10)	143.37 ± 16.83** (10)	258.30 ± 27.17** (10)	572.10 ± 59.06** (10)
Urine (µg/mg creatinine)	0.03 ± 0.00** (10)	107.36 ± 10.22** (10)	216.91 ± 30.97** (10)	338.59 ± 41.61** (10)
Twelve Months				
Kidney (µg/g)	BD	14.84 ± 2.09 (10)	20.42 ± 3.85 (10)	27.98 ± 2.71 (10)
Plasma (µg/mL) ^{b,c}	0.01 ± 0.00** (8)	2.22 ± 0.25** (10)	3.85 ± 0.62** (10)	7.05 ± 0.55** (10)
Kidney/Plasma Ratio	BD	7.50 ± 1.09 (10)	6.22 ± 1.26 (10)	4.11 ± 0.45 (10)
Urine ^{b,c}				
Urine (µg/mL urine)	1.29 ± 0.95** (8)	161.92 ± 15.97** (10)	368.70 ± 52.38** (10)	654.23 ± 125.91** (10)
Urine (µg/mg creatinine)	2.05 ± 1.76** (8)	94.50 ± 14.54** (10)	284.48 ± 66.96** (10)	435.73 ± 65.39** (10)
Eighteen Months				
Kidney (µg/g) ^{b,c}	0.05 ± 0.01** (6)	25.74 ± 3.34** (9)	30.27 ± 5.21** (8)	74.90 ± 20.04** (9)
Plasma (µg/mL)	BD	2.64 ± 0.35 (9)	2.34 ± 0.95 (8)	7.59 ± 2.66 (9)
Kidney/Plasma Ratio	BD	12.63 ± 3.18 (9)	24.26 ± 8.14 (8)	16.69 ± 4.27 (9)
Urine ^{b,c}				
Urine (µg/mL urine)	0.03 ± 0.01** (9)	121.77 ± 9.17** (9)	159.93 ± 28.11** (8)	390.11 ± 38.03** (9)
Urine (µg/mg creatinine)	0.03 ± 0.01** (8)	104.37 ± 7.08** (9)	156.40 ± 27.17** (7)	436.76 ± 109.49** (9)

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 **Statistically significant at $p \leq 0.01$.

6 BD = below detection; group did not have more than 20% of its values above the limit of detection (LOD).

7 ^aData presented as mean ± standard error (n).

8 ^bStatistical analysis performed by the Jonckheere (trend) and Shirley or Dunn (pairwise) tests.

9 ^cValues below the LOD (0.013 µg/mL) were substituted with 1/2 the LOD value. If 80% or more of the values in the vehicle control group were below the LOD, no mean or standard error were calculated, and no statistical analysis was performed.

10 ^dFor the kidney/plasma ratio calculation, a plasma density of 1 g/mL was assumed.

1 **Table 23. Summary of Plasma, Kidney, and Urine Tungsten Concentration Data for Female Rats**
 2 **Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months^a**

	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
Three Months				
Kidney (µg/g)	BD	3.92 ± 0.24 (10)	6.39 ± 0.52 (10)	11.93 ± 0.71 (10)
Plasma (µg/mL)	BD	1.92 ± 0.13 (10)	3.67 ± 0.58 (10)	10.99 ± 1.40 (10)
Kidney/Plasma Ratio ^b	BD	2.06 ± 0.08 (10)	1.92 ± 0.18 (10)	1.17 ± 0.09 (10)
Urine ^{c,d}				
Urine (µg/mL urine)	0.04 ± 0.01** (10)	167.30 ± 9.11** (10)	355.50 ± 26.36** (8)	512.20 ± 43.84** (10)
Urine (µg/mg creatinine)	0.05 ± 0.01** (10)	247.12 ± 14.01** (10)	484.15 ± 56.46** (8)	809.98 ± 103.55** (10)
Six Months				
Kidney (µg/g)	BD	6.65 ± 0.92 (10)	9.89 ± 0.67 (10)	17.11 ± 1.81 (10)
Plasma (µg/mL) ^{c,d}	0.05 ± 0.01** (10)	2.41 ± 0.43** (10)	4.01 ± 0.39** (10)	10.44 ± 1.38** (10)
Kidney/Plasma Ratio	BD	2.96 ± 0.32 (10)	2.56 ± 0.13 (10)	1.70 ± 0.11 (10)
Urine ^{c,d}				
Urine (µg/mL urine)	0.04 ± 0.01** (10)	174.10 ± 10.74** (10)	413.30 ± 20.05** (10)	619.20 ± 50.46** (10)
Urine (µg/mg creatinine)	0.05 ± 0.00** (10)	228.49 ± 20.07** (10)	467.46 ± 40.93** (10)	856.85 ± 58.29** (10)
Twelve Months				
Kidney (µg/g)	BD	10.62 ± 0.53 (10)	19.26 ± 1.89 (10)	31.65 ± 2.20 (10)
Plasma (µg/mL) ^{c,d}	0.03 ± 0.01** (8) ^e	2.47 ± 0.29** (10)	4.55 ± 0.33** (10)	12.06 ± 1.31** (10)
Kidney/Plasma Ratio	4.57 ± 0.85** (4)	4.76 ± 0.50 (10)	4.36 ± 0.42 (10)	2.96 ± 0.48 (10)
Urine ^{c,d}				
Urine (µg/mL urine)	0.07 ± 0.02** (9)	158.00 ± 8.45** (10)	291.60 ± 17.88** (10)	672.30 ± 51.25** (10)
Urine (µg/mg creatinine)	0.10 ± 0.02** (8)	298.03 ± 26.20** (10)	567.24 ± 42.20** (10)	1,036.16 ± 57.95** (10)
Eighteen Months				
Kidney (µg/g) ^{c,d}	0.09 ± 0.01** (3)	25.89 ± 7.69* (8)	27.96 ± 2.79** (10)	43.29 ± 2.90** (8)
Plasma (µg/mL) ^{c,d}	0.02 ± 0.00** (2)	5.45 ± 3.00* (8)	5.71 ± 0.59* (10)	11.78 ± 1.51** (8)
Kidney/Plasma Ratio	NR ^f	7.11 ± 1.17 (8)	5.47 ± 0.79 (10)	3.99 ± 0.42 (8)
Urine ^{c,d}				
Urine (µg/mL urine)	0.03 ± 0.00** (8)	162.00 ± 17.50** (8)	328.11 ± 37.85** (9)	612.38 ± 47.87** (8)
Urine (µg/mg creatinine)	0.05 ± 0.01** (8)	349.42 ± 34.89** (8)	628.91 ± 105.62** (9)	1,089.91 ± 100.25** (8)

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

6 BD = below detection; group did not have more than 20% of its values above the limit of detection (LOD).

7 ^aData presented as mean ± standard error (n).


8 ^bFor the kidney/plasma ratio calculation, a plasma density of 1 g/mL was assumed.

9 ^cStatistical analysis performed by the Jonckheere (trend) and Shirley or Dunn (pairwise) tests.

10 ^dValues below the LOD (0.013 µg/mL) were substituted with 1/2 the LOD value. If 80% or more of the values in the vehicle
 11 control group were below the LOD, no mean or standard error were calculated, and no statistical analysis was performed.

12 ^eThe plasma concentration value for one female in the 0 mg/L group at 12 months was excluded from the analysis as an
 13 implausible value.

14 ^fThe kidney/plasma ratio could not be calculated for females in the 0 mg/L group at 18 months because no animals in the group
 15 had concentration measures for both kidney and plasma.



1
2 **Figure 3. Tungsten Concentrations in Plasma, Kidney, and Urine in Rats Exposed to Sodium**
3 **Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months**

4 Study day (SD) 93 = 3 months; SD 185 = 9 months; SD 367 = 12 months; SD 549 = 18 months.

1 Two-year Study (Postweaning Phase)

2 Survival

3 Survival to study termination was significantly increased in the groups of exposed male rats
 4 compared to the vehicle control males (Table 24; Figure 4). The survival of the vehicle control
 5 males was lower than that typically seen in groups of control male Sprague Dawley
 6 (Hsd:Sprague Dawley[®] SD[®]) rats in previous 2-year NTP studies. Early deaths in the vehicle
 7 control males were attributed at necropsy primarily to CPN (22 animals). Pituitary gland
 8 adenomas were listed as the cause of death at necropsy for six vehicle control males, compared
 9 to two, two, and one animal in the 250, 500, and 1,000 mg/L males, respectively. The large
 10 number of vehicle control males with CPN as a cause of death at necropsy corresponded to an
 11 increased severity of nephropathy observed histologically in that group. There were no
 12 exposure-related differences in the survival of female groups.

13 **Table 24. Summary of Survival of Male and Female Rats in the Perinatal and Two-year Drinking**
 14 **Water Study of Sodium Tungstate Dihydrate**

	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
Male				
Animals Initially in Study	50	50	50	50
Moribund	17	14	10	8
Natural Deaths	21	10	16	13
Animals Surviving to Study Termination	12	26	24 ^a	29
Percent Probability of Survival at End of Study ^b	24.0%	52.0%	48.0%	58.0%
Mean Survival (Days) ^c	620.7 ± 17.4	656.6 ± 20.3	670.6 ± 13.7	657.4 ± 17.6
Survival Analysis ^d	p = 0.008N	p = 0.028N	p = 0.018N	p = 0.004N
Female				
Animals Initially in Study	50	50	50	50
Moribund	13	10	13	9
Natural Deaths	7	7	6	10
Animals Surviving to Study Termination	30	33	31	31 ^e
Percent Probability of Survival at End of Study	60.0%	66.0%	62.0%	62.0%
Mean Survival (Days)	667.0 ± 15.1	686.7 ± 12.3	661.8 ± 18.0	674.7 ± 14.8
Survival Analysis	p = 0.976N	p = 0.467N	p = 0.964	p = 0.794


15 ^aIncludes one animal that died naturally during the last week of the study.

16 ^bKaplan-Meier determinations.


17 ^cMean of litter means of all deaths (uncensored, censored, and study termination) ± standard error.

18 ^dThe result of the Cox proportional hazards trend test with random litter effects is in the vehicle control group column, and the
 19 results of the proportional hazards pairwise comparisons to the vehicle control group with random litter effects are in the exposed
 20 group columns. A negative trend or lower mortality in an exposure group is indicated by N.

21 ^eIncludes one animal that was euthanized moribund during the last week of the study.



1



2

3

4

Figure 4. Kaplan-Meier Survival Curves for Rats Exposed to Sodium Tungstate Dihydrate in Drinking Water for Two Years

1 **Body Weights, Water and Compound Consumption, and Clinical Observations**

2 At study termination, mean body weights of all groups of exposed males were within 10% of the
3 vehicle control group (Table 25; Figure 5). In females, mean body weights of the 500 mg/L and
4 1,000 mg/L groups at study termination were 88.9% and 78.4% of the vehicle control group,
5 respectively (Table 26; Figure 5).

6 Over the course of the chronic study, the respective mean water consumption for the 250, 500,
7 and 1,000 mg/L groups averaged 93%, 99%, and 84% of the vehicle control males and averaged
8 95%, 100%, and 91% of the vehicle control females (Table 27, Table 28). The daily ST
9 consumption averaged 14.2, 30.4, and 54.5 mg/kg/day for males, and 18.2, 39.3, and
10 74.3 mg/kg/day for females, respectively (Appendix G). Consumption at weeks 1, 13, 54, and
11 102 is presented in Table 27 and Table 28. Overall, the consumed dose of ST was of similar
12 proportionality (ranging from 1.9- to 2.1-fold increase from one dose to the next) to the increase
13 in ST concentration in drinking water (2-fold increases from one concentration to the next). Dose
14 proportionality was consistent for both male and female rats (Table 27, Table 28). No clinical
15 observations were considered exposure related (Appendix G).

1 **Table 25. Summary of Survival and Mean Body Weights of Male Rats in the Perinatal and**
 2 **Two-year Drinking Water Study of Sodium Tungstate Dihydrate**

Study Day ^a	0 mg/L		250 mg/L		500 mg/L			1,000 mg/L			
	Av. Wt. (g) ^a	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters
1	53.2	25	52.2	98.2	25	52.6	98.9	25	52.6	98.9	25
8	82.2	25	81.1	98.6	25	82.3	100.1	25	80.7	98.2	25
15	127.5	25	125.7	98.6	25	127.4	100.0	25	118.0	92.5	25
22	178.3	25	174.4	97.8	25	177.1	99.3	25	156.0	87.5	25
29	226.9	25	222.1	97.9	25	225.2	99.2	25	194.0	85.5	25
36	271.3	25	267.6	98.6	25	269.1	99.2	25	235.4	86.8	25
43	304.9	25	302.9	99.3	25	301.3	98.8	25	270.5	88.7	25
50	326.8	25	324.7	99.3	25	325.7	99.7	25	299.4	91.6	25
57	352.9	25	349.8	99.1	25	350.5	99.3	25	325.6	92.3	25
64	371.4	25	363.8	97.9	25	366.8	98.8	25	345.9	93.1	25
71	386.9	25	380.7	98.4	25	378.1	97.7	25	357.1	92.3	25
78	399.7	25	394.7	98.7	25	392.3	98.1	25	374.5	93.7	25
85	411.8	25	405.7	98.5	25	402.3	97.7	25	385.6	93.6	25
92	419.1	25	415.4	99.1	25	408.9	97.6	25	395.5	94.4	25
120	443.5	25	431.7	97.4	25	429.7	96.9	25	420.0	94.7	25
148	466.5	25	459.1	98.4	25	448.9	96.2	25	442.1	94.8	25
176	489.9	25	478.5	97.7	25	471.6	96.3	25	460.6	94.0	25
204	505.2	25	494.7	97.9	25	487.0	96.4	25	476.7	94.4	25
232	521.0	25	508.6	97.6	25	500.1	96.0	25	491.5	94.3	25
260	537.7	25	523.7	97.4	25	517.0	96.1	25	505.3	94.0	25
288	549.3	25	533.1	97.0	25	528.4	96.2	25	516.2	94.0	25
316	560.4	25	547.1	97.6	25	540.4	96.4	25	527.5	94.1	25
344	573.4	25	555.6	96.9	25	553.2	96.5	25	537.9	93.8	25
372	585.5	25	570.5	97.4	25	561.7	95.9	25	547.6	93.5	25
400	593.9	25	580.4	97.7	25	572.1	96.3	25	555.5	93.5	25
428	598.3	25	589.3	98.5	25	579.9	96.9	25	561.8	93.9	25
456	601.3	25	589.9	98.1	25	584.7	97.2	25	566.2	94.2	25
484	603.4	25	595.5	98.7	25	589.1	97.6	25	568.4	94.2	25
512	606.9	25	597.1	98.4	25	589.5	97.1	25	566.7	93.4	25
540	602.0	25	596.8	99.1	25	587.0	97.5	25	564.0	93.7	25
568	595.3	25	594.7	99.9	23	578.2	97.1	25	558.6	93.8	25
596	593.0	23	588.5	99.2	22	576.8	97.3	24	552.7	93.2	25
624	586.2	21	589.2	100.5	21	566.4	96.6	24	557.9	95.2	24
652	580.7	18	572.2	98.5	21	576.3	99.3	22	551.3	94.9	23
680	582.7	12	551.4	94.6	19	551.0	94.6	22	545.5	93.6	23
708	601.8	9	559.8	93.0	15	550.9	91.5	19	548.3	91.1	21
EOS	587.2	12	545.7	92.9	15	551.8	94.0	16	545.1	92.8	21

3 EOS = end of study; No. of litters = number of litters represented in weight average.

4 ^aStudy day 1 is the day animals were placed on study after pups were weaned.

5 ^bAverage weights shown are means of litter means.


1 **Table 26. Summary of Survival and Mean Body Weights of Female Rats in the Perinatal and**
 2 **Two-year Drinking Water Study of Sodium Tungstate Dihydrate**

Study Day ^a	0 mg/L		250 mg/L		500 mg/L		1,000 mg/L				
	Av. Wt. (g) ^b	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters	Av. Wt. (g)	Wt. (% of Controls)	No. of Litters
1	51.4	25	49.7	96.6	25	49.8	96.9	25	52.1	101.2	25
8	78.6	25	76.9	97.9	25	77.6	98.7	25	76.0	96.7	25
15	113.7	25	109.1	95.9	25	110.6	97.2	25	100.6	88.5	25
22	139.1	25	136.0	97.8	25	135.6	97.5	25	122.9	88.3	25
29	166.2	25	162.9	98.0	25	163.3	98.3	25	144.9	87.2	25
36	180.9	25	181.7	100.5	25	181.3	100.3	25	164.5	90.9	25
43	198.4	25	199.3	100.4	25	194.8	98.2	25	185.6	93.5	25
50	211.8	25	212.5	100.3	25	207.6	98.0	25	201.8	95.3	25
57	222.7	25	225.0	101.1	25	220.3	99.0	25	215.0	96.6	25
64	227.2	25	224.8	99.0	25	223.1	98.2	25	215.1	94.7	25
71	233.4	25	229.9	98.5	25	227.5	97.5	25	226.5	97.0	25
78	241.6	25	238.9	98.9	25	239.7	99.2	25	235.8	97.6	25
85	247.4	25	245.0	99.0	25	243.5	98.4	25	241.0	97.4	25
92	254.4	25	249.5	98.1	25	249.1	97.9	25	246.5	96.9	25
120	266.2	25	259.3	97.4	25	261.2	98.1	25	258.9	97.3	25
148	275.0	25	273.4	99.4	25	269.7	98.0	25	266.4	96.9	25
176	286.4	25	283.1	98.9	25	278.6	97.3	25	275.7	96.3	25
204	290.4	25	289.3	99.6	25	280.2	96.5	25	284.4	97.9	25
232	295.5	25	297.6	100.7	25	289.2	97.9	25	291.9	98.8	25
260	303.1	25	302.8	99.9	25	293.0	96.7	25	297.7	98.2	25
288	307.0	25	311.6	101.5	25	299.8	97.7	25	301.7	98.3	25
316	317.7	25	320.1	100.7	25	305.5	96.1	25	307.1	96.6	25
344	324.4	25	324.8	100.1	25	309.8	95.5	25	312.8	96.4	25
372	332.0	25	329.6	99.3	25	314.9	94.8	25	316.9	95.4	25
400	343.4	25	338.1	98.5	25	321.6	93.6	25	323.3	94.2	25
428	348.4	25	342.8	98.4	25	325.4	93.4	25	325.7	93.5	25
456	354.2	25	353.0	99.7	25	337.2	95.2	24	327.5	92.5	25
484	364.9	25	359.6	98.6	25	345.8	94.8	24	331.8	90.9	25
512	365.5	25	363.0	99.3	25	355.2	97.2	24	333.2	91.2	25
540	381.5	25	371.1	97.3	25	365.8	95.9	24	342.6	89.8	25
568	377.9	24	377.3	99.8	25	376.4	99.6	24	339.0	89.7	25
596	385.4	24	384.3	99.7	25	363.6	94.3	22	345.3	89.6	25
624	393.9	24	388.1	98.5	25	370.7	94.1	22	342.5	86.9	24
652	392.7	24	390.5	99.4	25	381.9	97.3	21	335.4	85.4	23
680	384.1	24	381.9	99.4	25	366.2	95.3	21	330.1	85.9	23
708	386.1	22	384.4	99.6	24	372.8	96.5	21	326.9	84.7	22
EOS	414.9	20	399.2	96.2	23	368.9	88.9	20	325.5	78.4	20


3 EOS = end of study; No. of litters = number of litters represented in weight average.

4 ^aStudy day 1 is the day animals were placed on study after pups were weaned.

5 ^bAverage weights shown are means of litter means.



1



2

3

4

Figure 5. Growth Curves for Rats Exposed to Sodium Tungstate Dihydrate in Drinking Water for Two Years

1 **Table 27. Summary of Water and Sodium Tungstate Dihydrate Consumption of Male Rats in the**
 2 **Perinatal and Two-year Drinking Water Study**

Week	0 mg/L		250 mg/L		500 mg/L		1,000 mg/L	
	Water (g/day) ^a	Water (g/day)	Dose (mg/kg/day) ^b	Water (g/day)	Dose (mg/kg/day)	Water (g/day)	Dose (mg/kg/day)	
1	8.9	8.8	42.1	8.3	78.9	9.6	182.5	
13	24.9	23.4	14.4	24.8	30.8	23.3	60.3	
54	26.5	24.9	10.9	25.4	22.6	22.9	41.7	
102	42.4	35.9	16.0	42.2	38.0	27.7	49.8	

3 ^aGrams of water consumed/animal/day.

4 ^bMilligrams of sodium tungstate dihydrate consumed/kilogram body weight/day.

5 **Table 28. Summary of Water and Sodium Tungstate Dihydrate Consumption of Female Rats in the**
 6 **Perinatal and Two-year Drinking Water Study of Sodium Tungstate Dihydrate**

Week	0 mg/L		250 mg/L		500 mg/L		1,000 mg/L	
	Water (g/day) ^a	Water (g/day)	Dose (mg/kg/day) ^b	Water (g/day)	Dose (mg/kg/day)	Water (g/day)	Dose (mg/kg/day)	
1	10.2	9.9	49.8	10.3	103.3	9.9	190.1	
13	20.0	18.2	18.6	18.9	38.8	17.6	73.0	
54	22.7	21.0	15.8	22.3	35.5	19.9	62.7	
102	27.0	24.7	16.1	29.0	39.6	28.1	87.4	

7 ^aGrams of water consumed per animal/day.

8 ^bMilligrams of sodium tungstate dihydrate consumed/kilogram body weight/day.

9 **Histopathology**

10 This section describes the significant or biologically noteworthy changes in the incidence of
 11 neoplasms and nonneoplastic lesions of the thyroid gland, kidney, and uterus. Summaries of the
 12 incidences of neoplasms and nonneoplastic lesions and statistical analyses of primary neoplasms
 13 are presented in CEBS (Appendix G).

14 *Thyroid Gland:* There was a significant increase in the incidence of thyroid gland C-cell
 15 adenomas in the 500 mg/L females relative to the vehicle control group (Table 29). There was no
 16 significant increase in C-cell carcinomas, although the 1,000 mg/L females had a slightly higher
 17 incidence than the other exposure groups (Table 29). There were no significant differences in the
 18 incidences of C-cell adenoma or carcinoma (combined) in any exposed group compared to the
 19 vehicle control group; however, the incidences in the 250 and 500 mg/L groups were outside of
 20 the historical control range (Table 29). The incidences of C-cell hyperplasia were not
 21 significantly increased in any exposed group when compared to the vehicle control group
 22 (Table 29). C-cell adenomas consisted of a discrete proliferation of C-cells that was larger than
 23 five follicles in diameter and caused compression of surrounding follicles (Figure 6). C-cell
 24 carcinomas tended to be large neoplasms that replaced the normal architecture of the thyroid
 25 gland and were diagnosed when there was evidence of invasion into the thyroid gland capsule or
 26 surrounding tissue (Figure 7).

1 **Table 29. Incidences of Neoplastic and Nonneoplastic Lesions of the Thyroid Gland in Female Rats**
 2 **in the Perinatal and Two-year Drinking Water Study of Sodium Tungstate Dihydrate**

	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
n^a	50	50	49	50
C-cell, Hyperplasia ^b	14 (2.5) ^c	13 (1.8)	9 (2.0)	12 (1.8)
C-cell Adenoma ^d				
Overall rate ^e	5/50 (10%)	13/50 (26%)	13/49 (27%)	8/50 (16%)
Rate per litters ^f	4/25 (16%)	12/25 (48%)	10/25 (40%)	8/25 (32%)
Adjusted rate ^g	12.1%	29.7%	31.9%	18.5%
Terminal rate ^h	2/30 (7%)	12/33 (36%)	8/31 (26%)	5/31 (16%)
First incidence (days)	676	701	585	583
Rao-Scott-adjusted Poly-3 test ⁱ	p = 0.451	p = 0.057	p = 0.040	p = 0.321
C-cell Carcinoma ^j				
Overall rate	2/50 (4%)	2/50 (4%)	2/49 (4%)	4/50 (8%)
Rate per litters	2/25 (8%)	2/25 (8%)	2/25 (8%)	4/25 (16%)
Adjusted rate	4.9%	4.6%	5%	9.5%
Terminal rate	1/30 (3%)	1/33 (3%)	1/31 (3%)	4/31 (13%)
First incidence (days)	708	630	716	730 (T)
Rao-Scott-adjusted Poly-3 test	p = 0.228	p = 0.651N	p = 0.659	p = 0.344
C-cell Adenoma or Carcinoma (Combined) ^k				
Overall rate	7/50 (14%)	15/50 (30%)	14/49 (29%)	11/50 (22%)
Rate per litters	6/25 (24%)	13/25 (52%)	11/25 (44%)	11/25 (44%)
Adjusted rate	16.9%	34%	34.4%	25.4%
Terminal rate	3/30 (10%)	13/33 (39%)	9/31 (29%)	8/31 (26%)
First incidence (days)	676	630	585	583
Rao-Scott-adjusted Poly-3 test	p = 0.369	p = 0.073	p = 0.072	p = 0.260

3 ^aNumber of animals with tissue examined microscopically.

4 ^bNumber of animals with lesion.

5 ^cAverage severity grade of lesions in affected animals: 1 = minimal, 2 = mild, 3 = moderate, 4 = marked.

6 ^dHistorical control incidence for all routes of 2-year studies (mean ± standard deviation): 70/488 (15.05% ± 7.65%);
 7 range: 4% to 24%.

8 ^eNumber of animals with neoplasm/number of animals necropsied.

9 ^fNumber of litters with neoplasm-bearing animals/number of litters examined at site.


10 ^gPoly-3 estimated neoplasm incidence after adjustment for intercurrent mortality.

11 ^hObserved incidence at terminal euthanasia.

12 ⁱBeneath the control incidence is the p value associated with the trend test. Beneath the exposed group incidences are the p values
 13 corresponding to pairwise comparisons between the vehicle control group and that exposed group. The Rao-Scott test adjusts the
 14 Poly-3 test (which accounts for differential mortality in animals that do not reach terminal euthanasia) for within-litter
 15 correlation. A negative trend or a lower incidence in an exposure group is indicated by N.

16 ^jHistorical control incidence: 7/488 (1.56% ± 1.67%); range: 0% to 4%.


17 ^kHistorical control incidence: 76/488 (16.38% ± 8.21%); range: 4% to 28%.




1
2 **Figure 6. C-cell Adenoma in the Thyroid Gland of a Female Sprague Dawley Rat Exposed to**
3 **1,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E)**

4 A) The C-cell adenoma is a mass of several nests, or clusters, of cells separated by a delicate fibrovascular stroma. B) Higher
5 magnification of the C-cell adenoma in panel A; at this magnification, the fibrovascular stroma that separates the nests of C-cells
6 is more apparent (arrows). The cells that make up the adenoma have oval nuclei and pale eosinophilic cytoplasm. A few follicles
7 are entrapped (asterisks).

A)



B)



1
2 **Figure 7. C-cell Carcinoma in the Thyroid Gland of a Female Sprague Dawley Rat Exposed to**
3 **1,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E)**

4 A) The C-cell carcinoma completely obliterates the normal architecture of the thyroid gland. B) Higher magnification of the C-
5 cell carcinoma in panel A; the cells tend to have slightly less cytoplasm than those of the C-cell adenoma (Figure 6), but the main
6 feature of C-cell carcinomas is invasion through the capsule (arrows).

7 *Kidney:* The incidences of suppurative inflammation of the renal tubules were significantly
8 increased in the 1,000 mg/L males and females, and the incidence of renal tubule regeneration
9 was significantly increased in the 1,000 mg/L females, relative to the respective vehicle control
10 groups (Table 30). Renal tubule suppurative inflammation consisted of dilated renal tubules
11 filled with neutrophils and necrotic debris (Figure 8). One to five affected renal tubules in a
12 section of kidney was graded as minimal; 6 to 15 affected tubules in a section of kidney was
13 graded as mild; and more than 15 affected tubules in a section of kidney was graded as moderate.
14 Renal tubule regeneration was characterized by cytoplasmic basophilia, karyomegaly,
15 hypertrophy and hyperplasia of the renal tubule epithelium (Table 30; Figure 9). Occasional
16 mitotic figures were also present. Severity grading was based on the amount of renal cortex
17 involved, with minimal regeneration involving <10% of the cortex; mild regeneration involving
18 approximately 10–25% of the cortex; moderate or marked regeneration was not observed. In
19 male rats, the widespread CPN made it impossible to identify regeneration as a distinct lesion;

1 hence, the incidence of renal tubule regeneration recorded might not reflect the actual number of
 2 animals with the lesion (Table 30).
 3 CPN was recorded in almost every male rat in the study; however, the mean severity score was
 4 the highest in the vehicle control group, and lowest in the 1,000 mg/L males (Table 30;
 5 Figure 10). Although no statistical comparison was conducted on the mean severity scores, the
 6 scores did parallel the incidences of several other lesions that are considered secondary to kidney
 7 failure associated with CPN. These lesions, which occurred at lower incidences in higher
 8 exposure concentration groups than in the vehicle control group, include diffuse hyperplasia of
 9 the parathyroid and mineral of the kidney, blood vessel, heart, large intestine (cecum), and
 10 glandular stomach (Appendix G).

11 **Table 30. Incidences of Nonneoplastic Lesions of the Kidney in Male and Female Rats in the**
 12 **Perinatal and Two-year Drinking Water Study of Sodium Tungstate Dihydrate**

	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
n^a	50	50	50	50
Male				
Renal Tubule, Regeneration ^b	0	1 (2.0) ^c	0	0
Nephropathy, Chronic Progressive	50 (3.4)	50 (3.2)	50 (3.2)	49 (2.5)
Renal Tubule, Inflammation, Suppurative	25** (1.2)	33 (1.3)	35 (1.3)	41** (1.6)
Female				
Renal Tubule, Regeneration	0**	0	0	18** (1.8)
Nephropathy, Chronic Progressive	49 (1.8)	49 (1.4)	48 (1.4)	47 (1.7)
Renal Tubule, Inflammation, Suppurative	8** (1.0)	9 (1.0)	6 (1.0)	19* (1.1)

13 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

14 Statistical significance for the vehicle control group indicates a significant trend test.


15 *Statistically significant at $p \leq 0.05$ by the Rao-Scott test; ** $p \leq 0.01$.

16 ^aNumber of animals with tissue examined microscopically.


17 ^bNumber of animals with lesion.

18 ^cAverage severity grade of lesions in affected animals: 1 = minimal, 2 = mild, 3 = moderate, 4 = marked.

A)



B)




1

2 **Figure 8. Suppurative Inflammation in the Renal Tubules of the Kidney from a Male Sprague**
3 **Dawley Rat Exposed to 1,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E)**


4

5 A) Numerous dilated tubules in the kidney are filled with neutrophils and necrotic debris (arrows). B) Higher magnification of
6 the renal tubule suppurative inflammation shown in panel A; the tubules are filled with neutrophils and cell debris, and there is a mixed inflammatory response in the interstitium.

A)




B)



1

2 **Figure 9. Renal Tubule Regeneration in the Kidney of a Female Sprague Dawley Rat Exposed to**
3 **1,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E)**

4 A) At this magnification, the regenerative tubules are evidenced by basophilic tubules (arrows). B) Higher magnification of the
5 renal tubule regeneration shown in panel A in which the affected tubules (asterisks) are basophilic, and there is slight crowding of
6 the nuclei compared with the cells in the normal, more eosinophilic tubules (arrowheads). No thickening of the basement
7 membrane is observed, as is seen in chronic progressive nephropathy; compare these tubules with those seen in Figure 10.



1
2 **Figure 10. Chronic Progressive Nephropathy in the Kidney of a Female Sprague Dawley Rat**
3 **Exposed to 1,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E)**

4 Compared with renal tubule regeneration (Figure 9), thickened basement membranes are visible surrounding the renal tubules
5 (arrows).

6 *Uterus:* A significant increase in the incidence of atypical hyperplasia of the uterus, relative to
7 the vehicle control group, occurred in the 500 mg/L females (Table 31). Atypical hyperplasia is
8 considered a preneoplastic lesion,⁸⁸ which can progress to uterine adenocarcinoma.

9 Atypical hyperplasia involved both the endometrial glands and the surface epithelium
10 (Figure 11). Affected endometrial glands were characterized by clusters of enlarged glands lined
11 by disorganized, large epithelial cells that were crowded and piled up on each other to form
12 multiple layers; this often resulted in blebbing (small protrusions of cells in the glandular lumen).
13 The involved cells displayed pleomorphism and anisokaryosis, and an occasional mitotic figure
14 was present.

15 Atypical hyperplasia affecting the surface epithelium consisted of branching, frond-like
16 projections of epithelial cells on a fibrovascular stalk that extended into the uterine lumen. These
17 thickened papillary projections were composed of cells that often contained clear vacuoles.

18 The severity (minimal, mild, moderate, marked) of atypical hyperplasia was graded based on the
19 amount of uterus that was involved in the lesion, as well as on the extent of the proliferation and
20 cellular atypia. Minimal lesions were small and focal, confined to one small area of the uterus,
21 and involved only one or a few glands or a small area of surface endometrium. Although the
22 cells might display some atypical characteristics (such as having differently sized nuclei or
23 abundant cytoplasm), the cells were not so anaplastic that they resembled those of an
24 adenocarcinoma. Blebbing of the cytoplasm was infrequent, and although the cells might have
25 been crowded, they did not typically form multiple layers. Increasing severity grades reflected
26 the involvement of a larger area of the uterus and a greater degree of cellular pleomorphism,
27 atypia, and proliferation, resulting in frequently vacuolated cells lining frond-like extensions
28 from the surface endometrium and crowded glands lined by multiple layers of abnormal looking
29 epithelial cells.

30 Often it is difficult to distinguish marked atypical hyperplasia from adenocarcinoma because
31 cellular features are similar with both lesions. Other than metastases, which only occur with

1 adenocarcinoma and not atypical hyperplasia, the main feature used to differentiate
 2 adenocarcinoma is invasion into the myometrium (Figure 11A). Invasion can be difficult to
 3 determine (its detection depends on the plane of the section taken through the uterus), and
 4 adenocarcinomas must be distinguished from adenomyosis, a condition in which nonneoplastic
 5 glands are present within the myometrium. Other features that can be observed in
 6 adenocarcinomas but are not seen with atypical hyperplasia include large areas of necrosis and
 7 hemorrhage, and scirrhous reactions; these features are more frequently present in
 8 adenocarcinomas that invade through the myometrium and serosa. In this study, the incidences
 9 of adenocarcinoma and adenoma were not increased in exposed groups compared to the vehicle
 10 control group (Table 31).

11 **Table 31. Incidences of Neoplastic and Nonneoplastic Lesions of the Uterus in Female Rats in the**
 12 **Perinatal and Two-year Drinking Water Study of Sodium Tungstate Dihydrate**

	0 mg/L	250 mg/L	500 mg/L	1,000 mg/L
n^a	50	50	50	50
Hyperplasia, Atypical ^b	4 (2.3) ^c	7 (1.4)	19** (1.7)	8 (2.3)
Adenocarcinoma ^d	3	0	2	5
Adenoma ^e	0	0	0	1

13 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

14 **Statistically significant at $p \leq 0.01$ by the Rao-Scott test.

15 ^aNumber of animals with tissue examined microscopically.


16 ^bNumber of animals with lesion.

17 ^cAverage severity grade of lesions in affected animals: 1 = minimal, 2 = mild, 3 = moderate, 4 = marked.


18 ^dHistorical control incidence for all routes of 2-year studies (mean \pm standard deviation): 20/350 (5.71% \pm 3.35%);
 19 range: 2% to 10%.

20 ^eHistorical control incidence: 1/350 (0.29% \pm 0.76%); range: 0% to 2%.


A)



B)



C)



1
2 **Figure 11. Atypical Hyperplasia in the Uterus of a Female Sprague Dawley Rat Exposed to**
3 **500 mg/L Sodium Tungstate Dihydrate for Two Years (H&E)**

4 A) A low magnification photomicrograph of atypical hyperplasia of the uterus shows proliferation of the surface and glandular
5 epithelium; the proliferative epithelium does not extend past the thick endometrial stroma into the underlying myometrial layers.
6 This lack of invasion is consistent with atypical hyperplasia; invasion into the myometrium would be indicative of an
7 adenocarcinoma. Even at this relatively low magnification, it is possible to notice the vacuoles in many of the epithelial cells; this
8 is a common feature of atypical hyperplasia. B) A higher magnification of the atypical hyperplasia seen in panel A in which the
9 vacuoles within the cytoplasm of numerous cells are easier to see. C) A photomicrograph of atypical hyperplasia affecting an
10 endometrial gland with large epithelial cells containing large nuclei, which are piling up on each other, resulting in protrusions
11 into the glandular lumen. The large number of degenerate neutrophils within the lumen are not a part of the atypical hyperplasia.

1 *Other Tissues:* In males, there was a significantly increased incidence of focal adrenal cortical
2 hyperplasia in the 1,000 mg/L group, relative to the vehicle control group, when unilateral and
3 bilateral lesions were combined (Appendix G). In the mandibular lymph node, there were
4 significantly increased incidences of plasma cell hyperplasia in the 500 and 1,000 mg/L males,
5 and of sinus dilation in the 250 mg/L males, relative to the vehicle control group (Appendix G).
6 In the nose, goblet cell hyperplasia of the nasopharyngeal duct was significantly increased in
7 incidence in the 250 mg/L males, and hyaline droplet accumulation in the respiratory epithelium
8 was significantly increased in incidence in all exposed groups of females, relative to the vehicle
9 control groups (Appendix G). The biological significance of these lesions is unknown.

10 In females, there was a significantly decreased incidence of mammary gland fibroadenomas in
11 the 1,000 mg/L group compared to the vehicle control group (Appendix G).

1 **Mice**

2 **Three-month Study**

3 There were no early deaths or exposure-related clinical observations in male or female
4 B6C3F1/N mice exposed to ST for 3 months (Table 32, Table 33; Appendix G). Over the course
5 of the study, group mean body weights were below 90% of the vehicle control group mean for
6 the 250, 1,000, and 2,000 mg/L females and the 2,000 mg/L males (Table 32, Table 33;
7 Figure 12). At study termination, the mean body weights of all exposed groups of males and
8 females were within 10% of the vehicle control groups.

9 Weekly mean water consumption was reduced slightly in the 1,000 mg/L male group (11%), the
10 2,000 mg/L males (16%), and the 2,000 mg/L females (11%), relative to the respective vehicle
11 control groups (Table 34; Appendix G). Drinking water concentrations of 125, 250, 500, 1,000,
12 and 2,000 mg/L resulted in average daily ST doses of approximately 14, 27, 57, 108, and
13 212 mg/kg/day for males and 14, 29, 58, 113, and 202 mg/kg/day for females.

1 **Table 32. Summary of Survival and Mean Body Weights of Male Mice in the Three-month Drinking Water Study of Sodium Tungstate**
 2 **Dihydrate**

Study Day ^a	0 mg/L			125 mg/L			250 mg/L			500 mg/L			1,000 mg/L			2,000 mg/L		
	Av. Wt. (g)	N		Av. Wt. (g)	Wt. (% of Controls)	N	Av. Wt. (g)	Wt. (% of Controls)	N	Av. Wt. (g)	Wt. (% of Controls)	N	Av. Wt. (g)	Wt. (% of Controls)	N	Av. Wt. (g)	Wt. (% of Controls)	N
1	22.2	10		22.4	100.9	10	22.6	101.7	10	22.7	102.3	10	22.1	99.8	10	22.4	100.9	10
8	24.4	10		24.8	101.5	10	24.6	100.7	10	24.3	99.6	10	23.9	98.0	10	23.9	97.7	10
15	25.9	10		26.3	101.7	10	26.0	100.5	10	25.7	99.5	10	25.3	97.6	10	24.8	95.7	10
22	27.9	10		28.7	102.7	10	28.3	101.5	10	27.9	99.9	10	27.1	97.2	10	26.5	95.0	10
29	28.9	10		29.6	102.4	10	29.4	101.7	10	29.0	100.3	10	28.4	98.1	10	27.3	94.4	10
36	30.4	10		31.4	103.3	10	31.0	102.0	10	30.3	99.8	10	29.7	97.7	10	28.4	93.3	10
43	32.7	10		33.5	102.7	10	33.0	101.0	10	32.4	99.2	10	31.6	96.8	10	30.0	91.9	10
50	34.6	10		35.6	103.0	10	34.7	100.4	10	34.0	98.3	10	33.9	98.1	10	31.8	92.1	10
57	35.8	10		37.3	104.2	10	36.5	101.9	10	35.3	98.7	10	35.3	98.7	10	32.8	91.5	10
64	36.2	10		38.0	105.0	10	37.0	102.3	10	35.9	99.1	10	35.9	99.4	10	32.9	91.0	10
71	37.6	10		39.3	104.5	10	38.1	101.3	10	37.2	98.8	10	36.9	98.2	10	34.0	90.4	10
78	39.0	10		40.7	104.6	10	39.2	100.5	10	38.3	98.4	10	38.2	98.0	10	35.0	89.7	10
85	39.4	10		41.2	104.5	10	40.0	101.5	10	39.1	99.3	10	39.1	99.3	10	35.2	89.4	10
EOS	41.2	10		42.4	102.9	10	41.6	100.8	10	40.7	98.6	10	40.4	98.0	10	37.5	91.0	10

3 EOS = end of study.


4 ^aStudy day 1 is the day animals were placed on study.

1 **Table 33. Summary of Survival and Body Weights of Female Mice in the Three-month Drinking Water Study of Sodium Tungstate**
 2 **Dihydrate**


Study Day ^a	0 mg/L			125 mg/L			250 mg/L			500 mg/L			1,000 mg/L			2,000 mg/L		
	Av. Wt. (g)	N		Av. Wt. (g)	Wt. (% of Controls)	N	Av. Wt. (g)	Wt. (% of Controls)	N	Av. Wt. (g)	Wt. (% of Controls)	N	Av. Wt. (g)	Wt. (% of Controls)	N	Av. Wt. (g)	Wt. (% of Controls)	N
1	17.1	10		16.7	97.6	10	17.2	100.1	10	17.3	101.0	10	16.9	98.8	10	16.8	98.1	10
8	19.4	10		18.6	95.9	10	18.8	96.9	10	19.1	98.3	10	18.8	96.5	10	18.5	95.1	10
15	20.3	10		19.8	97.6	10	19.6	96.5	10	20.2	99.5	10	19.5	95.8	10	19.8	97.3	10
22	21.9	10		21.3	97.4	10	21.2	96.8	10	22.0	100.5	10	20.8	95.1	10	20.8	95.2	10
29	23.0	10		22.1	96.2	10	21.2	92.2	10	22.4	97.5	10	21.0	91.6	10	21.6	94.0	10
36	24.8	10		22.8	91.9	10	22.6	91.0	10	23.6	94.9	10	22.1	89.0	10	22.2	89.3	10
43	26.0	10		24.3	93.6	10	23.5	90.5	10	24.3	93.4	10	23.1	89.0	10	23.4	90.2	10
50	27.8	10		26.0	93.6	10	24.6	88.4	10	26.2	94.3	10	24.9	89.5	10	24.5	88.1	10
57	28.9	10		27.9	96.5	10	25.0	86.6	10	27.0	93.4	10	26.1	90.4	10	26.1	90.3	10
64	30.3	10		28.5	93.9	10	26.0	85.7	10	28.2	92.9	10	26.4	86.9	10	27.8	91.6	10
71	31.3	10		29.4	93.7	10	26.7	85.3	10	30.3	96.6	10	27.9	89.0	10	28.1	89.8	10
78	32.1	10		31.1	96.9	10	27.6	86.2	10	30.3	94.4	10	29.2	91.0	10	28.7	89.6	10
85	31.7	10		32.7	103.1	10	29.0	91.5	10	31.1	98.0	10	29.5	93.1	10	28.7	90.6	10
EOS	31.7	10		33.8	106.6	10	29.9	94.1	10	31.7	99.9	10	29.5	92.8	10	29.1	91.6	10

3 EOS = end of study.

4 ^aStudy day 1 is the day animals were placed on study.



1



2

3

4

Figure 12. Growth Curves for Mice Exposed to Sodium Tungstate Dihydrate in Drinking Water for Three Months

1 **Table 34. Summary of Water and Sodium Tungstate Dihydrate Consumption of Male and Female Mice in the Three-month Drinking**
 2 **Water Study**

Week	0 mg/L		125 mg/L		250 mg/L		500 mg/L		1,000 mg/L		2,000 mg/L	
	Water (g/day) ^a	Water (g/day)	Dose (mg/kg/day) ^b	Water (g/day)	Dose (mg/kg/day)	Water (g/day)	Dose (mg/kg/day)	Water (g/day)	Dose (mg/kg/day)	Water (g/day)	Dose (mg/kg/day)	
Male												
1	4.3 ^d	3.8	21.2	3.7	41.0	3.5	77.2	3.6	162.6	3.4	303.7	
4	3.7	3.5	15.3	3.4	30.0	3.4	61.0	3.3	121.7	3.1	234.0	
13	3.5	3.5	10.6	3.6	22.5	3.4	43.4	3.1	79.2	3.0	170.4	
Female												
1	2.5	2.4	17.9	2.6	37.9	2.5	72.2	2.2	129.9	2.0	237.8	
4	3.0	2.6	15.2	2.6	30.7	2.7	61.4	2.7	129.8	2.3	220.9	
13	2.2	2.6	9.9	2.6	22.4	2.6	41.8	2.4	81.4	2.3	160.1	

3 ^aWater consumption data are presented as grams/animal/day.

4 ^bMilligrams of sodium tungstate dihydrate consumed/kilogram body weight/day.

1 Blood was collected from up to 10 animals per group on the morning of day 91. Total blood
 2 tungsten concentrations were determined using a validated analytical method (Appendix E); data
 3 are presented in Table 35. Tungsten was not detected in males in the vehicle control group above
 4 the LOD of the assay (0.0016 µg/g); however, low concentrations of tungsten were detected in
 5 females in the vehicle control group. In both males and females, the blood tungsten
 6 concentrations increased proportionally with the exposure concentration; there was no observed
 7 sex difference. In females, the tungsten concentrations in exposed groups were significantly
 8 higher than in the corresponding vehicle control group.

9 **Table 35. Summary of Blood Tungsten Concentration Data for Male and Female Mice in the**
 10 **Three-month Drinking Water Study of Sodium Tungstate Dihydrate^a**

	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Male						
n	10	10	10	10	10	10
Blood (µg/g)	BD	0.17 ± 0.04	0.28 ± 0.04	0.47 ± 0.10	1.56 ± 0.27	2.63 ± 0.70
Female						
n	10	10	10	9	10	10
Blood (µg/g) ^{b,c}	0.03 ± 0.01**	0.13 ± 0.02**	0.27 ± 0.05**	0.44 ± 0.07**	1.06 ± 0.11**	2.87 ± 0.40**

11 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

12 Statistical significance for the vehicle control group indicates a significant trend test.

13 **Statistically significant at $p \leq 0.01$.

14 BD = below detection; group did not have more than 20% of its values above the limit of detection (LOD).

15 ^aData presented as mean ± standard error.

16 ^bValues below the LOD (0.0016 µg/g), were substituted with 1/2 the LOD value. If 80% or more of the values in the vehicle
 17 control group were below the LOD, no mean or standard error were calculated, and no statistical analysis was performed.

18 ^cStatistical analysis performed by the Jonckheere (trend) and Shirley or Dunn (pairwise) tests.

19 In male mice, the white blood cell count was significantly decreased in the 1,000 and 2,000 mg/L
 20 groups relative to the vehicle control group (Table 36). These decreases were driven by a
 21 significant decrease in the lymphocyte count in the 1,000 and 2,000 mg/L groups and a
 22 significant decrease in the monocyte count in the 500 mg/L and higher groups (Table 36).
 23 Additionally, the eosinophil counts were significantly decreased in all ST-exposed male groups.
 24 These leukocyte changes are consistent with a stress leukogram (i.e., effects of chronic increase
 25 in endogenous corticosterone).⁸⁷

1 **Table 36. Summary of Select Hematology Data for Male Mice in the Three-month Drinking Water**
 2 **Study of Sodium Tungstate Dihydrate^{a,b}**

	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
n	10	10	10	10	10	10
White Blood Cells ($10^3/\mu\text{L}$)	5.91 ± 0.52**	4.87 ± 0.61	5.32 ± 0.42	4.94 ± 0.51	4.20 ± 0.37**	3.83 ± 0.56**
Lymphocytes ($10^3/\mu\text{L}$)	4.83 ± 0.45**	3.93 ± 0.49	4.38 ± 0.34	4.07 ± 0.43	3.41 ± 0.31**	3.12 ± 0.45**
Monocytes ($10^3/\mu\text{L}$)	0.20 ± 0.02**	0.13 ± 0.02	0.14 ± 0.02	0.10 ± 0.02**	0.09 ± 0.02**	0.09 ± 0.03**
Eosinophils ($10^3/\mu\text{L}$)	0.15 ± 0.03**	0.08 ± 0.01*	0.08 ± 0.01*	0.07 ± 0.02*	0.07 ± 0.01**	0.06 ± 0.02**

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

6 ^aData displayed as mean ± standard error.

7 ^bStatistical analysis performed by the Jonckheere (trend) and Shirley or Dunn (pairwise) tests.

8 A higher group mean relative testis weight was observed in the 2,000 mg/L male group, relative
 9 to the vehicle control group, and was likely due to the lower mean body weights in that group
 10 (Appendix G). In female mice, there were several sporadic increases in group mean organ
 11 weights, but they lacked an exposure concentration response or other supporting evidence that
 12 they represented anything but biological variation (Appendix G). Relative kidney weights were
 13 significantly increased in the 1,000 and 2,000 mg/L females and 2000 mg/L males but were most
 14 likely due to reduced mean body weights in those groups compared to the vehicle control group
 15 (Table 37).

16 No exposure-related gross lesions were recorded (Appendix G). The only histological lesion
 17 associated with exposure was in the kidney. The incidences of renal tubule regeneration were
 18 higher in the 1,000 and 2,000 mg/L male and female groups compared to the respective vehicle
 19 control groups; the increases in the male groups were significant (Table 37). The lesion consisted
 20 of hyperplastic tubules, predominantly in the deep cortical to medullary region, lined by
 21 epithelial cells with increased cytoplasmic basophilia, nuclear crowding, prominent nucleoli,
 22 margined chromatin, and karyomegaly. There were occasional mitotic figures.

23 **Table 37. Summary of Renal Findings for Male and Female Mice in the Three-month Drinking**
 24 **Water Study of Sodium Tungstate Dihydrate**

	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
n	10	10	10	10	10	10
Male						
Necropsy Body Wt. (g) ^a	41.2 ± 1.0**	42.4 ± 0.9	41.6 ± 1.5	40.7 ± 1.1	40.4 ± 0.8	37.5 ± 1.1*
R. Kidney Weight ^a						
Absolute (g)	0.30 ± 0.00	0.30 ± 0.01	0.31 ± 0.01	0.31 ± 0.01	0.31 ± 0.01	0.29 ± 0.00
Relative (mg/g) ^b	7.28 ± 0.15**	7.15 ± 0.16	7.47 ± 0.18	7.53 ± 0.22	7.75 ± 0.16	7.84 ± 0.18*
Histological Findings						
Kidney ^c	10	10	10	10	10	10
Renal tubule, regeneration ^d	0**	0	0	0	6** (1.0) ^e	10** (1.6)

	0 mg/L	125 mg/L	250 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Female						
Necropsy Body Wt. (g)	31.7 ± 1.7*	33.8 ± 1.3	29.9 ± 1.1	31.7 ± 1.3	29.5 ± 1.1	29.1 ± 1.2
R. Kidney Weight						
Absolute (g)	0.15 ± 0.00	0.17 ± 0.00	0.16 ± 0.00	0.16 ± 0.00	0.16 ± 0.00	0.16 ± 0.00
Relative (mg/g)	4.88 ± 0.19**	4.91 ± 0.11	5.28 ± 0.11	5.22 ± 0.13	5.48 ± 0.16*	5.42 ± 0.19*
Histological Findings						
Kidney	10	10	10	10	10	10
Renal tubule, regeneration	0*	0	0	0	1 (1.0)	2 (1.0)

1 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

2 Statistical significance for the vehicle control group indicates a significant trend test.

3 Statistical analysis for organ weight data performed by the Jonckheere (trend) and Williams or Dunnett (pairwise) tests.

4 Statistical analysis for histological findings performed by the Poly-3 test.

5 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

6 ^aData presented as mean ± standard error.

7 ^bRelative organ weights (organ-weight-to-body-weight ratios) are given as mg organ weight/g body weight.

8 ^cNumber of animals examined microscopically.

9 ^dNumber of animals with lesion.

10 ^eAverage severity grade of observed lesion in affected animals: 1 = minimal; 2 = mild; 3 = moderate; 4 = marked.

11 In male mice, no significant differences were observed between exposed groups and the vehicle
 12 control group in left testis weight, left epididymal weight, left cauda weight, or any of the sperm
 13 parameters, including number of sperm/mg cauda epididymis, total number sperm/cauda, sperm
 14 motility, number of homogenization-resistant spermatids/mg testis, and total number of
 15 spermatids (Appendix G). The testes and epididymides were evaluated to a no-effect level, and
 16 no histological findings associated with ST exposure were present at 3 months. Under the
 17 conditions of this 3-month study, ST administration via drinking water did not exhibit the
 18 potential to be a reproductive toxicant in B6C3F1/N mice. Testicular lesions observed in male
 19 mice after 2 years of exposure to ST, however, might possibly impair reproductive performance.

20 Exposure Concentration Selection Rationale for Two-year Studies in Mice

21 In the 3-month studies, there were no effects on survival and mean body weights in any exposure
 22 groups that were considered exposure concentration-limiting (Table 32, Table 33). There was a
 23 16% reduction in water consumption in male mice at the top exposure group of 2,000 mg/L
 24 (Table 34), and minimal to mild renal tubule regeneration was noted in the 1,000 and 2,000 mg/L
 25 groups, but neither effect was considered exposure concentration-limiting (Table 37). Hence,
 26 exposure concentrations selected for the 2-year study were 0, 500, 1,000, and 2,000 mg/L.

1 **Two-year Study (Interim Evaluations – 3, 6, 12, and 18 Months)**

2 Mean body weights of the 2,000 mg/L males were significantly decreased by approximately 12%
3 relative to the vehicle control group at 6 months, and mean body weights of the 2,000 mg/L
4 females were significantly decreased by approximately 13% relative to the vehicle control group
5 at 12 months (Table 38, Table 39). Mean body weights of all other groups, both males and
6 females, were within 10% of the vehicle control groups at all time points.

7 At 3 months in males, mean absolute kidney weights were significantly increased in the
8 500 mg/L and 1,000 mg/L groups by approximately 10% and 13%, respectively, and mean
9 relative kidney weight was significantly increased in the 500 mg/L group relative to the vehicle
10 control group (Table 38). In females at 3 months, mean relative kidney weight was significantly
11 decreased in the 1,000 mg/L group relative to the vehicle control group (Table 39). Mean relative
12 kidney weights were significantly increased at 6 months in the 500 and 1,000 mg/L males
13 relative to the vehicle control group; there were no significant differences in kidney weights
14 among female groups. At 12 months, there were no significant differences in the kidney weights
15 among male groups, but in females, mean relative kidney weights were significantly increased in
16 the 1,000 and 2,000 mg/L groups relative to the vehicle control group. At 18 months, there were
17 no significant pairwise differences in kidney weights in either male or female groups compared
18 to the vehicle control groups.

19 At 3 months, mean relative liver weights were significantly increased in the 2,000 mg/L males
20 and females relative to the vehicle control groups. At 6 months, mean absolute liver weights
21 were significantly decreased by approximately 13% in the 1,000 mg/L males and significantly
22 decreased by approximately 19% in the 2,000 mg/L males relative to the vehicle control males
23 (Table 38, Table 39). The biological importance of these changes in liver weights is unknown,
24 but they are likely of little toxicological significance.

25 Other sporadic differences in organ weights were considered isolated changes of no toxicological
26 significance (Appendix G).

1 **Table 38. Summary of Select Organ Weights and Organ-Weight-to-Body-Weight Ratios for Male**
 2 **Mice Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months^{a,b}**

	0 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Three Months				
Necropsy Body Wt. (g)	34.7 ± 0.8 (10)	35.3 ± 1.2 (10)	37.0 ± 1.1 (10)	31.2 ± 1.0 (10)
Kidneys				
Absolute (g)	0.52 ± 0.01 (10)	0.57 ± 0.01* (10)	0.59 ± 0.01** (10)	0.50 ± 0.01 (10)
Relative (mg/g) ^c	14.95 ± 0.33* (11)	16.40 ± 0.48* (10)	15.91 ± 0.36 (10)	16.19 ± 0.27 (10)
Liver				
Absolute (g)	1.37 ± 0.03 (10)	1.43 ± 0.03 (10)	1.51 ± 0.06 (10)	1.30 ± 0.03 (10)
Relative (mg/g)	39.52 ± 0.27** (11)	40.72 ± 0.95 (10)	40.74 ± 0.62 (10)	41.83 ± 0.67* (10)
Six Months				
Necropsy Body Wt. (g)	45.0 ± 1.0** (11)	42.5 ± 0.8 (10)	42.1 ± 1.1* (10)	39.8 ± 0.8** (10)
Kidneys				
Absolute (g)	0.67 ± 0.02 (11)	0.67 ± 0.02 (10)	0.69 ± 0.02 (10)	0.62 ± 0.01 (10)
Relative (mg/g)	14.80 ± 0.26* (12)	15.87 ± 0.20* (10)	16.47 ± 0.30** (10)	15.69 ± 0.33 (10)
Liver				
Absolute (g)	1.98 ± 0.07** (11)	1.77 ± 0.10 (10)	1.72 ± 0.09* (10)	1.60 ± 0.05** (10)
Relative (mg/g)	43.59 ± 0.94 (12)	41.50 ± 1.63 (10)	40.67 ± 1.12 (10)	40.23 ± 0.66 (10)
Twelve Months				
Necropsy Body Wt. (g)	49.4 ± 1.0 (10)	48.1 ± 0.7 (10)	48.1 ± 0.8 (10)	46.9 ± 0.8 (10)
Kidneys				
Absolute (g)	0.81 ± 0.02 (10)	0.80 ± 0.02 (10)	0.82 ± 0.03 (10)	0.78 ± 0.02 (10)
Relative (mg/g)	16.43 ± 0.29 (10)	16.70 ± 0.25 (10)	17.12 ± 0.47 (10)	16.62 ± 0.29 (10)
Liver				
Absolute (g)	2.42 ± 0.15 (10)	2.18 ± 0.05 (10)	2.36 ± 0.14 (10)	2.13 ± 0.08 (10)
Relative (mg/g)	48.93 ± 3.00 (10)	45.37 ± 0.62 (10)	49.18 ± 3.15 (10)	45.56 ± 1.79 (10)
Eighteen Months				
Necropsy Body Wt. (g)	49.4 ± 1.7* (9)	48.4 ± 0.9 (10)	48.3 ± 1.2 (9)	46.4 ± 1.0 (6)
Kidneys				
Absolute (g)	0.79 ± 0.02 (9)	0.79 ± 0.02 (10)	0.85 ± 0.05 (9)	0.74 ± 0.02 (6)
Relative (mg/g)	16.07 ± 0.36 (9)	16.35 ± 0.24 (10)	17.69 ± 0.94 (9)	15.90 ± 0.35 (6)
Liver				
Absolute (g)	2.25 ± 0.19 (9)	2.45 ± 0.20 (10)	2.79 ± 0.30 (9)	1.86 ± 0.14 (6)
Relative (mg/g)	46.98 ± 6.36 (9)	50.98 ± 4.76 (10)	58.20 ± 6.42 (9)	39.95 ± 2.41 (6)

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

6 ^aData presented as mean ± standard error (n).

7 ^bStatistical analysis performed by the Jonckheere (trend) and the Williams or Dunnett (pairwise) tests.

8 ^cRelative organ weights (organ-weight-to-body-weight ratios) are given as mg organ weight/g body weight.

1 **Table 39. Summary of Select Organ Weights and Organ-Weight-to-Body-Weight Ratios for Female**
 2 **Mice Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months^{a,b}**

	0 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Three Months				
Necropsy Body Wt. (g)	27.3 ± 1.5 (10)	24.8 ± 0.7 (10)	29.2 ± 1.0 (10)	24.8 ± 0.9 (10)
Kidneys				
Absolute (g)	0.32 ± 0.01 (10)	0.31 ± 0.01 (10)	0.32 ± 0.01 (10)	0.31 ± 0.01 (10)
Relative (mg/g) ^c	11.88 ± 0.33 (10)	12.58 ± 0.28 (10)	10.89 ± 0.30 (10)*	12.41 ± 0.21 (10)
Liver				
Absolute (g)	1.13 ± 0.04 (10)	1.09 ± 0.03 (10)	1.17 ± 0.02 (10)	1.13 ± 0.04 (10)
Relative (mg/g)	41.97 ± 1.23 (10)	43.76 ± 0.73 (10)	40.24 ± 0.92 (10)	45.60 ± 0.89* (10)
Six Months				
Necropsy Body Wt. (g)	39.8 ± 1.3 (10)	38.7 ± 1.0 (10)	39.3 ± 1.5 (10)	39.4 ± 1.4 (10)
Kidneys				
Absolute (g)	0.36 ± 0.01 (10)	0.36 ± 0.01 (10)	0.36 ± 0.01 (10)	0.36 ± 0.01 (10)
Relative (mg/g)	9.07 ± 0.26 (10)	9.36 ± 0.24 (10)	9.10 ± 0.24 (10)	9.15 ± 0.30 (10)
Liver				
Absolute (g)	1.37 ± 0.03 (10)	1.34 ± 0.02 (10)	1.36 ± 0.04 (10)	1.34 ± 0.03 (10)
Relative (mg/g)	34.63 ± 0.77 (10)	34.75 ± 0.59 (10)	34.83 ± 0.69 (10)	34.28 ± 0.86 (10)
Twelve Months				
Necropsy Body Wt. (g)	53.9 ± 0.8** (10)	50.3 ± 1.7 (10)	51.1 ± 1.6 (10)	46.7 ± 1.6** (10)
Kidneys				
Absolute (g)	0.41 ± 0.01 (10)	0.39 ± 0.02 (10)	0.41 ± 0.01 (10)	0.39 ± 0.01 (10)
Relative (mg/g)	7.51 ± 0.19** (10)	7.80 ± 0.14 (10)	8.10 ± 0.16* (10)	8.34 ± 0.27** (10)
Liver				
Absolute (g)	1.61 ± 0.08 (10)	1.48 ± 0.07 (10)	1.58 ± 0.05 (10)	1.43 ± 0.03 (10)
Relative (mg/g)	29.84 ± 1.34 (10)	29.38 ± 0.98 (10)	30.91 ± 0.71 (10)	30.97 ± 1.09 (10)
Eighteen Months				
Necropsy Body Wt. (g)	56.3 ± 1.3 (10)	55.0 ± 1.9 (9)	53.3 ± 2.6 (9)	56.3 ± 1.2 (10)
Kidneys				
Absolute (g)	0.44 ± 0.01 (10)	0.44 ± 0.02 (9)	0.45 ± 0.02 (9)	0.47 ± 0.01 (10)
Relative (mg/g)	7.72 ± 0.12* (10)	8.06 ± 0.23 (9)	8.55 ± 0.56 (9)	8.33 ± 0.17 (10)
Liver				
Absolute (g)	1.57 ± 0.07 (10)	1.51 ± 0.06 (9)	1.54 ± 0.06 (9)	1.61 ± 0.05 (10)
Relative (mg/g)	27.91 ± 0.78 (10)	27.61 ± 0.72 (9)	29.12 ± 1.08 (9)	28.63 ± 0.57 (10)

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

6 ^aData presented as mean ± standard error (n).

7 ^bStatistical analysis performed by the Jonckheere (trend) and the Williams or Dunnett (pairwise) tests.

8 ^cRelative organ weights (organ-weight-to-body-weight ratios) are given as mg organ weight/g body weight.

1 Plasma, kidney, and urine from up to 10 animals per group were collected at 3, 6, 12, and
2 18 months from interim animals. Tungsten concentrations in all matrices were determined using
3 validated analytical methods (Appendix E). In male mice, plasma tungsten concentrations
4 increased proportionally with the exposure concentration at all time points (except at the
5 18 months in the 1,000 mg/L group) (Table 40; Figure 13A). In female rats, plasma tungsten
6 concentrations increased proportionally with the exposure concentration up to 1,000 mg/L for all
7 time points (except at 18 months in the 500 mg/L group); however, at 2,000 mg/L, the trend was
8 toward a more-than-proportional increase in plasma tungsten concentration with increasing
9 exposure concentration (Table 41; Figure 13A). There was also a trend toward decreasing plasma
10 tungsten concentrations with increasing exposure duration in both male and female mice. Low
11 tungsten concentrations were observed in the vehicle control groups at some interim evaluations;
12 however, they were significantly lower than those in the lowest exposure groups (Table 40,
13 Table 41). There was no observed sex difference in plasma tungsten concentration in mice
14 (Table 40, Table 41; Figure 13A).

15 In male and female mice, tungsten concentrations in the kidney increased with increasing
16 exposure concentration, but the trend was less than proportional (Figure 13B). There was also a
17 trend toward decreasing tungsten concentrations with increasing exposure duration in both male
18 and female mice, with the highest concentrations at 3 months. Low tungsten concentrations were
19 observed in the kidneys of some animals in the vehicle control group; however, they were
20 significantly lower than those in the lowest exposure group (Table 40, Table 41). The
21 kidney-to-plasma ratios ranged from 1.43 to 4.36 suggesting retention of tungsten in the kidney.
22 There were no consistent trends in the kidney-to-plasma ratios with increasing exposure
23 concentration or duration. There was no observed sex difference in kidney tungsten
24 concentrations in mice (Table 40, Table 41; Figure 13B).

25 The concentrations of tungsten in urine are presented as both $\mu\text{g/g}$ of urine and $\mu\text{g/mg}$ creatinine.
26 Creatinine-corrected tungsten concentrations in urine increased proportionally with the exposure
27 concentration for both males and females (Figure 13C). As with plasma and kidney, the trend
28 was toward decreasing tungsten concentrations in urine with increasing exposure duration in
29 both male and female mice. Low tungsten concentrations were observed in the urine of some
30 vehicle control groups; however, tungsten concentrations in exposed groups were significantly
31 higher than those in corresponding vehicle control groups (Table 40, Table 41). There was no
32 observed sex difference in urinary tungsten concentrations in mice (Table 40, Table 41;
33 Figure 13C).

1 **Table 40. Summary of Plasma, Kidney, and Urine Tungsten Concentration Data for Male Mice**
 2 **Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months^a**

	0 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Three Months				
Kidney (µg/g)	BD	5.71 ± 0.88 (10)	6.50 ± 0.57 (10)	12.15 ± 2.16 (10)
Plasma (µg/mL)	BD	2.62 ± 0.47 (10)	4.11 ± 0.58 (10)	8.10 ± 1.14 (10)
Kidney/Plasma Ratio ^b	BD	2.45 ± 0.31 (10)	1.73 ± 0.16 (10)	1.51 ± 0.13 (10)
Urine ^{c,d}				
Urine (µg/mL urine)	0.02 ± 0.01** (7)	191.56 ± 41.89** (7)	592.00 ± 69.45** (10)	1,335.33 ± 148.26** (6)
Urine (µg/mg creatinine)	0.03 ± 0.01** (7)	353.16 ± 27.44** (7)	805.94 ± 70.58** (10)	1,672.99 ± 147.92** (8)
Six Months				
Kidney (µg/g) ^{c,d}	0.06 ± 0.01** (10)	3.79 ± 0.29** (10)	5.89 ± 1.23** (10)	12.10 ± 1.39** (10)
Plasma (µg/mL) ^{c,d}	0.02 ± 0.00** (10)	1.68 ± 0.22** (10)	2.43 ± 0.57** (10)	5.82 ± 0.96** (10)
Kidney/Plasma Ratio ^c	3.46 ± 0.42* (10)	2.40 ± 0.18 (10)	2.58 ± 0.24 (10)	2.24 ± 0.19 (10)
Urine ^{c,d}				
Urine (µg/mL urine)	3.92 ± 2.77** (8)	204.50 ± 28.06** (8)	407.89 ± 92.69** (8)	816.50 ± 144.16** (8)
Urine (µg/mg creatinine)	6.43 ± 4.14** (8)	375.17 ± 45.85** (8)	709.28 ± 117.50** (7)	1,368.62 ± 149.28** (7)
Twelve Months				
Kidney (µg/g)	BD	3.67 ± 0.61 (10)	6.69 ± 0.59 (10)	10.45 ± 1.06 (10)
Plasma (µg/mL) ^{c,d}	0.02 ± 0.00** (10)	1.07 ± 0.16** (10)	2.37 ± 0.33** (10)	4.84 ± 0.73** (10)
Kidney/Plasma Ratio	BD	3.66 ± 0.48 (10)	3.10 ± 0.27 (10)	2.35 ± 0.18 (10)
Urine ^{c,d}				
Urine (µg/mL urine)	4.00 ± 1.69** (10)	152.57 ± 27.46** (10)	349.00 ± 38.70** (10)	661.09 ± 123.57** (10)
Urine (µg/mg creatinine)	15.57 ± 8.39** (10)	404.82 ± 34.78** (9)	905.34 ± 65.48** (10)	1,390.13 ± 183.53** (8)
Eighteen Months				
Kidney (µg/g) ^{c,d}	0.06 ± 0.01** (9) ^e	2.67 ± 0.23** (9)	5.50 ± 0.73** (8)	8.71 ± 0.99** (6) ^f
Plasma (µg/mL) ^{c,d}	0.01 ± 0.00** (9)	1.27 ± 0.19** (10)	4.95 ± 2.03** (9)	4.86 ± 0.65** (6)
Kidney/Plasma Ratio ^c	6.28 ± 1.23** (9)	2.41 ± 0.21** (9)	2.42 ± 0.34** (8)	1.85 ± 0.16** (6)
Urine ^{c,d}				
Urine (µg/mL urine)	0.02 ± 0.00** (9)	104.52 ± 11.81** (10)	194.00 ± 14.23** (8)	661.00 ± 161.34** (6)
Urine (µg/mg creatinine)	0.05 ± 0.01** (9)	278.92 ± 30.45** (10)	579.75 ± 52.95** (8)	1,279.52 ± 155.83** (6)

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 *Statistically significant at $p \leq 0.05$; ** $p \leq 0.01$.

6 BD = below detection; group did not have more than 20% of its values above the limit of detection (LOD).

7 ^aData presented as mean ± standard error (n).

8 ^bFor the kidney/plasma ratio calculation, a plasma density of 1 g/mL was assumed.

9 ^cStatistical analysis performed by the Jonckheere (trend) and Shirley or Dunn (pairwise) tests.

10 ^dValues below the LOD (0.013 µg/mL) were substituted with 1/2 the LOD value. If 80% or more of the values in the vehicle control group were below the LOD, no mean or standard error were calculated, and no statistical analysis was performed.

11 ^eThe kidney concentration value for one male in the 500 mg/L group at 18 months was excluded from the analysis as an implausible value.

12 ^fThe kidney concentration value for one male in the 1,000 mg/L group at 18 months was excluded from the analysis as an implausible value.

1 **Table 41. Summary of Plasma, Kidney, and Urine Tungsten Concentration Data for Female Mice**
 2 **Exposed to Sodium Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months^a**

	0 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Three Months				
Kidney (µg/g)	BD	3.54 ± 0.27 (10)	5.34 ± 0.49 (10)	11.44 ± 1.22 (10)
Plasma (µg/mL)	BD	1.84 ± 0.23 (10)	3.25 ± 0.49 (10)	8.72 ± 1.25 (10)
Kidney/Plasma Ratio ^b	BD	2.05 ± 0.12 (10)	1.86 ± 0.17 (10)	1.43 ± 0.11 (10)
Urine ^{c,d}				
Urine (µg/mL urine)	0.08 ± 0.02** (10)	442.78 ± 33.39** (9)	681.00 ± 118.08** (7)	1,470.00 ± 114.30** (10)
Urine (µg/mg creatinine)	0.14 ± 0.04** (9)	596.49 ± 57.37** (9)	872.11 ± 105.33** (7)	2,149.65 ± 183.59** (10)
Six Months				
Kidney (µg/g) ^{c,d}	0.08 ± 0.02** (10)	2.90 ± 0.17** (10)	5.52 ± 0.40** (10)	10.00 ± 0.91** (10)
Plasma (µg/mL) ^{c,d}	0.02 ± 0.00** (10)	1.11 ± 1.13** (9)	1.95 ± 0.23** (10)	4.81 ± 0.61** (10)
Kidney/Plasma Ratio ^c	5.50 ± 2.51 (10)	2.79 ± 0.23 (9)	3.19 ± 0.40 (10)	2.19 ± 0.14 (10)
Urine ^{c,d}				
Urine (µg/mL urine)	0.11 ± 0.03** (9)	217.11 ± 20.48** (9)	419.60 ± 49.71** (10)	855.40 ± 106.56** (10)
Urine (µg/mg creatinine)	0.21 ± 0.05** (9)	432.30 ± 32.48** (9)	788.33 ± 89.19** (10)	1,594.25 ± 192.88** (10)
Twelve Months				
Kidney (µg/g) ^{c,d}	0.05 ± 0.01** (10)	2.30 ± 0.23** (10)	3.64 ± 0.22** (10)	6.59 ± 0.54** (10)
Plasma (µg/mL) ^{c,d}	0.04 ± 0.00** (10)	0.58 ± 0.09** (10)	1.35 ± 0.14** (10)	4.59 ± 1.25** (9) ^e
Kidney/Plasma Ratio ^c	1.13 ± 0.15 (10)	4.36 ± 0.42** (10)	2.84 ± 0.18** (10)	2.05 ± 0.32 (9)
Urine ^{c,d}				
Urine (µg/mL urine)	0.13 ± 0.03** (10)	167.01 ± 26.29** (10)	397.40 ± 63.15** (10)	1,070.20 ± 182.92** (10)
Urine (µg/mg creatinine)	0.26 ± 0.04** (10)	309.67 ± 42.88** (10)	645.14 ± 83.24** (10)	1,642.14 ± 306.17** (10)
Eighteen Months				
Kidney (µg/g)	BD	2.90 ± 0.26 (9)	3.85 ± 0.80 (9)	5.32 ± 0.44 (10)
Plasma (µg/mL)	BD	1.78 ± 0.32 (9)	1.66 ± 0.39 (9)	3.29 ± 1.15 (10)
Kidney/Plasma Ratio	BD	1.83 ± 0.16 (9)	2.45 ± 0.16 (9)	2.76 ± 0.60 (10)
Urine ^{c,d}				
Urine (µg/mL urine)	0.04 ± 0.01** (9) ^f	74.25 ± 15.40** (9)	106.61 ± 28.16** (9)	245.04 ± 37.25** (10)
Urine (µg/mg creatinine) ^{c,d,g}	0.19 ± 0.04** (4)	262.73 ± 28.01** (8)	397.77 ± 38.06** (6)	851.15 ± 59.51** (9)

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 **Statistically significant at $p \leq 0.01$.

6 BD = below detection, group did not have more than 20% of its values above the limit of detection (LOD).

7 ^aData presented as mean ± standard error (n).

8 ^bFor the kidney/plasma ratio calculation, a plasma density of 1 g/mL was assumed.


9 ^cStatistical analysis performed by the Jonckheere (trend) and Shirley or Dunn (pairwise) tests

10 ^dValues below the LOD (0.013 µg/mL) were substituted with 1/2 the LOD value. If 80% or more of the values in the vehicle
 11 control group were below the LOD, no mean or standard error were calculated, and no statistical analysis was performed.

12 ^eThe plasma concentration value for one female in the 2,000 mg/L group at 12 months was excluded from the analysis as an
 13 implausible value.

14 ^fThe urine concentration value for one female in the 0 mg/L group at 18 months was excluded from the analysis as an implausible
 15 value.

16 ^gThe urine concentration values for five females in the 0 mg/L group, one female in the 500 mg/L group, three females in the
 17 1,000 mg/L group, and one female in the 2,000 mg/L group were excluded because samples were determined to be dilute.



1
2 **Figure 13. Tungsten Concentrations in Plasma, Kidney, and Urine in Mice Exposed to Sodium**
3 **Tungstate Dihydrate in Drinking Water for 3, 6, 12, and 18 Months**

4 Study day (SD) 93 = 3 months; SD 184 = 9 months; SD 366 = 12 months; SD 548 = 18 months.

1 Two-year Study

2 Survival

3 More mice in the exposed groups of males survived to study termination than in the vehicle
4 control group of males; however, the differences were not significant (Table 42; Figure 14).
5 Survival in females was similar across all groups.

6 **Table 42. Summary of Survival of Male and Female Mice in the Two-year Drinking Water Study of**
7 **Sodium Tungstate Dihydrate**

	0 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
Male				
Animals Initially in Study	50	50	50	50
Moribund	5	3	4	6
Natural Deaths	18	16	11	9
Accidental Deaths	1	–	–	–
Animals Surviving to Study Termination	26	31	35	35
Percent Probability of Survival at End of Study ^a	53.1% ^b	62.0%	70.0%	70.0%
Mean Survival (Days) ^c	683.4 ± 9.7	684.4 ± 11.6	710.4 ± 6.4	686.4 ± 13.1
Survival Analysis ^d	p = 0.107N	p = 0.530N	p = 0.081N	p = 0.158N
Female				
Animals Initially in Study	50	50	50	50
Moribund	2	1	2	0
Natural Deaths	10	8	10	10
Animals Surviving to Study Termination	38	41 ^e	38	40
Percent Probability of Survival at End of Study	76.0%	82.0%	76.0%	80.0%
Mean Survival (Days)	699.9 ± 10.2	713.5 ± 5.9	702.0 ± 10.9	704.0 ± 14.8
Survival Analysis	p = 0.809N	p = 0.582N	p = 1.000N	p = 0.743N


8 ^aKaplan-Meier determinations.

9 ^bCalculation does not include the accidental death animal.


10 ^cMean of litter means of all deaths (uncensored, censored, and study termination) ± standard error.

11 ^dThe result of the Cox proportional hazards trend test with random litter effects is in the vehicle control group column, and the
12 results of the proportional hazards pairwise comparisons to the vehicle control with random litter effects are in the exposed group
13 columns. A negative trend or lower mortality in an exposure group is indicated by N.

14 ^eIncludes one animal that died naturally during the last week of the study.



1



2

3 **Figure 14. Kaplan-Meier Survival Curves for Mice Exposed to Sodium Tungstate Dihydrate in**
4 **Drinking Water for Two Years**

1 **Body Weights, Water and Compound Consumption, and Clinical Observations**

2 At study termination, the mean body weight of the 2,000 mg/L males was 88% of the vehicle
3 control group; all other groups of exposed males and all groups of exposed females had mean
4 body weights within 10% of their respective vehicle control groups (Table 43, Table 44;
5 Figure 15).

6 Group mean water consumption over the course of the study for ST-exposed males for the 500,
7 1,000, and 2,000 mg/L groups averaged 96%, 90%, and 85% of the vehicle control group,
8 respectively (Table 45). For the ST-exposed females, group water consumption values for the
9 500, 1,000, and 2,000 mg/L groups averaged 105%, 97%, and 93% of the vehicle control group,
10 respectively (Table 46). Daily ST consumption for the 500, 1,000, and 2,000 mg/L groups
11 averaged 42.5, 80.0 and 158.1 mg/kg/day, respectively, for the males and 29.0, 56.2, and
12 107.0 mg/kg/day, respectively, for the females (Table 45, Table 46). In general, ingested dose
13 increased proportionally with the exposure concentration for both sexes. More occurrences of
14 thinness and ruffled fur were recorded in exposed groups of male mice compared to vehicle
15 control males; clinical observations were similar in all groups of females (Appendix G).

1 **Table 43. Summary of Survival and Mean Body Weights of Male Mice in the Two-year Drinking**
 2 **Water Study of Sodium Tungstate Dihydrate**

Study Day ^a	0 mg/L		500 mg/L			1,000 mg/L			2,000 mg/L		
	Av. Wt. (g)	No. of Survivors	Av. Wt. (g)	Wt. (% of Controls)	No. of Survivors	Av. Wt. (g)	Wt. (% of Controls)	No. of Survivors	Av. Wt. (g)	Wt. (% of Controls)	No. of Survivors
1	21.7	50	21.6	99.1	50	21.6	99.3	50	21.5	98.7	50
8	23.5	50	23.3	99.1	50	22.9	97.4	50	22.5	95.4	50
15	25.0	50	24.8	99.4	50	24.4	97.9	50	23.5	94.1	50
22	26.3	50	25.8	98.2	50	25.6	97.5	50	24.5	93.3	50
29	27.5	50	27.1	98.7	50	26.7	97.3	50	25.3	92.2	50
36	28.6	50	28.3	98.9	50	27.8	97.3	50	26.3	92.0	50
43	29.9	50	29.6	99.0	50	28.9	96.6	50	26.9	90.1	50
50	31.4	50	31.2	99.2	50	30.4	96.6	50	28.1	89.3	50
57	32.1	50	32.1	100.2	50	31.3	97.5	50	28.7	89.4	50
64	33.6	50	33.6	100.1	50	32.2	95.9	50	29.2	87.0	50
71	34.9	50	34.1	97.7	50	33.8	96.7	50	31.1	89.6	50
78	36.4	50	35.7	97.9	50	34.8	95.4	50	31.5	86.6	50
85	37.4	50	36.6	97.9	50	35.6	95.3	50	31.9	85.2	50
92	38.7	50	37.7	97.3	50	37.0	95.5	50	33.1	85.4	50
120	43.2	50	42.2	97.6	50	41.0	94.9	50	36.8	85.1	50
148	45.9	50	45.1	98.1	50	43.8	95.4	50	39.3	85.5	50
176	48.0	50	46.8	97.4	50	46.0	95.9	50	42.5	88.5	50
204	49.3	50	48.3	98.0	50	47.3	96.1	50	44.6	90.5	50
232	50.0	50	48.9	97.8	50	48.2	96.2	50	45.9	91.7	50
260	50.7	50	49.7	97.9	50	48.8	96.3	50	46.6	91.8	50
288	51.7	50	50.2	97.2	50	49.4	95.7	50	47.4	91.7	50
316	52.5	50	51.2	97.6	50	50.5	96.3	50	48.8	93.0	50
344	52.9	50	51.1	96.6	50	50.9	96.1	50	49.3	93.2	50
372	53.5	50	51.6	96.5	50	51.4	96.2	50	49.3	92.3	49
400	54.1	50	51.7	95.5	50	52.1	96.2	50	50.1	92.7	49
428	54.2	50	51.4	94.7	50	52.2	96.3	50	50.3	92.7	47
456	54.3	50	51.6	95.0	48	52.5	96.7	50	50.5	92.9	47
484	54.5	49	51.9	95.3	47	52.5	96.3	50	50.1	91.9	47
512	54.5	47	52.4	96.1	46	52.5	96.4	49	49.3	90.5	47
540	54.6	47	52.2	95.7	45	52.5	96.2	49	49.0	89.7	45
568	53.8	46	50.9	94.6	44	52.3	97.3	49	47.7	88.8	43
596	53.2	43	50.0	93.9	42	51.5	96.7	48	46.6	87.6	43
624	51.8	41	49.7	96.0	41	50.5	97.5	46	46.5	89.9	42
652	50.9	38	49.0	96.2	39	50.1	98.3	44	46.3	91.0	40
680	49.1	34	46.6	94.8	37	47.4	96.5	44	45.3	92.1	39
708	48.8	27	44.6	91.4	35	45.2	92.6	38	42.4	87.0	38
EOS	49.8	26	47.1	94.4	31	45.9	92.0	35	43.9	88.0	35

3 EOS = end of study.


4 ^aStudy day 1 is the day animals were placed on study.

1 **Table 44. Summary of Survival and Mean Body Weights of Female Mice in the Two-year Drinking**
 2 **Water Study of Sodium Tungstate Dihydrate**


Study Day ^a	0 mg/L		500 mg/L		1,000 mg/L			2,000 mg/L			
	Av. Wt. (g)	No. of Survivors	Av. Wt. (g)	Wt. (% of Controls)	No. of Survivors	Av. Wt. (g)	Wt. (% of Controls)	No. of Survivors	Av. Wt. (g)	Wt. (% of Controls)	No. of Survivors
1	15.9	50	15.7	99.2	50	16.1	101.4	50	16.1	101.3	50
8	17.5	50	17.5	99.8	50	17.0	97.1	50	17.3	98.9	50
15	18.5	50	18.4	99.6	50	18.2	98.3	50	18.2	98.6	50
22	19.5	50	19.4	99.3	50	18.9	96.7	50	19.2	98.5	50
29	20.5	50	20.5	100.3	50	20.1	98.3	50	19.9	97.2	50
36	21.2	50	21.1	99.4	50	21.0	99.0	50	20.7	97.6	50
43	22.6	50	22.4	99.1	50	22.0	97.7	50	21.7	96.2	50
50	22.9	50	22.8	99.9	50	22.2	97.1	50	22.4	98.1	49
57	23.7	50	23.3	98.4	50	23.0	97.2	50	23.3	98.5	49
64	24.6	50	24.7	100.0	50	23.9	97.2	50	23.9	97.1	49
71	25.5	50	25.5	99.8	50	25.0	98.0	50	25.1	98.2	49
78	26.7	50	26.5	99.2	50	26.0	97.2	50	25.7	96.4	49
85	27.8	50	27.0	96.9	50	26.8	96.2	50	26.9	96.6	49
92	28.6	50	28.2	98.7	50	27.5	96.0	50	27.5	96.2	49
120	33.5	50	32.5	97.2	50	31.5	94.2	50	31.5	94.2	49
148	38.1	50	37.6	98.7	50	35.6	93.5	50	36.3	95.2	49
176	41.7	50	41.2	98.8	50	38.9	93.4	50	39.7	95.3	49
204	45.6	50	44.6	97.8	50	42.0	92.2	50	42.8	93.9	49
232	48.2	50	47.0	97.5	50	44.3	91.8	50	45.2	93.8	49
260	50.2	50	49.3	98.2	50	47.0	93.7	49	47.9	95.3	49
288	52.3	50	51.7	98.9	50	48.5	92.7	49	49.5	94.5	49
316	54.9	50	53.9	98.3	50	51.1	93.0	49	51.8	94.3	49
344	55.0	50	54.3	98.7	50	51.7	93.9	49	52.2	94.8	49
372	56.8	50	56.7	99.8	50	53.7	94.4	49	54.3	95.6	49
400	58.5	50	58.2	99.4	50	55.3	94.4	49	55.1	94.2	49
428	57.9	49	58.2	100.5	50	55.3	95.5	49	55.9	96.6	49
456	59.8	48	59.1	98.9	50	56.8	95.0	49	56.9	95.1	48
484	60.8	48	60.1	98.8	50	57.9	95.2	49	57.6	94.8	48
512	61.4	48	60.0	97.8	50	58.2	94.9	49	57.7	94.1	48
540	61.6	48	61.1	99.2	49	59.1	96.0	49	58.6	95.2	48
568	61.2	45	60.7	99.0	49	58.6	95.7	49	57.8	94.3	48
596	61.2	44	59.4	97.2	49	57.6	94.2	49	58.1	95.0	48
624	61.1	44	59.0	96.5	48	58.2	95.2	44	57.4	93.8	48
652	59.9	44	59.5	99.4	45	58.3	97.3	43	56.3	93.9	48
680	59.0	42	57.6	97.6	44	57.8	97.9	42	56.0	95.0	45
708	56.6	39	53.6	94.7	42	54.3	96.0	41	52.7	93.1	43
EOS	57.1	38	54.9	96.1	40	54.9	96.2	38	53.8	94.2	40

3 EOS = end of study.

4 ^aStudy day 1 is the day animals were placed on study.



1



2

3

4

Figure 15. Growth Curves for Mice Exposed to Sodium Tungstate Dihydrate in Drinking Water for Two Years

1 **Table 45. Summary of Water and Sodium Tungstate Dihydrate Consumption of Male Mice in the**
 2 **Two-year Drinking Water Study**

Week	0 mg/L		500 mg/L		1,000 mg/L		2,000 mg/L	
	Water (g/day) ^a	Water (g/day)	Dose (mg/kg/day) ^b	Water (g/day)	Dose (mg/kg/day)	Water (g/day)	Dose (mg/kg/day)	
1	–	–	–	–	–	–	–	
13	3.2	3.3	45.1	3.0	84.2	2.8	175.7	
54	4.1	4.1	39.7	3.7	72.0	3.3	133.8	
102	4.9	4.1	46.0	4.3	95.1	3.8	179.1	

3 ^aGrams of water consumed/animal/day.

4 ^bMilligrams of sodium tungstate dihydrate consumed/kilogram body weight/day.

5 **Table 46. Summary of Water and Sodium Tungstate Dihydrate Consumption of Female Mice in the**
 6 **Two-year Drinking Water Study**

Week	0 mg/L		500 mg/L		1,000 mg/L		2,000 mg/L	
	Water (g/day) ^a	Water (g/day)	Dose (mg/kg/day) ^b	Water (g/day)	Dose (mg/kg/day)	Water (g/day)	Dose (mg/kg/day)	
1	1.9	1.8	57.2	1.8	111.9	1.7	211.5	
13	2.5	2.6	48.2	2.3	85.9	2.2	163.7	
54	2.4	2.5	22.0	2.3	42.9	2.4	88.4	
102	2.9	3.4	31.7	2.8	51.5	2.6	98.6	

7 ^aGrams of water consumed/animal/day.

8 ^bMilligrams of sodium tungstate dihydrate consumed/kilogram body weight/day.

9 **Histopathology**

10 This section describes the significant or biologically noteworthy changes in the incidences of
 11 neoplasms and nonneoplastic lesions of the kidney, liver, large intestine (cecum), and testes.
 12 Summaries of the incidences of neoplasms and nonneoplastic lesions and statistical analyses of
 13 primary neoplasms are in CEBS (Appendix G).

14 *Kidney:* Renal tubule neoplasms were only recorded in exposed males; there was one renal
 15 tubule adenoma in the 1,000 mg/L males and two renal tubule carcinomas in the 2,000 mg/L
 16 males (Table 47). Additional step sections of the kidneys were examined from all male and
 17 female mice, but additional renal tubule neoplasms were not observed. Proliferative lesions
 18 identified by the kidney step section pathologist were considered part of the spectrum of lesions
 19 associated with CPN by the pathology working group (PWG). Renal tubule regeneration was
 20 significantly increased in all exposed groups of males and in the 1,000 and 2,000 mg/L females,
 21 relative to vehicle control groups (Table 47). The incidences of pigment in the kidney were
 22 significantly increased compared to the vehicle control groups in the 2,000 mg/L males and
 23 1,000 mg/L females, and lower in the 500 and 1,000 mg/L males compared to the vehicle control
 24 group (Table 47). Given the high incidence of pigment in the male vehicle control group, and the
 25 lack of an exposure concentration response in either the males or females, the finding of pigment
 26 in the kidney of exposed animals is of questionable toxicological importance. Also, in males,
 27 there was a positive trend for renal tubule necrosis with increased exposure (Table 47).

1 The renal tubule adenoma was a discrete nodule that protruded slightly above the cortical surface
2 (Figure 16). It was a well-circumscribed cellular mass composed of well-differentiated renal
3 tubule epithelial cells. The renal cell carcinomas, in contrast to the adenoma, were large and
4 invasive, effacing and replacing more than 60% of the renal parenchyma (Figure 17). The
5 carcinomas had solid and cystic areas and displayed marked cellular pleomorphism and a high
6 mitotic rate.

7 A normal kidney from a B6C3F1/N mouse is shown in Figure 18. Renal tubule regeneration was
8 characterized by basophilia, karyomegaly, hypertrophy, and hyperplasia of the renal tubule
9 epithelium (Figure 19). Occasional mitotic figures were also present. Severity grading was based
10 on the amount of renal cortex involved, with minimal regeneration involving <10% of the cortex;
11 mild regeneration involving approximately 10–25% of the cortex; moderate regeneration
12 involving approximately 26–75% of the cortex, and marked regeneration involving over 75% of
13 the cortex. Renal tubule regeneration often occurred in kidneys that also had CPN, and the
14 lesions had to be separated diagnostically. Basophilic and hyperplastic tubules could also be seen
15 in CPN, but with CPN, the tubules would also have thickened basement membranes. Affected
16 renal tubules of CPN tended to lack the amount of karyomegaly, hypertrophy, and hyperplasia
17 that was characteristic of the epithelium seen in regeneration. Also, the pattern of the
18 regeneration was somewhat different from that of CPN—with low magnification, moderate to
19 severe regeneration could be seen to affect the cortex in a thick band sparing the medulla
20 (generally affecting the proximal convoluted tubule); the lesions of CPN typically were not so
21 evenly distributed.

22 Pigment in the kidney was light golden brown and was usually located in or around tubules with
23 CPN or regeneration, either in the renal tubule epithelium or in macrophages. All of the
24 occurrences of kidney pigment in females, and almost all in males, were of minimal severity.

25 Renal tubule necrosis was recorded in three 2,000 mg/L males and was characterized by renal
26 tubule epithelial cells with brightly eosinophilic cytoplasm and pyknotic or missing nuclei;
27 affected cells were often shrunken, irregular in shape, or sloughed off of their basement
28 membrane (Table 47). Mineral in the kidney was composed of very small irregular foci of darkly
29 basophilic material, typically found in the cortex or outer medulla. It was almost always of
30 minimal severity.

1 **Table 47. Incidences of Neoplastic and Nonneoplastic Lesions of the Kidney in Male and Female**
 2 **Mice in the Two-year Drinking Water Study of Sodium Tungstate Dihydrate**

	0 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
n^a	50	50	50	50
Male				
Renal Tubule, Regeneration ^b	2** (1.0) ^c	21** (1.4)	32** (1.4)	38** (1.6)
Renal Tubule, Necrosis	0*	0	0	3 (1.3)
Pigment	30** (1.0)	10 (1.0)	18 (1.2)	44** (1.0)
Renal Tubule, Adenoma ^d	0	0	1	0
Renal Tubule, Carcinoma ^e				
Overall rate ^f	0/50 (0%)	0/50 (0%)	0/50 (0%)	2/50 (4%)
Adjusted rate ^g	0%	0%	0%	4.6%
Terminal rate ^h	0/26 (0%)	0/31 (0%)	0/35 (0%)	2/35 (6%)
First incidence (days)	– ⁱ	–	–	730 (T)
Poly-3 test ^j	p = 0.047	(e)	(e)	p = 0.244
Renal Tubule, Adenoma or Carcinoma (Combined) ^k				
Overall rate	0/50 (0%)	0/50 (0%)	1/50 (2%)	2/50 (4%)
Adjusted rate	0%	0%	2.2%	4.6%
Terminal rate	0/26 (0%)	0/31 (0%)	1/35 (3%)	2/35 (6%)
First incidence (days)	–	–	730 (T)	730 (T)
Poly-3 test	p = 0.073	(e)	p = 0.520	p = 0.244
Female				
Renal Tubule, Regeneration	0**	1 (3.0)	7** (1.1)	7** (1.3)
Renal Tubule, Necrosis	1 (2.0)	5 (1.0)	1 (1.0)	3 (2.7)
Pigment	0	0	6* (1.0)	1 (1.0)

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.

5 *Statistically significant at $p \leq 0.05$ by the Poly-3 test; ** $p \leq 0.01$.

6 (T) = study termination; (e) = statistic could not be computed.

7 ^aNumber of animals examined microscopically.

8 ^bNumber of animals with lesion.

9 ^cAverage severity grade of observed lesion in affected animals: 1 = minimal; 2 = mild; 3 = moderate; 4 = marked.

10 ^dHistorical control incidence for all routes of 2-year studies (mean \pm standard deviation): 1/687 (0.15% \pm 0.55%);
 11 range: 0% to 2%.

12 ^eHistorical control incidence: 2/687 (0.31% \pm 1.11%); range: 0% to 4%.

13 ^fNumber of animals with neoplasm/number of animals necropsied.


14 ^gPoly-3 estimated neoplasm incidence after adjustment for intercurrent mortality.

15 ^hObserved incidence at terminal euthanasia.

16 ⁱNot applicable; no neoplasms in group.

17 ^jBeneath the control incidence is the p value associated with the trend test. Beneath the exposed group incidences are the p values
 18 corresponding to pairwise comparisons between the vehicle control group and that exposed group. The Poly-3 test accounts for
 19 differential mortality in animals that do not reach terminal euthanasia. A negative trend or a lower incidence in an exposure group
 20 is indicated by N.


21 ^kHistorical control incidence: 3/687 (0.46% \pm 1.66%); range: 0% to 6%.




1
2 **Figure 16. Renal Tubule Adenoma in a Male B6C3F1/N Mouse Exposed to 1,000 mg/L Sodium**
3 **Tungstate Dihydrate for Two Years (H&E)**

4 The adenoma is a well-circumscribed mass, which is protruding above the surface of the kidney. It is composed of a solid sheet
5 of well-differentiated epithelial cells.

A)




B)




6
7 **Figure 17. Renal Tubule Carcinoma in a Male B6C3F1/N Mouse Exposed to 2,000 mg/L Sodium**
8 **Tungstate Dihydrate for Two Years (H&E)**

9 A) The neoplasm is very large and has almost completely obliterated the kidney; only a small amount of uninvolved kidney
10 remains (arrows). B) Higher magnification of the renal tubule carcinoma in panel A; the neoplasm is densely cellular and one
11 area in particular (circle) displays cells with a very high nuclear-to-cytoplasmic ratio.

A)




B)




1
2 **Figure 18. Normal Kidney from a Vehicle Control Female B6C3F1/N Mouse in the Two-year Study**
3 **of Sodium Tungstate Dihydrate (H&E)**

4 A) This normal kidney is from a vehicle control female B6C3F1/N mouse. B) This shows a higher magnification of the normal
5 kidney in panel A.

A)



B)



1
2 **Figure 19. Moderate Renal Tubule Regeneration in a Female B6C3F1/N Mouse Exposed to**
3 **2,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E)**

4 A) This low magnification photomicrograph shows a thick band of basophilia due to regeneration of the renal tubules involving
5 the cortex (arrows). Compare with the kidney in Figure 18A. B) This higher magnification of the renal tubule regeneration in
6 panel A shows that the tubules are basophilic and the cells are crowded, consistent with hyperplasia and regeneration. There are
7 no thickened basement membranes or other evidence of chronic progressive nephropathy, such as associated interstitial
8 inflammation. Compare with the section of kidney in Figure 18B.

9 *Large Intestine, Cecum:* The incidences of pigment in the cecum were significantly increased in
10 the 1,000 and 2,000 mg/L males and females (Table 48). The pigment was found in
11 macrophages, and possibly fibroblasts, within the lamina propria of the cecum, and was similar
12 in character to that observed in the kidney; it was light golden brown (Figure 20). Cells
13 containing pigment were usually observed in small clusters, but were not associated with other
14 lesions, such as mucosal inflammation or ulceration. Special staining with Perl's stain for iron
15 did not reveal iron in the pigment, therefore, the pigment was not hemosiderin. Other
16 possibilities include pigments associated with cell breakdown, such as lipofuscin or ceroid, or ST
17 itself. Steps to further identify the pigment were not taken.

1 *Testes:* There were increased incidences of germinal epithelium degeneration recorded in the
2 testes of all exposed groups of males compared to the vehicle control group, and the increase was
3 significant in the 500 mg/L group (Table 48). Germinal epithelial degeneration consisted of
4 several changes, including vacuolation of the germinal epithelium, disorganization of the
5 germinal epithelium, or depletion of germs cells (Figure 21). Typically, especially with minimal
6 and mild lesions, only one or two seminiferous tubules were affected; however, the involvement
7 of only one seminiferous tubule might present as several cross sections of tubules within a
8 section of testis. Marked lesions were characterized by seminiferous tubules with only Sertoli
9 cells remaining (atrophy); this was usually a widespread to diffuse change throughout the testis.

10 *Bone Marrow and Spleen:* Hypercellularity of the bone marrow was significantly increased in
11 incidence in the 500 and 1,000 mg/L males; the incidence of extramedullary hematopoiesis in the
12 spleen was significantly increased in the 500 and 1,000 mg/L females (Table 48).
13 Hypercellularity of the bone marrow was characterized by cells filling the marrow cavity,
14 throughout the length of the femur. Little adipose tissue was observed. Both the erythroid and
15 myeloid cell lines appeared to have increased in most cases, although fixation with subsequent
16 decalcification made determining the exact type of many of the cells difficult. Megakaryocytes
17 were also abundant in affected animals. Most of the cases were minimal to mild in severity,
18 although an occasional animal had moderate to marked hypercellularity recorded; severity was
19 graded subjectively and was based on the amount of blood cell precursors and the lack of
20 adipocytes compared to the vehicle control animals and what would be expected in a chronic
21 study. Extramedullary hematopoiesis in the spleen is common in mice, even at 2 years. As such,
22 extramedullary hematopoiesis was recorded when it appeared increased over that which would
23 be expected in the spleen, although this was a subjective evaluation. It usually involved an
24 expansion of the red pulp, which was filled with cells containing little cytoplasm and dense
25 basophilic nuclei. Evidence of both erythroid and myeloid lineages were typically present, as
26 were abundant megakaryocytes. The biological significance of the differences in the incidences
27 of bone marrow hyperplasia and extramedullary hematopoiesis in the spleen is unknown.

28 *Other Tissues:* There were significantly increased incidences of hepatocellular adenomas and
29 carcinomas, combined, in the 500 mg/L females. This group also had a significant increase in the
30 incidences of eosinophilic foci and focal inflammation. There was no exposure concentration
31 relationship in the incidences of either the neoplastic or nonneoplastic lesions observed in the
32 liver, and the incidences of hepatocellular adenomas and hepatocellular carcinomas were within
33 the historical control range for each group (Appendix G).

34 A couple of tissues had a significantly increased incidence of a nonneoplastic lesion in one group
35 when compared to the vehicle control group, but these were not considered toxicologically
36 significant. They include inflammation of the tooth, which was significantly increased in
37 incidence in 2,000 mg/L males; and hyaline droplet accumulation in the respiratory epithelium of
38 the nose, which was significantly increased in 500 mg/L males (Appendix G).

1 **Table 48. Incidences of Select Nonneoplastic Lesions in Male and Female Mice in the Two-year**
 2 **Drinking Water Study of Sodium Tungstate Dihydrate**

	0 mg/L	500 mg/L	1,000 mg/L	2,000 mg/L
n^a	50	50	50	50
Male				
Large Intestine, Cecum				
Pigment ^b	3** (1.0) ^c	7 (1.0)	17** (1.0)	32** (1.0)
Testis				
Germinal epithelium, degeneration	11 (1.5)	20* (1.3)	20 (1.5)	20 (2.2)
Bone Marrow				
Hypercellularity	15 (1.2)	35** (1.7)	26* (1.6)	19 (1.4)
Female				
Large Intestine, Cecum				
Pigment	0**	3 (1.0)	7** (1.0)	14** (1.0)
Spleen				
Extramedullary hematopoiesis	5 (1.6)	18** (1.2)	13* (1.5)	8 (1.3)

3 Statistical significance for an exposure group indicates a significant pairwise test compared to the vehicle control group.

4 Statistical significance for the vehicle control group indicates a significant trend test.


5 *Statistically significant at $p \leq 0.05$ by the Poly-3 test; ** $p \leq 0.01$.

6 ^aNumber of animals examined microscopically.


7 ^bNumber of animals with lesion.

8 ^cAverage severity grade of observed lesion in affected animals: 1 = minimal; 2 = mild; 3 = moderate; 4 = marked.

A)



B)




1


2 **Figure 20. Pigment in the Cecum of B6C3F1/N Mice Exposed to 0 or 2,000 mg/L Sodium Tungstate**
3 **Dihydrate for Two Years (H&E)**

4 A) This normal cecum is from a B6C3F1/N vehicle control female mouse. B) In this cecum in a female B6C3F1/N mouse
5 exposed to 2,000 mg/L sodium tungstate dihydrate, the golden-brown pigment is contained primarily in macrophages within the
6 lamina propria (circle). Compare with the section of cecum in panel A.

A)



B)



1
2 **Figure 21. Germinal Epithelium Degeneration in the Testis from B6C3F1/N Mice Exposed to**
3 **2,000 mg/L Sodium Tungstate Dihydrate for Two Years (H&E)**

4 A) Testis from a B6C3F1/N mouse exposed to 2,000 mg/L sodium tungstate dihydrate (ST) is within normal limits. B) Testis
5 from a B6C3F1/N mouse exposed to 2,000 mg/L ST shows widespread degeneration of the germinal epithelium. A thinning of
6 the germinal epithelium of the seminiferous tubules is observed due to a decrease in the number of cells; compare with the testis
7 in panel A.

8 Genetic Toxicology

9 The genetic toxicity of ST was evaluated in bacterial reverse mutation assays and in both the
10 peripheral blood micronucleus test and the comet assay in rats and mice. ST significantly
11 increased DNA damage in the comet assay in liver cells from male and female rats, and in cells
12 from liver and ileum tissues in male mice (Appendix D).

13 ST (12.5 to 6,000 $\mu\text{g}/\text{plate}$) was not mutagenic in *Salmonella typhimurium* strains TA98 or
14 TA100, or in *Escherichia coli* WP2 *uvrA*/pKM101, when tested with or without exogenous
15 metabolic activation provided by phenobarbital/benzoflavone-induced rat S9 and cofactors
16 (Table D-1).

1 At the end of the 3-month studies, peripheral blood samples were obtained from male and female
2 rats and mice and analyzed for the frequency of micronucleated reticulocytes and erythrocytes
3 (Table D-2, Table D-3). In male and female rats, the reticulocyte population (polychromatic
4 erythrocytes, or PCEs)—which is the only red blood cell population that can be accurately
5 assessed for micronucleus frequency in peripheral blood of rats due to efficient splenic
6 scavenging of damaged erythrocytes—did not show an increase in micronucleated cells after
7 3 months of exposure to ST via drinking water (0, 125, 250, 500, 1,000, and 2,000 mg/L)
8 (Table D-2). Significant increases in the percent reticulocytes were seen in both male and female
9 rats suggesting that ST could have stimulated erythropoiesis in the bone marrow; however, the
10 absolute increases in the percentages were small compared to the vehicle control animals.

11 In male and female mice, there were no significant increases in micronucleated reticulocytes or
12 in micronucleated erythrocytes in either sex following 3 months of exposure to ST via drinking
13 water (0, 125, 250, 500, 1,000, and 2,000 mg/L) (Table D-3). A significant increase in the
14 percent reticulocytes was seen in male mice suggesting that ST could have stimulated
15 erythropoiesis in the bone marrow; however, the absolute increase was small compared to the
16 vehicle control group.

17 In addition to evaluating the potential for chromosomal damage, the potential for DNA damage
18 was assessed using the comet assay in the same animals in which micronucleus induction was
19 evaluated. DNA damage from exposure to ST was assessed in liver, ileum, and kidney cell
20 samples and in blood leukocytes (Table D-4, Table D-5). Significant increases in DNA damage,
21 measured as percent tail DNA, were observed in liver cells from male and female rats.
22 Significant increases in DNA damage were also observed in liver and ileum cells from male
23 mice. No increases in percent tail DNA were observed in female mice. Although cells from
24 kidney tissue were evaluated for male and female rats, and ileum tissue was evaluated from male
25 rats, results from these tissues were considered invalid due to unusually high percent tail DNA in
26 the vehicle control group.

1 Discussion

2 Tungsten was nominated to the National Toxicology Program (NTP) by the Center of Disease
3 Control and Prevention for toxicity and carcinogenicity studies due to inadequate data on effects
4 in humans and concern about potential human exposure via drinking water.^{89; 90} Tungstate
5 (WO_4^{2-}) is the most common of the naturally occurring forms of soluble tungsten; sodium
6 tungstate dihydrate (ST) was selected for these studies because it is the most water-soluble form
7 of tungstate.

8 In the current perinatal and 3-month rat study, no exposure-related effects were noted on
9 pregnancy status, maternal survival, or the number of dams that littered at any of the exposure
10 concentrations tested. There were significant decreases in the mean body weights of dams in the
11 1,000 and 2,000 mg/L groups (approximately 10% and 18%, respectively) at the end of lactation.
12 The mean body weights of pups (male and female combined) on postnatal day 21 in the
13 2,000 mg/L group were also significantly decreased by approximately 14%. There were no ST-
14 related findings in the testes or epididymides, or changes in sperm parameters in either rat or
15 mouse after 3 months of exposure. A previous study of Sprague Dawley rats exposed daily to ST
16 at 5 or 125 mg/kg/day by gavage for 70 days (including through mating, gestation, and lactation)
17 similarly found no reproductive effects.⁹¹

18 In the 3-month drinking water studies, the kidney was the major target organ of toxicity in both
19 rats and mice. In rats, renal tubule regeneration was increased in the male and female 1,000 and
20 2,000 mg/L groups; the increases in the 2,000 mg/L groups were significant. Renal tubule
21 regeneration was distinct from chronic progressive nephropathy (CPN), which had similar
22 incidences and severities between exposed and vehicle control groups. Renal tubule regeneration
23 was also significantly increased in 1,000 and 2,000 mg/L male mice. In female mice, although
24 there was a significant positive trend, there were no pairwise significant differences between
25 vehicle control animals and exposed animals, indicating sex-related differences in ST sensitivity
26 in mice.

27 The urine xanthine/creatinine ratios were significantly increased in all the male and female rat
28 groups at 13 weeks. This finding was not unexpected as ST has been shown to antagonize the
29 normal metabolic action of molybdate in its role as a cofactor for several enzymes, including
30 xanthine oxidase.^{92; 93; 2} Xanthine oxidase catalyzes the oxidation of xanthine to uric acid; thus,
31 inhibition of this enzyme can lead to increases of xanthine in the urine.²

32 Serum insulin concentrations were also measured in the perinatal and 3-month rat study. Serum
33 insulin concentrations were significantly decreased in the high-exposure male group whereas
34 serum glucose was unchanged. Studies have shown that ST has antidiabetic effects in rats in that
35 when administered to streptozotocin-induced (STZ) diabetic rats, serum glucose normalizes
36 without changes to the serum insulin concentrations.^{94; 95} In these same studies, both glucose and
37 insulin concentrations were unchanged when ST was administered to healthy rats. These studies
38 suggest that in STZ diabetic rats, serum glucose is lowered in part due to ST's ability to restore
39 hepatic glucose metabolism by increasing the capacity of the liver to use glucose through
40 glycolysis and glycogenesis. The effects have previously been shown to occur through an insulin
41 receptor-independent pathway.⁹ Given the known effects of tungstate on glucose metabolism, the

1 finding of decreased insulin was unexpected; the reason for this finding in the current study is
2 not known.

3 Significant decreases noted in body weight gain in rat dams during the lactation phase, and
4 reductions in final mean body weight in weaned pups, informed the decision to lower the top
5 exposure concentration in rats to 1,000 mg/L in the chronic studies and to expose mice up to
6 2,000 mg/L.

7 In the perinatal and 2-year rat study, no exposure-related effects were noted on pregnancy status,
8 maternal survival, or the number of dams that littered at any of the exposure concentrations
9 tested (up to approximately 143 mg/kg/day). These findings are consistent with previous findings
10 in the literature, which showed no reproductive effects in Sprague Dawley rats exposed to ST at
11 5 or 125 mg/kg/day.⁹¹

12 At study termination, an approximately 22% reduction in mean body weight was observed in the
13 1,000 mg/L female rats compared to the vehicle control group. Mean body weights of all groups
14 of exposed males were within 10% of the vehicle control group. Mean water consumption was
15 reduced by approximately 16% in the 1,000 mg/L males; in all other groups it was within 10% of
16 the vehicle control group.

17 In the chronic studies, female rats in the 500 mg/L group had a significant increase in the
18 incidence of thyroid C-cell adenomas compared to the vehicle control group, and there was a
19 doubling of the incidence of C-cell adenomas in the 250 and 500 mg/L groups compared to that
20 seen in the vehicle control group. Although there was no significant difference between the
21 combined incidences of C-cell adenoma or carcinoma, the incidences in the 250 and 500 mg/L
22 groups were above the historical control range. The highest incidence of C-cell carcinomas was
23 found in the 1,000 mg/L group, and there was an earlier onset of C-cell neoplasms in exposed
24 groups compared to the vehicle control group. No significant differences of C-cell hyperplasia
25 were observed in female rats between exposed groups and the vehicle control group, however,
26 nor were there any significant differences between exposed and vehicle control male rats in the
27 incidences of C-cell lesions. NTP concluded that there was equivocal evidence of carcinogenic
28 activity based on the incidences of C-cell adenoma or carcinoma (combined) in female rats; this
29 was driven primarily by the adenomas.

30 Similar to the 3-month studies, the kidney was confirmed as a target organ of toxicity in both
31 male and female rats following chronic exposure to ST. Suppurative inflammation of the renal
32 tubules was significantly increased in the 1,000 mg/L males and females. Renal tubule
33 regeneration was significantly increased in the 1,000 mg/L females but was not observed in male
34 rats. By the end of the chronic study, the widespread and severe CPN present in male rats made it
35 impossible to distinguish changes of renal tubule regeneration from those seen with CPN. Given
36 the findings in the 3-month study and the findings in the female rats, however, the renal tubular
37 epithelium is likely a target of ST in both male and female rats. Despite the observation of renal
38 tubule regeneration at 3 months, there did not appear to be a progression to renal tubule
39 neoplasms at 2 years. Consistent with the nephrotoxicity, kidney tungsten concentration
40 increased with the exposure concentration and the kidney/plasma ratios were higher than 1.0 at
41 all exposure concentrations and time points demonstrating retention of tungsten in the kidney.

42 Exposure to ST for 2 years also resulted in a significantly increased incidence of atypical
43 epithelial hyperplasia in the uterus of female rats in the 500 mg/L group. The incidences of

1 atypical hyperplasia were greater than that of the vehicle control group, but not significantly so,
2 in the 250 and 1,000 mg/L groups. Atypical epithelial hyperplasia in the uterus is considered a
3 preneoplastic finding, but neither the incidences of adenoma nor adenocarcinoma of the uterus
4 were increased in exposed groups compared to the vehicle control group. Therefore, the
5 relationship between atypical hyperplasia and exposure to ST in this study is unknown.

6 In mice, toxicological effects included moderate reductions in final mean body weight and water
7 consumption for males in the 2,000 mg/L group. There were no exposure-related reductions in
8 mean body weights or water consumption for females in any of the ST exposed groups. Daily
9 estimated ST consumption indicated that the consumed doses of females were approximately
10 70% of males at a given exposure concentration. Absolute kidney weights were significantly
11 increased in the 500 and 1,000 mg/L males compared to the vehicle control group at 3 months,
12 but by 18 months, there were no significant pairwise differences in kidney weights in either
13 males or females compared to the vehicle control groups.

14 Histologically, the kidney was a target of ST administration in mice after chronic exposure.
15 Similar to what was seen at 3 months, renal tubule regeneration was recorded in significantly
16 increased incidences with increasing exposure. Three renal tubule neoplasms were observed in
17 exposed male mice; no renal tubule neoplasms occurred in vehicle control mice, and no renal
18 tubule neoplasms were found in female mice. Step sectioning was conducted on the kidney, but
19 no additional neoplasms were observed. Although there was no statistical difference between the
20 incidences of renal tubule neoplasms in exposed groups versus the vehicle control group, they
21 are uncommon in mice, with only three males out of a total of 687 control male mice in the
22 historical database having renal tubule neoplasms. Given that renal tubule regeneration was
23 observed in the 3-month study, and there was a limited exposure response—with one renal
24 tubule adenoma present in the mid-exposure group and two renal tubule carcinomas present in
25 the high-exposure group—the occurrences of the renal tubule adenoma or carcinoma (combined)
26 were considered equivocal evidence of carcinogenic activity. Consistent with this, the
27 kidney/plasma tungsten ratios of higher than 1 suggest retention of tungsten in the kidney.

28 Overall, the kidney was considered a major target organ of toxicity in rats and mice in the
29 subchronic and chronic studies. As indicated by the tissue burden data, tungsten accumulated in
30 the kidney in an exposure concentration-dependent manner. Some species differences in tungsten
31 accumulation in the kidney was observed in these studies. In mice, kidney/plasma ratios (1.4–
32 4.4) remained similar regardless of the exposure duration with no observed sex difference;
33 however, in rats, the ratios increased with exposure duration, with male rats showing a higher
34 ratio (1.83–2.51 at 3 months to 12.63–24.26 at 18 months) than female rats (1.17–2.06 at 3
35 months to 3.99–7.11 at 18 months). The general trend was toward a decreasing kidney/plasma
36 ratio with increasing exposure concentration in both rats and mice. Accumulation of tungsten in
37 the kidney corresponded to toxicity as evidenced by renal tubule inflammation and regeneration
38 in male and female rats and mice in the subchronic and chronic studies. In rats, renal tubule
39 changes were significant at the highest exposure concentration (1,000 mg/L), but mice receiving
40 higher exposure concentrations than rats (up to 2,000 mg/L) exhibited a distinct effect of
41 exposure concentration-dependent renal toxicity. Additionally, renal tubule adenomas and
42 carcinomas were noted in the chronic studies in male mice.

43 Although human biomonitoring data on tungsten concentrations in serum are relatively limited in
44 the literature, the plasma tungsten concentration in the current studies following exposure of

1 male rats (2.64 $\mu\text{g}/\text{mL}$) and male mice (1.27 $\mu\text{g}/\text{mL}$) to 250 ppm ST are approximately 18,000
2 and 8,500 times higher, respectively, compared to humans (arithmetic mean 0.15 $\mu\text{g}/\text{L}$ or
3 0.00015 $\mu\text{g}/\text{mL}$, n= 290).⁹⁶ Several studies have evaluated urinary tungsten concentrations in
4 humans (summarized in Lemus et al.⁹⁷). Urinary tungsten concentrations in male rat
5 (104.37 $\mu\text{g}/\text{mg}$ creatinine) and male mice (278.92 $\mu\text{g}/\text{mg}$ creatinine) exposed to 250 ppm ST in
6 these studies are >1,000,000 times the urinary concentrations reported by a National Health and
7 Nutrition Examination Survey program (2015–2016) (geometric mean 0.076 $\mu\text{g}/\text{g}$ creatinine or
8 0.000076 $\mu\text{g}/\text{mg}$ creatinine) based on a representative sample of 3,057 individuals.⁹⁸

9 Other nonneoplastic lesions observed in the chronic mouse study associated with exposure to ST
10 included pigment in the cecum, germinal epithelium degeneration in the testes, hypercellularity
11 of the bone marrow in males, and extramedullary hematopoiesis in the spleen of females. Hence,
12 there was evidence of systemic exposure to ST following oral administration in the drinking
13 water based on the tissue distribution data and on toxicity in multiple tissues.

1 **Conclusions**

- 2 Under the conditions of these 2-year drinking water studies, there was *no evidence of*
3 *carcinogenic activity* of sodium tungstate dihydrate (ST) in male Hsd:Sprague Dawley[®] SD[®] rats
4 at exposure concentrations of 250, 500, or 1,000 mg/L. There was *equivocal evidence of*
5 *carcinogenic activity* of ST in female Hsd:Sprague Dawley[®] SD[®] rats based on increased
6 incidences of C-cell adenoma or carcinoma (combined) of the thyroid gland.
- 7 There was *equivocal evidence of carcinogenic activity* of ST in male B6C3F1/N mice based on
8 the occurrences of renal tubule adenoma or carcinoma (combined) in exposed animals. There
9 was *no evidence of carcinogenic activity* of ST in female B6C3F1/N mice at exposure
10 concentrations of 500, 1,000, or 2,000 mg/L.
- 11 Exposure to ST in drinking water caused increased incidences of nonneoplastic lesions in the
12 kidney of male and female rats and mice, in the uterus of female rats, in the testes and bone
13 marrow of male mice, and in the spleen of female mice.

1 **References**

- 2 1. Higgins ES, Richert DA, Westerfield WW. Competitive role of tungsten in molybdenum
3 nutrition. *Fed Proc* 1956; 15:274-275.
- 4 2. Higgins ES, Richert DA, Westerfield WW. Molybdenum deficiency and tungstate inhibition
5 studies. *J Nutr.* 1956; 59(539-559).
- 6 3. Johnson JL, Cohen HJ, Rajagopalan KV. Molecular basis of the biological function of
7 molybdenum-free sulfite oxidase from livers of tungsten-treated rats. *J Biol Chem.* 1974;
8 249:5046-5055.
- 9 4. Owen EC, Proudfoot R. The effect of tungstate ingestion on xanthine oxidase in milk and
10 liver. *Br J Nutr.* 1968; 22:331-340.
- 11 5. Hazardous Substances Data Bank (HSDB). Sodium tungstate CASRN: 13472-45-2. Bethesda,
12 MD: National Library of Medicine; 2009.
13 https://hero.epa.gov/hero/index.cfm/reference/details/reference_id/2346118
- 14 6. Agency for Toxic Substances & Disease Registry (ATSDR). Toxicological profile for
15 tungsten. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service;
16 2005. <https://www.atsdr.cdc.gov/ToxProfiles/tp.asp?id=806&tid=157>
- 17 7. Agency for Toxic Substances & Disease Registry (ATSDR). Addendum to the toxicological
18 profile for tungsten. Atlanta, GA: U.S. Department of Health and Human Services, Public Health
19 Service; 2015. https://www.atsdr.cdc.gov/toxprofiles/Tungsten_Addendum_508.pdf
- 20 8. Altirriba J, Barbera A, Del Zotto H, Badal B, Piquer S, Sanchez-Pla A, Gagliardino JJ, Gomis
21 R. Molecular mechanisms of tungstate-induced pancreatic plasticity: A transcriptomics
22 approach. *BMC Genomics.* 2009; 10:406. 10.1186/1471-2164-10-406
- 23 9. Dominguez JE, Munoz MC, Zafra D, Sanchez-Perez I, Baque S, Caron M, Mercurio C,
24 Barbera A, Perona R, Gomis R et al. The antidiabetic agent sodium tungstate activates glycogen
25 synthesis through an insulin receptor-independent pathway. *J Biol Chem.* 2003; 278(44):42785-
26 42794. 10.1074/jbc.M308334200
- 27 10. Bertinat R, Nualart F, Li X, Yanez AJ, Gomis R. Preclinical and clinical studies for sodium
28 tungstate: Application in humans. *J Clin Cell Immunol.* 2015; 6(1):pii: 285. 10.4172/2155-
29 9899.1000285
- 30 11. National Institute for Occupational Safety and Health (NIOSH). Tungsten. U.S. Department
31 of Health and Human Services, Public Health Service, Centers for Disease Control and
32 Prevention, National Institute for Occupational Safety and Health; 2011.
33 <https://www.cdc.gov/niosh/pel88/7440-33s.html>
- 34 12. U.S. Environmental Protection Agency (USEPA). Technical fact sheet-tungsten. 2017.
35 [https://www.epa.gov/sites/production/files/2017-10/documents/ffro_ecfactsheet_tungsten_9-15-
36 17_508.pdf](https://www.epa.gov/sites/production/files/2017-10/documents/ffro_ecfactsheet_tungsten_9-15-17_508.pdf)

- 1 13. McDonald JD, Weber WM, Marr R, Kracko D, Khain H, Arimoto R. Disposition and
2 clearance of tungsten after single-dose oral and intravenous exposure in rodents. *Journal of*
3 *Toxicology and Environmental Health Part A*. 2007; 70(10):829-836.
4 10.1080/15287390701211762
- 5 14. Weber WM, Marr R, Kracko D, Gao Z, McDonald JD, Chearnaigh KU. Disposition of
6 tungsten in rodents after repeat oral and drinking water exposures. *Toxicological &*
7 *Environmental Chemistry*. 2008; 90(3):445-455. 10.1080/02772240701565891
- 8 15. Lagarde F, Leroy M. Metabolism and toxicity of tungsten in humans and animals. *Met Ions*
9 *Biol Syst*. 2002; 39:741-759.
- 10 16. Bowen HJM. The elemental content of human diets and excreta In: Bowen HJM, Berrow
11 ML, Burton JD, Cawse PA, Patterson DSP, Statham PJ, Ure AM, editors. *Environmental*
12 *Chemistry, Vol 2 A Review of the Literature Published up to mid-1980*. London: The Royal
13 Society of Chemistry, Burlington House; 1982. p. 70-93.
- 14 17. Friberg L, Nordberg GR. *Handbook on the Toxicology of Metals*. New York, NY: Elsevier
15 North Holland; 1979.
- 16 18. Fernandez-Alvarez J, Zapatero J, Pifiol C. Acute oral and intravenous toxicity of sodium
17 tungstate: A potential agent to treat diabetes mellitus. Abstracts of the Symposium on the
18 Insulinomimetic Effects of Metal Ions: Potential Therapy for Diabetes Mellitus Sitges, Spain.
19 2000a:24.
- 20 19. Nadeenko VG. Maximum permissible concentrations of tungsten in water basins. *Hyg Sanit*.
21 1966; 31:197-203
- 22 20. McCain WC, Crouse LC, Bazar MA, Roszell LE, Leach GJ, Middleton JR, Reddy G.
23 Subchronic oral toxicity of sodium tungstate in Sprague-Dawley rats. *Int J Toxicol*. 2015;
24 34(4):336-345. 10.1177/1091581815585568
- 25 21. Hazardous Substances Data Bank (HSDB). Sodium tungstate. HSDB No. 5057. Bethesda,
26 MD: National Library of Medicine (NLM); 2002.
- 27 22. Domingo JL. Vanadium and tungsten derivatives as antidiabetic agents: A review of their
28 toxic effects. *Biol Trace Elem Res*. 2002; 88(2):97-112. 10.1385/BTER:88:2:097
- 29 23. Fernandez-Alvarez J, Zapatero J, Pinol C. Subacute and subchronic sodium tungstate toxicity
30 studies. Abstracts of the Symposium on the Insulinomimetic Effects of Metal Ions: Potential
31 Therapy for Diabetes Mellitus Sitges, Spain. 2000b:25.
- 32 24. Registry of Toxic Effects of Chemical Substances (RTECS). Tungstic acid, disodium salt.
33 RTECS No. YO7875000. Last updated in December 2000. Cincinnati, OH: National Institute for
34 Occupational Safety and Health (NIOSH); 2002.
- 35 25. Frawley RP, Smith MJ, White KL, Jr., Elmore SA, Herbert R, Moore R, Staska LM, Behl M,
36 Hooth MJ, Kissling GE et al. Immunotoxic effects of sodium tungstate dihydrate on female
37 B₆C₃F₁/N mice when administered in drinking water. *J Immunotoxicol*. 2016; 13(5):666-675.
38 10.3109/1547691X.2016.1154118

- 1 26. Schroeder HA, Mitchener M. Life-term studies in rats: Effects of aluminum, barium,
2 beryllium, and tungsten. *J Nutr.* 1975; 105(4):421-427. 10.1093/jn/105.4.421
- 3 27. Sato K, Ichimasa M, Miyahara K, Shiomi M, Nishimura Y, Ichimasa Y. Radioprotective
4 effects of sodium tungstate on hematopoietic injury by exposure to ⁶⁰Co gamma-rays in Wistar
5 rats. *Journal of Radiation Research.* 1999; 40(2):101-113. 10.1269/jrr.40.101
- 6 28. American Conference of Government Industrial Hygienists (ACGIH). Tungsten. Threshold
7 limit values for chemical substances and physical agents and biological exposure indices.
8 Cincinnati, OH; 1991.
- 9 29. Wide M. Effect of short-term exposure to five industrial metals on the embryonic and fetal
10 development of the mouse. *Environ Res.* 1984; 33(1):47-53.
- 11 30. Nadeenko VG, Lenchenko VG, Oschepkova AN. New data for standardization of tungsten
12 and molybdenum in their separate and simultaneous presence in water bodies. *Gig Sanit.* 1978;
13 3:7-11.
- 14 31. McInturf SM, Bekkedal MY, Wilfong E, Arfsten D, Chapman G, Gunasekar PG. The
15 potential reproductive, neurobehavioral and systemic effects of soluble sodium tungstate
16 exposure in Sprague-Dawley rats. *Toxicol Appl Pharmacol.* 2011; 254(2):133-137.
17 10.1016/j.taap.2010.04.021
- 18 32. DiPaolo JA, Casto BC. Quantitative studies of in vitro morphological transformation of
19 Syrian hamster cells by inorganic metal salts. *Cancer Res.* 1979; 39(3):1008-1013.
- 20 33. Larramendy ML, Popescu NC, DiPaolo JA. Induction by inorganic metal salts of sister
21 chromatid exchanges and chromosome aberrations in human and Syrian hamster cell strains.
22 *Environmental and Molecular Mutagenesis.* 1981; 3(6):597-606. 10.1002/em.2860030602
- 23 34. Barbera A, Gomis RR, Prats N, Rodriguez-Gil JE, Domingo M, Gomis R, Guinovart JJ.
24 Tungstate is an effective antidiabetic agent in streptozotocin-induced diabetic rats: A long-term
25 study. *Diabetologia.* 2001; 44(4):507-513. 10.1007/s001250100479
- 26 35. Maronpot RR, Boorman GA. Interpretation of rodent hepatocellular proliferative alterations
27 and hepatocellular tumors in chemical safety assessment. *Toxicol Pathol.* 1982; 10(2):71-78.
28 10.1177/019262338201000210
- 29 36. Boorman GA, Montgomery CA, Jr., Eustis SL, Wolfe MJ, McConnell EE, Hardisty JF.
30 Quality assurance in pathology for rodent carcinogenicity studies In: Milman HA, Weisburger
31 EK, editors. *Handbook of Carcinogen Testing.* Park Ridge, NJ: Noyes Publications; 1985. p.
32 345-357.
- 33 37. Boorman GA, Haseman JK, Waters MD, Hardisty JF, Sills RC. Quality review procedures
34 necessary for rodent pathology databases and toxicogenomic studies: The National Toxicology
35 Program experience. *Toxicol Pathol.* 2002; 30(1):88-92. 10.1080/01926230252824752
- 36 38. Brix AE, Hardisty JF, McConnell EE. Combining neoplasms for evaluation of rodent
37 carcinogenesis studies In: Hsu C-H, Stedeford T, editors. *Cancer Risk Assessment.* Hoboken,
38 NJ: John Wiley & Sons, Inc.; 2010. p.

- 1 39. Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *J Am Stat*
2 *Assoc.* 1958; 53(282):457-481. 10.2307/2281868
- 3 40. Tarone RE. Tests for trend in life table analysis. *Biometrika.* 1975; 62(3):679-690.
- 4 41. Cox DR. Regression models and life-tables. *Journal of Royal Statistical Society Series B.*
5 1972; 34(2):187-202.
- 6 42. Bailer AJ, Portier CJ. Effects of treatment-induced mortality and tumor-induced mortality on
7 tests for carcinogenicity in small samples. *Biometrics.* 1988; 44(2):417-431.
- 8 43. Piegorsch W, Bailer AJ. *Statistics for environmental biology and toxicology.* Section 6.3.2.
9 London, England: CRC Press; 1997.
- 10 44. Portier CJ, Bailer AJ. Testing for increased carcinogenicity using a survival-adjusted quantal
11 response test. *Fundam Appl Toxicol.* 1989; 12(4):731-737.
- 12 45. Portier CJ, Hedges JC, Hoel DG. Age-specific models of mortality and tumor onset for
13 historical control animals in the National Toxicology Program's carcinogenicity experiemnts.
14 *Cancer Res.* 1986; 46(9):4372-4378.
- 15 46. Bieler GS, Williams RL. Ratio estimates, the delta method, and quantal response tests for
16 increased carcinogenicity. *Biometrics.* 1993; 49(3):793-801.
- 17 47. Nam JM. A simple approximation for calculating sample sizes for detecting linear trend in
18 proportions. *Biometrics.* 1987; 43(3):701-705.
- 19 48. Rao JN, Scott AJ. A simple method for the analysis of clustered binary data. *Biometrics.*
20 1992; 48(2):577-585.
- 21 49. Fung KY, Krewski D, Rao JN, Scott AJ. Tests for trend in developmental toxicity
22 experiments with correlated binary data. *Risk Anal.* 1994; 14(4):639-648.
- 23 50. Gart JJ, Chu KC, Tarone RE. Statistical issues in interpretation of chronic bioassay tests for
24 carcinogenicity. *J Natl Cancer Inst.* 1979; 62(4):957-974.
- 25 51. Dixon WJ, Massey FJ, Jr. *Introduction to Statistical Analysis.* New York, NY: McGraw-Hill
26 Book Company; 1957.
- 27 52. Tukey J. Easy summaries-numerical and graphical. *Exploratory Data Analysis.* Reading,
28 MA: Addison-Wesley; 1977. p. 43-44.
- 29 53. Dunnett CW. A multiple comparison procedure for comparing several treatments with a
30 control. *J Am Stat Assoc.* 1955; 50(272):1096-1121. 10.1080/01621459.1955.10501294
- 31 54. Williams DA. A test for differences between treatment means when several dose levels are
32 compared with a zero dose control. *Biometrics.* 1971; 27(1):103-117.
- 33 55. Williams DA. The comparison of several dose levels with a zero dose control. *Biometrics.*
34 1972; 28(2):519-531.

- 1 56. Shirley E. A non-parametric equivalent of Williams' test for contrasting increasing dose
2 levels of a treatment. *Biometrics*. 1977; 33(2):386-389.
- 3 57. Williams DA. A note on Shirley's nonparametric test for comparing several dose levels with
4 a zero-dose control. *Biometrics*. 1986; 42(1):183-186.
- 5 58. Dunn OJ. Multiple comparison using RANK sums. *Technometrics*. 1964; 6:241-252.
6 10.1080/00401706.1964.10490181
- 7 59. Jonckheere AR. A distribution-free k-sample test against ordered alternatives. *Biometrika*.
8 1954; 41(1-2):133-145. 10.1093/biomet/41.1-2.133
- 9 60. Hsu JC. The factor analytic approach to simultaneous inference in the general linear model. *J*
10 *Comput Graph Stat*. 1992; 1(2):151-168. 10.1080/10618600.1992.10477011
- 11 61. Davison AC, Hinkley DV. *Bootstrap Methods and Their Application*. Cambridge Univ.
12 Press.; 1997.
- 13 62. Datta S, Satten GA. Rank-sum tests for clustered data. *J Am Stat Assoc* 2005; 100:908-915.
14 10.1198/016214504000001583
- 15 63. Hommel G. A stagewise rejective multiple test procedure based on a modified Bonferroni
16 test. *Biometrika*. 1988; 75(2):383-386.
- 17 64. Haseman JK. Value of historical controls in the interpretation of rodent tumor data. *Drug Inf*
18 *J*. 1992; 26(2):191-200.
- 19 65. Haseman JK. Data analysis: Statistical analysis and use of historical control data. *Regul*
20 *Toxicol Pharmacol*. 1995; 21(1):52-59; discussion 81-56. 10.1006/rtp.1995.1009
- 21 66. Haseman JK, Rao GN. Effects of corn oil, time-related changes, and inter-laboratory
22 variability on tumor occurrence in control Fischer 344 (F344/N) rats. *Toxicol Pathol*. 1992;
23 20(1):52-60. 10.1177/019262339202000107
- 24 67. Code of Federal Regulations (CFR). 21(Part 58).
- 25 68. Miller JA, Miller EC. Ultimate chemical carcinogens as reactive mutagenic electrophiles In:
26 Hiatt HH, Watson JD, Winsten JA, editors. *Origins of Human Cancer*. Spring Harbor, NY: Cold
27 Spring Harbor Laboratory; 1977. p. 605-627.
- 28 69. Crawford BD. Perspectives on the somatic mutation model of carcinogenesis In: Mehlman
29 MA, Flamm WG, Lorentzen RJ, editors. *Advances in Modern Environmental Toxicology*
30 *Mechanisms and Toxicity of Chemical Carcinogens and Mutagens*. Princeton, NJ: Princeton
31 Scientific Publishing Co. Inc; 1985. p. 13-59.
- 32 70. Straus DS. Somatic mutation, cellular differentiation, and cancer causation. *J Natl Cancer*
33 *Inst*. 1981; 67:233-241.
- 34 71. Ashby J, Tennant RW. Definitive relationships among chemical structure, carcinogenicity
35 and mutagenicity for 301 chemicals tested by the US NTP. *Mutat Res*. 1991; 257(3):229-306.

- 1 72. Tennant RW, Margolin BH, Shelby MD, Zeiger E, Haseman JK, Spalding J, Caspary W,
2 Resnick M, Stasiewicz S, Anderson B. Prediction of chemical carcinogenicity in rodents from in
3 vitro genetic toxicity assays. *Science* 1987; 236(4804):933-941.
- 4 73. Zeiger E, Haseman JK, Shelby MD, Margolin BH, Tennant RW, Holden H. Evaluation of
5 four in vitro genetic toxicity tests for predicting rodent carcinogenicity: Confirmation of earlier
6 results with 41 additional chemicals. *Environ Mol Mutag*. 1990; 16(S18):1-14.
- 7 74. Heddle JA, Hite M, Kirkhart B, Mavournin K, MacGregor JT, Newell GW, Salamone MF.
8 The induction of micronuclei as a measure of genotoxicity: A report of the U.S. environmental
9 protection agency Gene-Tox program. *Mutat Res/Rev Mutat Res*. 1983; 123(1):61-118.
10 [https://doi.org/10.1016/0165-1110\(83\)90047-7](https://doi.org/10.1016/0165-1110(83)90047-7)
- 11 75. Schmid W. The micronucleus test. *Mutation Research/Environmental Mutagenesis and*
12 *Related Subjects*. 1975; 31(1):9-15. [https://doi.org/10.1016/0165-1161\(75\)90058-8](https://doi.org/10.1016/0165-1161(75)90058-8)
- 13 76. Shelby MD, Erexson GL, Hook GJ, Tice RR. Evaluation of a three-exposure mouse bone
14 marrow micronucleus protocol: Results with 49 chemicals. *Environ Mol Mutagen*. 1993; 21:160-
15 179.
- 16 77. Shelby MD, Witt KL. Comparison results from mouse bone marrow chromosome aberration
17 and micronucleus tests. *Environ Mol Mutagen*. 1995; 25:302-313.
- 18 78. Witt KL, Knapton A, Wehr CM, Hook GJ, Mirsalis J, Shelby MD, MacGregor JT.
19 Micronucleated erythrocyte frequency in peripheral blood of B6C3F1 mice from short-term,
20 prechronic, and chronic studies of the NTP Carcinogenesis Bioassay Program. *Environ Mol*
21 *Mutagen*. 2000; 36:163-194.
- 22 79. Organisation for Economic Cooperation and Development (OECD). OCED guideline for the
23 testing of chemicas. In vivo mammalian alkaline comet assay. Paris, France: OECD Publishing;
24 2014. Testing Guideline 489. [https://www.oecd.org/env/test-no-489-in-vivo-mammalian-](https://www.oecd.org/env/test-no-489-in-vivo-mammalian-alkaline-comet-assay-9789264264885-en.htm)
25 [alkaline-comet-assay-9789264264885-en.htm](https://www.oecd.org/env/test-no-489-in-vivo-mammalian-alkaline-comet-assay-9789264264885-en.htm)
- 26 80. Brendler-Schwaab S, Hartmann A, Pfuhrer S, Speit G. The in vivo comet assay: Use and
27 status in genotoxicity testing. *Mutagenesis*. 2005; 20(4):245-254. 10.1093/mutage/gei033
- 28 81. Burlinson B, Tice RR, Speit G, Agurell E, Brendler-Schwaab SY, Collins AR, Escobar P,
29 Honma M, Kumaravel TS, Nakajima M et al. Fourth International Workgroup on Genotoxicity
30 testing: Results of the in vivo comet assay workgroup. *Mutat Res*. 2007; 627(1):31-35.
31 10.1016/j.mrgentox.2006.08.011
- 32 82. Collins AR. The comet assay for DNA damage and repair: Principles, applications, and
33 limitations. *Mol Biotechnol*. 2004; 26(3):249-261. 10.1385/mb:26:3:249
- 34 83. Tice RR, Agurell E, Anderson D, Burlinson B, Hartmann A, Kobayashi H, Miyamae Y,
35 Rojas E, Ryu JC, Sasaki YF. Single cell gel/comet assay: Guidelines for in vitro and in vivo
36 genetic toxicology testing. *Environ Mol Mutagen*. 2000; 35(3):206-221. 10.1002/(sici)1098-
37 2280(2000)35:3<206::aid-em8>3.0.co;2-j
- 38 84. Hartmann A, Agurell E, Beevers C, Brendler-Schwaab S, Burlinson B, Clay P, Collins A,
39 Smith A, Speit G, Thybaud V et al. Recommendations for conducting the in vivo alkaline comet

- 1 assay. 4th International Comet Assay Workshop. *Mutagenesis*. 2003; 18(1):45-51.
2 10.1093/mutage/18.1.45
- 3 85. Pfuhrer S, Wolf HU. Detection of DNA-crosslinking agents with the alkaline comet assay.
4 *Environ Mol Mutagen*. 1996; 27(3):196-201. 10.1002/(sici)1098-2280(1996)27:3<196::Aid-
5 em4>3.0.Co;2-d
- 6 86. National Toxicology Program (NTP). TR-599: Pathology tables, survival and growth curves
7 from NTP short-term and long-term studies. Research Triangle Park, NC: U.S. Department of
8 Health and Human Services, National Institute of Environmental Health Sciences, National
9 Toxicology Program; 2020. <https://doi.org/10.22427/NTP-DATA-TR-599>
- 10 87. Everds NE, Snyder PW, Bailey KL, Bolon B, Creasy DM, Foley GL, Rosol TJ, Sellers T.
11 Interpreting stress responses during routine toxicity studies: A review of the biology, impact, and
12 assessment. *Toxicol Pathol*. 2013; 41(4):560-614. 10.1177/0192623312466452
- 13 88. Dixon D, Alison R, Bach U, Colman K, Foley GL, Harleman JH, Haworth R, Herbert R,
14 Heuser A, Long G et al. Nonproliferative and proliferative lesions of the rat and mouse female
15 reproductive system. *J Toxicol Pathol*. 2014; 27(3-4 Suppl):1s-107s. 10.1293/tox.27.1S
- 16 89. Centers for Disease Control and Prevention (CDC). Exposure to tungsten in three Nevada
17 communities. Centerse for Disease Control and Prevention study results. 2003.
18 <https://www.cdc.gov/nceh/clusters/Fallon/w-summary.pdf>
- 19 90. Centers for Disease Control and Prevention (CDC). Update: Community meeting on
20 February 6, 2003. Cross-sectional exposure assessment of environmental contaminants in
21 Churchill County, Nevada. Centers for Disease Control and Prevention study results. 2003.
22 <https://www.cdc.gov/nceh/clusters/fallon/factsheet.pdf>
- 23 91. McInturf SM, Bekkedal MY, Wilfong E, Arfsten D, Gunasekar PG, Chapman GD.
24 Neurobehavioral effects of sodium tungstate exposure on rats and their progeny. *Neurotoxicol*
25 *Teratol*. 2008; 30(6):455-461. 10.1016/j.ntt.2008.07.003
- 26 92. Cohen HJ, Drew RT, Johnson JL, Rajagopalan KV. Molecular basis of the biological
27 function of molybdenum: The relationship between sulfite oxidase and the acute toxicity of
28 bisulfite and SO₂. *Proc Natl Acad Sci U S A*. 1973; 70(12):3655-3659. 10.1073/pnas.70.12.3655
- 29 93. Cohen HJ, Johnson JL, Rajagopalan KV. Molecular basis of the biological function of
30 molybdenum. Developmental patterns of sulfite oxidase and xanthine oxidase in the rat. *Arch*
31 *Biochem Biophys*. 1974; 164(2):440-446. 10.1016/0003-9861(74)90053-8
- 32 94. Barberà A, Fernández-Alvarez J, Truc A, Gomis R, Guinovart JJ. Effects of tungstate in
33 neonatally streptozotocin-induced diabetic rats: mechanism leading to normalization of
34 glycaemia. *Diabetologia*. 1997; 40(2):143-149. 10.1007/s001250050655
- 35 95. Barberà A, Rodríguez-Gil JE, Guinovart JJ. Insulin-like actions of tungstate in diabetic rats.
36 Normalization of hepatic glucose metabolism. *J Biol Chem*. 1994; 269(31):20047-20053.
- 37 96. Bocca B, Mattei D, Pino A, Alimonti A. Italian network for human biomonitoring of metals:
38 preliminary results from two Regions. *Ann Ist Super Sanita*. 2010; 46(3):259-265.
39 10.4415/ann_10_03_06

- 1 97. Lemus R, Venezia CF. An update to the toxicological profile for water-soluble and sparingly
2 soluble tungsten substances. *Crit Rev Toxicol.* 2015; 45(5):388-411.
3 10.3109/10408444.2014.1003422
- 4 98. Centers for Disease Control and Prevention (CDC). Fourth national report on human
5 exposure to environmental chemicals. 2013. <https://www.cdc.gov/exposurereport/>
- 6 99. Witt KL, Livanos E, Kissling GE, Torous DK, Caspary W, Tice RR, Recio L. Comparison of
7 flow cytometry- and microscopy-based methods for measuring micronucleated reticulocyte
8 frequencies in rodents treated with nongenotoxic and genotoxic chemicals. *Mutat Res.* 2008;
9 649(1-2):101-113. 10.1016/j.mrgentox.2007.08.004
- 10 100. Dertinger SD, Camphausen K, Macgregor JT, Bishop ME, Torous DK, Avlasevich S,
11 Cairns S, Tometsko CR, Menard C, Muanza T et al. Three-color labeling method for flow
12 cytometric measurement of cytogenetic damage in rodent and human blood. *Environ Mol*
13 *Mutagen.* 2004; 44(5):427-435. 10.1002/em.20075
- 14 101. Kissling GE, Dertinger SD, Hayashi M, MacGregor JT. Sensitivity of the erythrocyte
15 micronucleus assay: Dependence on number of cells scored and inter-animal variability. *Mutat*
16 *Res.* 2007; 634(1-2):235-240. 10.1016/j.mrgentox.2007.07.010
- 17 102. Recio L, Kissling GE, Hobbs CA, Witt KL. Comparison of Comet assay dose-response for
18 ethyl methanesulfonate using freshly prepared versus cryopreserved tissues. *Environ Mol*
19 *Mutagen.* 2012; 53(2):101-113. 10.1002/em.20694

20

1 **Appendix A. Chemical Characterization and Dose**

2 **Formulation Studies**

3 **Table of Contents**

4	A.1. Procurement and Characterization of Sodium Tungstate Dihydrate	A-2
5	A.2. Preparation and Analysis of Dose Formulations.....	A-2
6		

7 **Tables**

8	Table A-1. Chromatography Systems Used in the Three-month and Two-year Studies of	
9	Sodium Tungstate Dihydrate.....	A-4
10	Table A-2. Preparation and Storage of Dose Formulations in the Three-month and Two-	
11	year Studies of Sodium Tungstate Dihydrate.....	A-4
12	Table A-3. Results of Analyses of Dose Formulations Administered to Rats in the	
13	Perinatal and Three-month Study of Sodium Tungstate Dihydrate	A-5
14	Table A-4. Results of Analyses of Dose Formulations Administered to Mice in the Three-	
15	month Study of Sodium Tungstate Dihydrate.....	A-7
16	Table A-5. Results of Analyses of Dose Formulations Administered to Rats in the	
17	Perinatal and Two-year Study of Sodium Tungstate Dihydrate	A-9
18	Table A-6. Results of Analyses of Dose Formulations Administered to Mice in the Two-	
19	year Study of Sodium Tungstate Dihydrate	A-11
20		

21 **Figures**

22	Figure A-1. Infrared Absorption Spectrum of Sodium Tungstate Dihydrate	A-13
23		

1 **A.1. Procurement and Characterization of Sodium Tungstate** 2 **Dihydrate**

3 Sodium tungstate dihydrate (ST) was procured from Sigma-Aldrich (St. Louis, MO) in two lots
4 (lot 12330JO and lot MKBG9975V). Lot 12330JO was obtained directly from Sigma-Aldrich
5 (St. Louis, MO), whereas lot MKBG9975V was produced by Sigma-Aldrich and obtained from
6 Government Scientific Source, Inc. (Reston, VA). Identity, purity, and stability analyses were
7 conducted by the analytical chemistry laboratory and study laboratory at Battelle (Columbus,
8 OH). Reports on analyses performed in support of the ST studies are on file at the National
9 Institute of Environmental Health Sciences.

10 ST is a white solid composed of fine crystals. The 3-month studies used lot 12330JO, which was
11 homogenized by mixing for 15 minutes and transferred to 1 L amber storage bottles. For the
12 2-year studies, the remainder of lot 12330JO was combined with lot MKBG9975V to create
13 lot 07072011 by mixing in a blender for 15 minutes.

14 The identities of the lots were confirmed using infrared spectroscopy (Figure A-1). The x-ray
15 diffraction patterns of both lots were in good agreement with the reference pattern. Purity
16 assessment with proton-induced x-ray emission identified magnesium (0.7–0.9 %) and aluminum
17 (approximately 0.3%) impurities in both lots. The average concentrations of tungsten and sodium
18 in lot 12330JO were 54.09% and 18.24%, respectively. Lot 07072011 had average tungsten and
19 sodium concentrations of 47.6% and 15.8%, respectively. Elemental analysis using inductively
20 coupled plasma atomic emission spectrometry yielded purities of approximately 99%, based on
21 weight percentages of tungsten (55.2–56.4%) and sodium (13.4–13.8%). Karl Fisher titration
22 yielded a water content of 9.5% for lot 12330JO and 10.0–10.3% for lot 07072011, slightly
23 lower than the anticipated 10.9%. Titration with lead nitrate indicated a purity of 97.6% for
24 lot 12330JO and 98.2% for lot 07072011. Ion chromatography (IC) with a suppressed
25 conductivity detector and liquid chromatography with an inductively coupled plasma (ICP) mass
26 spectrometer indicated a purity of 100% for both lots. Additional information on
27 chromatography systems used can be found in Table A-1.

28 Accelerated stability studies were conducted on samples of ST stored at 60°C, 25°C, 5°C, and
29 –20°C using IC with a suppressed conductivity detector. Stability was confirmed for at least
30 2 weeks when stored in sealed amber glass bottles at 25°C, 5°C, and –20°C. Given these
31 findings, bulk ST was stored in sealed amber glass bottles at 25°C. Periodic analyses of the bulk
32 chemical using IC with a suppressed conductivity detector were conducted during the 3-month
33 and 2-year studies by the study laboratory and confirmed that no degradation occurred.

34 **A.2. Preparation and Analysis of Dose Formulations**

35 The presence of tungsten and molybdenum in animal feed (NIH-07 and NTP-2000), tap water,
36 and deionized water used in the 3-month and 2-year studies were evaluated with ICP optical
37 emission spectrometry at Galbraith Laboratories, Inc. (Knoxville, TN). NIH-07 feed contained
38 approximately 2 ppm tungsten, and the concentration in NTP-2000 feed was at the detection
39 limit of the assay (0.80 ppm). Concentrations of tungsten in the tap water and deionized water,
40 and the concentration of molybdenum in all feed and water samples, were below the limits of
41 detection of the assay (0.20 to 0.80 ppm).

1 Stability analysis conducted on the 20 µg/mL (20 mg/mL) formulation found that the
2 formulations were stable when sealed and stored in Nalgene bottles for 42 days at 5°C and room
3 temperature (approximately 25°C). An animal room simulation was conducted using the
4 20 mg/mL formulation stored in a drinking water bottle filled near capacity with aliquots
5 periodically removed to simulate animal drinking. There was no significant loss in tungsten over
6 7 days at room temperatures.

7 Dose formulations of ST were prepared monthly (Table A-2). Formulations were prepared with
8 deionized water, except for the first two preparations used in the 3-month studies, which used tap
9 water instead. The 3-month studies used formulations of 0, 125, 250, 500, 1,000, and 2,000 mg/L
10 for both mice and rats. These formulations were prepared four times in the mouse study (May–
11 July 2009) and five times in the rat study (May–August 2009). The 2-year mouse study used 0,
12 500, 1,000, and 2,000 mg/L formulations prepared 27 times from January 2012 to January 2014.
13 The 2-year rat study used 0, 250, 500, and 1,000 mg/L formulations prepared 28 times from
14 December 2011 to January 2014. Formulations were determined to be homogeneous and of
15 appropriate concentration using IC with a suppressed conductivity detector. Stability was
16 confirmed for 42 days at room temperature.

17 Preadministration and postadministration (animal room) analysis of formulations was conducted
18 monthly throughout the 3-month studies (Table A-3, Table A-4). During the 2-year studies,
19 preadministration formulations were analyzed every 1–3 months, whereas postadministration
20 (animal room) formulation were analyzed every 6–8 months (Table A-5, Table A-6). All
21 preadministration formulations in the 3-month rat and mouse studies were within 10% of the
22 target concentration. In the 3-month mouse study, four postadministration samples were more
23 than 10% below the target concentration in postadministration samples collected from bottles
24 used to expose females and carboys, with the largest difference being 12.8% below the target
25 (Table A-4). Postadministration samples collected from bottles or carboys in the 3-month rat
26 study prepared in May 2009 (125 mg/L), July 2009 (500 mg/L), and August 2009 (2,000 mg/L)
27 were 12.3%, 11.5%, and 10.8% below the target concentrations, respectively. All
28 preadministration and postadministration samples in the 2-year studies were within 10% of the
29 target concentration.

1 **Table A-1. Chromatography Systems Used in the Three-month and Two-year Studies of Sodium**
 2 **Tungstate Dihydrate**

Chromatography	Detection System	Column	Mobile Phase
System A			
Ion chromatography	Suppressed conductivity (50 mA)	Dionex IonPac AS11-HC, 250 mm × 2 mm ID	20 mM sodium hydroxide, flow rate 0.4 mL/min
System B			
Ion chromatography	Suppressed conductivity (40°C)	Dionex IonPac AS11-HC, 250 mm × 4 mm ID	27 mM sodium hydroxide, flow rate 0.9 mL/min
System C			
Liquid chromatography	Inductively coupled plasma-mass spectrometer	IonPac AS11-HC, 250 × 2 mm ID	Approximately 14.4 mM sodium hydroxide, flow rate 0.4 mL/min

3 ID = internal diameter.

4 **Table A-2. Preparation and Storage of Dose Formulations in the Three-month and Two-year**
 5 **Studies of Sodium Tungstate Dihydrate**

Preparation

Stock solutions of sodium tungstate dihydrate were created by weighing an appropriate amount of lot 12330JO (3-month studies) or lot 07072011 (2-year studies) in a weighing container. The contents were transferred to a volumetric flask and rinsed with water to ensure complete transfer. Flasks were brought to volume with deionized water. In place of deionized water, tap water was used for the first two formulations in the 3-month studies. Dose formulations were prepared monthly throughout the 3-month and 2-year studies.

Chemical Lot Number

3-month: 12330JO

2-year: 07072011 (Sigma lot 12330JO and MKBG9975V)

Maximum Storage Time

42 days

Storage Conditions

Formulations were stored in sealed Nalgene bottles at 5°C or 25°C

Study Laboratory

Battelle (Columbus, OH)

1 **Table A-3. Results of Analyses of Dose Formulations Administered to Rats in the Perinatal and**
 2 **Three-month Study of Sodium Tungstate Dihydrate**

Date Prepared	Date Analyzed	Target Concentration (mg/L)	Determined Concentration (mg/L) ^a	Difference from Target (%)
May 11, 2009	May 12, 2009	125	118 ± 9	-5.6
		250	257 ± 2	2.7
		500	511 ± 12	2.2
		1,000	1,070 ± 20	6.7
		2,000	2,200 ± 10	10.0
	June 25, 2009 (bottle) ^b	125	123 ± 6	-1.9
		250	259 ± 2	3.7
		500	515 ± 9	3.0
		1,000	945 ± 53	-5.5
		2,000	1,890 ± 130	-5.5
	June 25, 2009 (carboy) ^c	125	110 ± 8	-12.3
		250	245 ± 8	-2.1
		500	485 ± 27	-2.9
		1,000	973 ± 19	-2.7
		2,000	2,120 ± 70	6.0
June 5, 2009	June 16, 2009	125	123 ± 3	-1.6
		250	243 ± 4	-2.7
		500	494 ± 4	-1.2
		1,000	1,000 ± NA ^d	0.0
		2,000	1,940 ± 40	-3.2
	July 24, 2009 (bottle) ^b	125	121 ± 4	-3.5
		250	245 ± 8	-1.9
		500	508 ± 11	1.6
		1,000	1,040 ± 10	3.7
		2,000	2,070 ± 20	3.7
	July 24, 2009 (carboy) ^c	125	127 ± 7	1.3
		250	248 ± NA ^e	-0.8
		500	523 ± NA ^e	4.5
		1,000	1,050 ± 10	4.7
		2,000	2,080 ± 30	4.2

Date Prepared	Date Analyzed	Target Concentration (mg/L)	Determined Concentration (mg/L) ^a	Difference from Target (%)		
July 31, 2009	August 3, 2009	125	118 ± 17	-5.6		
		250	254 ± 4	1.6		
		500	510 ± 13	2.0		
		1,000	1010 ± 80	1.3		
		2,000	1,950 ± 170	-2.5		
	September 22, 2009 (bottles) ^b	September 22, 2009 (bottles) ^b	125	117 ± 4	-6.7	
			250	243 ± 22	-2.7	
			500	442 ± 32	-11.5	
			1,000	1,050 ± 30	4.7	
			2,000	1,870 ± 130	-6.7	
		September 22, 2009 (carboy) ^c	September 22, 2009 (carboy) ^c	125	129 ± 8	3.2
				250	271 ± 8	8.5
				500	479 ± 24	-4.2
				1,000	1,030 ± 80	2.9
				2,000	1,920 ± 20	-3.8
August 27, 2009	August 31, 2009	125	114 ± 4	-8.5		
		250	244 ± 3	-2.4		
		500	467 ± 19	-6.5		
		1,000	964 ± 4	-3.6		
		2,000	1,890 ± 40	-5.7		
	September 30, 2009 (bottle) ^b	September 30, 2009 (bottle) ^b	125	114 ± 12	-8.5	
			250	240 ± 7	-4.0	
			500	486 ± 4	-2.9	
			1,000	1,010 ± 10	1.3	
			2,000	2,050 ± 20	2.3	
		September 30, 2009 (carboy) ^c	September 30, 2009 (carboy) ^c	125	126 ± 3	1.1
				250	263 ± 1	5.1
				500	522 ± 10	4.3
				1,000	1,020 ± 50	2.2
				2,000	1,780 ± 240	-10.8

1 NA = not applicable.

2 ^aAverage of triplicate analysis.

3 ^bAnimal room sample from the formulation remaining in the drinking water bottle.

4 ^cAnimal room sample from the formulation collected from the carboy.

5 ^dDuplicate sample analyzed with a precision of duplicates value of 0.96.

6 ^eDuplicate sample analyzed with a precision of duplicates value of 1.0.

1 **Table A-4. Results of Analyses of Dose Formulations Administered to Mice in the Three-month**
 2 **Study of Sodium Tungstate Dihydrate**

Date Prepared	Date Analyzed	Target Concentration (mg/L)	Determined Concentration (mg/L) ^a	Difference from Target (%)
May 11, 2009	May 12, 2009	125	118 ± 9	-5.6
		250	257 ± 2	2.7
		500	511 ± 12	2.2
		1,000	1,070 ± 20	6.7
		2,000	2,200 ± 10	10.0
	June 25, 2009 (bottle) ^b	125	123 ± 2	-1.9
		250	263 ± 3	5.1
		500	508 ± 3	1.7
		1,000	1,050 ± 10	4.7
		2,000	2,140 ± 30	7.0
	June 25, 2009 (carboy) ^c	125	128 ± 9	2.7
		250	269 ± 5	7.7
		500	530 ± 5	6.0
		1,000	1,090 ± 20	9.3
		2,000	2,190 ± 50	9.3
June 5, 2009	June 16, 2009	125	123 ± 3	-1.6
		250	243 ± 4	-2.7
		500	494 ± 4	-1.2
		1,000	1,000 ± NA ^d	0.0
		2,000	1,940 ± 40	-3.2
	July 24, 2009 (bottle) ^b	125	127 ± 4	1.9
		250	257 ± 1	2.9
		500	513 ± 18	2.6
		1,000	1,060 ± 0	6.0
		2,000	2,100 ± 10	5.0
	July 24, 2009 (carboy) ^c	125	128 ± 4	2.1
		250	265 ± 7	5.9
		500	537 ± 12	7.3
		1,000	1,100 ± 20	9.7
		2,000	2,110 ± 30	5.5

Date Prepared	Date Analyzed	Target Concentration (mg/L)	Determined Concentration (mg/L) ^a	Difference from Target (%)
July 31, 2009	August 3, 2009	125	118 ± 17	-5.6
		250	254 ± 4	1.6
		500	510 ± 13	2.0
		1,000	1,010 ± 80	1.3
		2,000	1,950 ± 170	-2.5
	September 1, 2009 (bottles, males) ^b	125	115 ± 6	-8.0
		250	239 ± 5	-4.3
		500	486 ± 8	-2.7
		1,000	981 ± 17	-1.9
		2,000	1,850 ± 80	-7.7
	September 1, 2009 (bottles, females) ^b	125	109 ± 7	-12.8
		250	220 ± 5	-12.0
		500	456 ± 17	-8.8
		1,000	901 ± 6	-9.9
		2,000	1,770 ± 30	-11.3
	September 1, 2009 (carboy) ^c	125	112 ± 3	-10.1
		250	241 ± 2	-3.7
		500	494 ± 5	-1.3
1,000		1,000 ± 10	0.4	
2,000		1,920 ± 60	-4.0	

1
2
3
4

^aAverage of triplicate analysis.

^bAnimal room sample from the formulation remaining in the drinking water bottle.

^cAnimal room sample from the formulation collected from the carboy.

^dDuplicate sample analyzed with a precision of duplicates value of 0.96.

1 **Table A-5. Results of Analyses of Dose Formulations Administered to Rats in the Perinatal and**
 2 **Two-year Study of Sodium Tungstate Dihydrate**

Date Prepared	Date Analyzed	Target Concentration (mg/L)	Determined Concentration (mg/L) ^a	Difference from Target (%)
December 12, 2011	December 15, 2011	250	246 ± 6 / 247 ± 2	-1.7 / -1.1
		500	495 ± 3 / 483 ± 13	-0.9 / -3.3
		1,000	963 ± 10 / 945 ± 17	-3.7 / -3.7
	January 19, 2012 ^b (bottle)	250	236 ± 6	-5.7
		500	486 ± 2	-2.8
		1,000	998 ± 13	-0.2
	January 19, 2012 ^c (carboy)	250	239 ± 7	-4.3
		500	485 ± 3	-3.1
		1,000	967 ± 6	-3.3
January 5, 2012	January 6, 2012	250	238 ± 10 / 243 ± 5	-4.8 / -2.7
		500	473 ± 28 / 471 ± 24	-5.3 / -5.7
		1,000	918 ± 57 / 960 ± 32	-8.2 / -4.0
March 29, 2012	March 29, 2012	250	245 ± 1 / 238 ± 3	-2.0 / -4.7
		500	480 ± 3 / 479 ± 1	-4.1 / -4.1
		1,000	957 ± 3 / 962 ± 3	-4.3 / -3.8
May 22, 2012	May 22, 2012	250	242 ± 4 / 242 ± 1	-3.1 / -3.2
		500	490 ± 9 / 477 ± 7	-2.0 / -4.6
		1,000	958 ± 3 / 957 ± 16	-4.2 / -4.3
August 15, 2012	August 20, 2012	250	236 ± 1 / 249 ± 1	-5.6 / -0.4
		500	508 ± 0 / 504 ± 8	1.6 / 0.9
		1,000	997 ± 13 / 1,000 ± 10	0.0 / -0.8
	September 25, 2012 ^b (bottle)	250	233 ± 7	-6.7
		500	480 ± 8	-4.1
		1,000	1,000 ± 20	0.0
	September 25, 2012 ^c (carboy)	250	240 ± 8	-4.1
		500	498 ± 0	-0.4
		1,000	1,010 ± 10	0.7
October 11, 2012	October 14, 2012	250	237 ± 15 / 256 ± 6	-5.1 / 2.4
		500	490 ± 5 / 499 ± 4	-1.9 / -0.3
		1,000	998 ± 3 / 1,050 ± 20	-0.2 / 5.3
December 4, 2012	December 6, 2012	250	240 ± 10 / 253 ± 3	-4.0 / 1.1
		500	515 ± 8 / 538 ± 6	3.1 / 7.5
		1,000	1,050 ± 20 / 1,060 ± 10	5.0 / 5.7

Date Prepared	Date Analyzed	Target Concentration (mg/L)	Determined Concentration (mg/L) ^a	Difference from Target (%)
February 26, 2013	February 27, 2012	250	231 ± 8 / 238 ± 3	-7.6 / -4.9
		500	478 ± 3 / 476 ± 7	-4.5 / 4.8
		1,000	970 ± 22 / 950 ± 17	-5.0
	April 9, 2012 ^b (bottle)	250	233 ± 1	-6.7
		500	482 ± 3	-3.7
		1,000	953 ± 18	-4.7
	April 9, 2012 ^c (carboy)	250	241 ± 1	-3.6
		500	468 ± 3	-6.5
		1,000	945 ± 18	-5.5
April 25, 2013	April 26, 2013	250	240 ± 5 / 250 ± 7	-4.1 / -0.1
		500	527 ± 23 / 494 ± 17	5.4 / -1.3
		1,000	995 ± 9 / 988 ± 6	-0.5 / -1.2
July 16, 2013	July 18, 2013	250	258 ± 10 / 240 ± 14	3.1 / -4.0
		500	479 ± 17 / 509 ± 21	-4.1 / 1.7
		1,000	955 ± 71 / 970 ± 10	-4.5 / -3.0
September 10, 2013	September 14, 2013	250	228 ± 5 / 227 ± 6	-8.9 / -9.1
		500	454 ± 8 / 501 ± 23	-9.1 / 0.3
		1,000	1,090 ± 20 / 1,080 ± 0	8.7 / 8.0
	October 23, 2012 ^b (bottle)	250	228 ± 12	-8.8
		500	491 ± 5	-1.8
		1,000	1,070 ± 10	6.7
	October 23, 2012 ^c (carboy)	250	246 ± 4	-1.7
		500	485 ± 16	-2.9
		1,000	1,060 ± 30	6.0
December 4, 2013	December 6, 2013	250	232 ± 9 / 256 ± 12	-7.3 / 2.5
		500	495 ± 8 / 480 ± 21	-1.1 / -4.1
		1,000	973 ± 13 / 900 ± 6.9	-2.7 / -10.0

1 Values on either side of the / represent multiple formulations of the same dose prepared on that date.

2 ^aAverage of triplicate analysis.

3 ^bAnimal room sample from the formulation remaining in the drinking water bottle.

4 ^cAnimal room sample from the formulation collected from the carboy.

1 **Table A-6. Results of Analyses of Dose Formulations Administered to Mice in the Two-year Study**
 2 **of Sodium Tungstate Dihydrate**

Date Prepared	Date Analyzed	Target Concentration (mg/L)	Determined Concentration (mg/L) ^a	Difference from Target (%)
January 5, 2012	January 6, 2012	500	473 ± 28 / 471 ± 24	-5.3 / -5.7
		1,000	918 ± 57 / 960 ± 32	-8.2 / -4.0
		2,000	1,840 ± 60	-8.2
	February 16, 2012 ^b (bottle)	500	476 ± 26	-4.7
		1,000	938 ± 57	-6.2
		2,000	2,020 ± 30	1.0
	February 16, 2012 ^c (carboy)	500	456 ± 30	-8.8
		1,000	987 ± 32	-1.3
		2,000	2,030 ± 90	1.7
March 29, 2012	March 29, 2012	500	480 ± 3 / 479 ± 1	-4.1 / -4.1
		1,000	957 ± 3 / 962 ± 3	-4.3 / -3.8
		2,000	1,830 ± 40	-8.7
May 22, 2012	May 22, 2012	500	490 ± 9 / 477 ± 7	-2.0 / -4.6
		1,000	958 ± 3 / 957 ± 16	-4.2 / -4.3
		2,000	1,940 ± 10	-3.0
August 15, 2012	August 20, 2012	500	508 ± 0 / 504 ± 8	1.6 / 0.9
		1,000	997 ± 13 / 1,000 ± 10	0.0 / -0.8
		2,000	1,980 ± 10	-0.8
	September 25, 2012 ^b (bottle)	500	494 ± 22	-1.3
		1,000	1,020 ± 10	2.0
		2,000	2,070 ± 20	3.7
	September 25, 2012 ^c (carboy)	500	529 ± 4	5.9
		1,000	NA ^d	NA ^d
		2,000	2,090 ± 30	4.7
October 11, 2012	October 14, 2012	500	490 ± 5 / 499 ± 4	-1.9 / -0.3
		1,000	998 ± 3 / 1,050 ± 20	-0.2 / 5.3
		2,000	2,110 ± 30	5.7
December 4, 2012	December 6, 2012	500	515 ± 8 / 538 ± 6	3.1 / 7.5
		1,000	1,050 ± 20 / 1,060 ± 10	5.0 / 5.7
		2,000	2,060 ± 30	3.2

Date Prepared	Date Analyzed	Target Concentration (mg/L)	Determined Concentration (mg/L) ^a	Difference from Target (%)
February 26, 2013	February 27, 2012	500	478 ± 3 / 476 ± 7	-4.5 / 4.8
		1,000	970 ± 22 / 950 ± 17	-5.0
		2,000	1,890 ± 10	-5.3
	April 9, 2012 ^b (bottle)	500	453 ± 16	-9.5
		1,000	915 ± 13	-8.5
		2,000	1,900 ± 20	-5.0
	April 9, 2012 ^c (carboy)	500	468 ± 3	-6.5
		1,000	937 ± 10	-6.3
		2,000	1,880 ± 20	-5.8
April 25, 2013	April 26, 2013	500	527 ± 23 / 494 ± 17	5.4 / -1.3
		1,000	995 ± 9 / 988 ± 6	-0.5 / -1.2
		2,000	1,900 ± 10	-5.2
July 16, 2013	July 18, 2013	500	479 ± 17 / 509 ± 21	-4.1 / 1.7
		1,000	955 ± 71 / 970 ± 10	-4.5 / -3.0
		2,000	2,000 ± 70	0.0
September 10, 2013	September 14, 2013	500	454 ± 8 / 501 ± 23	-9.1 / 0.3
		1,000	1,090 ± 20 / 1,080 ± 0	8.7 / 8.0
		2,000	2,150 ± 100	7.5
	October 23, 2012 ^b (bottle)	500	503 ± 7	0.5
		1,000	1,040 ± 10	3.7
		2,000	2,090 ± 10	4.5
	October 23, 2012 ^c (carboy)	500	485 ± 16	-2.9
		1,000	1,060 ± 30	6.0
		2,000	1,980 ± 30	-0.8
December 4, 2013	December 6, 2013	500	495 ± 8 / 480 ± 21	-1.1 / -4.1
		1,000	973 ± 13 / 900 ± 6.9	-2.7 / -10.0
		2,000	1,910 ± 30	-4.5

1 Values on either side of the / represent multiple formulations of the same dose prepared on that date.


2 NA = not analyzed.

3 ^aAverage of triplicate analysis.

4 ^bAnimal room sample from the formulation remaining in the drinking water bottle.

5 ^cAnimal room sample from the formulation collected from the carboy.

6 ^dSample not collected from carboy.



1

2 **Figure A-1. Infrared Absorption Spectrum of Sodium Tungstate Dihydrate**

3

1 **Appendix B. Ingredients, Nutrient Composition, and**
2 **Contaminant Levels in NIH-07 Rat Ration and NTP-2000 Rat**
3 **and Mouse Ration**

4 **Tables**

5 Table B-1. Ingredients of NIH-07 Rat RationB-2
6 Table B-2. Vitamins and Minerals in NIH-07 Rat RationB-2
7 Table B-3. Nutrient Composition of NIH-07 Rat RationB-3
8 Table B-4. Contaminant Levels in NIH-07 Rat RationB-5
9 Table B-5. Ingredients of NTP-2000 Rat and Mouse RationB-6
10 Table B-6. Vitamins and Minerals in NTP-2000 Rat and Mouse RationB-7
11 Table B-7. Nutrient Composition of NTP-2000 Rat and Mouse RationB-8
12 Table B-8. Contaminant Levels in NTP-2000 Rat and Mouse RationB-10
13

1 **B.1. NIH-07 Feed**2 **Table B-1. Ingredients of NIH-07 Rat Ration**

Ingredients	Percent by Weight
Ground Hard Winter Wheat	23.00
Ground #2 Yellow Shelled Corn	24.25
Wheat Middlings	10.0
Alfalfa Meal (Dehydrated, 17% Protein)	4.0
Soybean Meal (47% Protein)	12.0
Fish Meal (62% Protein)	10.0
Soy Oil (Without Preservatives)	2.5
Dried Brewer's Yeast	2.0
Calcium Carbonate (USP)	0.5
Vitamin Premix ^a	0.25
Mineral Premix ^b	0.15
Calcium Phosphate, Dibasic (USP)	1.25
Sodium Chloride	0.5
Choline Chloride (70% Choline)	0.10
Dried Skim Milk	5.00
Dried Molasses	1.50
Corn Gluten Meal (60% Protein)	3.00
Methionine	0.0

3 USP = United States Pharmacopeia.

4 ^aWheat middlings as carrier.5 ^bCalcium carbonate as carrier.6 **Table B-2. Vitamins and Minerals in NIH-07 Rat Ration**

	Amount ^a	Source
Vitamins		
Vitamin A	6,062 IU	Stabilized vitamin A palmitate or acetate
Vitamin D	5,070 IU	D-activated animal sterol
Vitamin K	3.1 mg	Menadione sodium bisulfite complex
Vitamin E	22 IU	α -Tocopheryl acetate
Niacin	33 mg	–
Folic Acid	2.4 mg	–
d-Pantothenic Acid	19.8 mg	d-Calcium pantothenate
Riboflavin	3.8 mg	–
Thiamine	11 mg	Thiamine mononitrate

	Amount ^a	Source
B ₁₂	50 µg	–
Pyridoxine	6.5 mg	Pyridoxine hydrochloride
Biotin	0.15 mg	d-Biotin
Minerals		
Iron	132 mg	Iron sulfate
Zinc	18 mg	Zinc oxide
Manganese	66 mg	Manganese oxide
Copper	4.4 mg	Copper sulfate
Iodine	2.0 mg	Calcium iodate
Cobalt	0.44 mg	Cobalt carbonate

1 ^aPer kg of finished product.

2 **Table B-3. Nutrient Composition of NIH-07 Rat Ration**

Nutrient	Mean ± Standard Deviation	Range	Number of Samples
Protein (% by Weight)	24.65 ± 1.344	23.7–25.6	2
Crude Fat (% by Weight)	5.3 ± 0.141	5.2–5.4	2
Crude Fiber (% by Weight)	3.57 ± 0.184	3.44–3.70	2
Ash (% by Weight)	6.565 ± 0.049	6.53–6.60	2
Amino Acids (% of Total Diet)			
Arginine	1.380 ± 0.06	1.3–1.49	10
Cystine	0.322 ± 0.031	0.274–0.372	10
Glycine	1.150 ± 0.070	1.06–1.31	10
Histidine	0.518 ± 0.024	0.497–0.553	10
Isoleucine	0.984 ± 0.024	0.952–1.03	10
Leucine	2.018 ± 0.067	1.93–2.13	10
Lysine	1.243 ± 0.051	1.13–1.32	10
Methionine	0.488 ± 0.016	0.468–0.515	10
Phenylalanine	1.097 ± 0.022	1.07–1.12	10
Threonine	0.918 ± 0.031	0.883–0.961	10
Tryptophan	0.277 ± 0.020	0.265–0.326	10
Tyrosine	0.860 ± 0.037	0.785–0.894	10
Valine	1.134 ± 0.025	1.11–1.17	10
Essential Fatty Acids (% of Total Diet)			
Linoleic	2.30 ± 0.219	1.99–2.59	10
Linolenic	0.25 ± 0.275	0.217–0.296	10

Nutrient	Mean \pm Standard Deviation	Range	Number of Samples
Vitamins			
Vitamin A (IU/kg)	4,545 \pm 67.2	4,070–5,020	2
α -Tocopherol (ppm)	6,704 \pm 21,045	40.3–66,600	10
Thiamine (ppm) ^a	14.95 \pm 0.778	14.4–15.5	2
Riboflavin (ppm)	14.47 \pm 3.352	10.0–19.8	10
Niacin (ppm)	99.33 \pm 8.235	87.0–112.0	10
Pantothenic Acid (ppm)	44.38 \pm 3.806	38.2–51.1	10
Pyridoxine (ppm) ^a	12.876 \pm 3.171	9.63–19.7	10
Folic Acid (ppm)	2.482 \pm 0.487	1.68–3.09	10
Biotin (ppm)	0.3283 \pm 0.172	0.0–0.638	10
B ₁₂ (ppb)	49.4 \pm 6.83	41.8–61.6	10
Choline (as chloride) (ppm)	1,821.0 \pm 197.5	1,570–2,200	10
Minerals			
Calcium (%)	1.205 \pm 0.078	1.15–1.26	2
Phosphorus (%)	0.938 \pm 0.036	0.912–0.963	2
Potassium (%)	0.830 \pm 0.036	0.769–0.88	10
Chloride (%)	0.652 \pm 0.106	0.441–0.8	10
Sodium (%)	0.378 \pm 0.46	0.318–0.469	10
Magnesium (%)	0.187 \pm 0.014	0.17–0.218	10
Iron (ppm)	385.1 \pm 54.9	276.0–469.0	10
Manganese (ppm)	90.81 \pm 7.566	80.7–104.0	10
Zinc (ppm)	64.15 \pm 10.07	52.4–89.2	10
Copper (ppm)	14.13 \pm 2.57	11.9–21.1	10
Iodine (ppm)	1.811 \pm 0.992	0.54–3.45	10
Chromium (ppm)	3.946 \pm 0.036	3.89–4.0	8
Cobalt (ppm)	0.5155 \pm 0.267	0.01–0.963	10

1 ^aAs hydrochloride.

1 **Table B-4. Contaminant Levels in NIH-07 Rat Ration**

	Mean ± Standard Deviation	Range	Number of Samples
Contaminants			
Arsenic (ppm)	0.2965 ± 0.029	0.26–0.317	2
Cadmium (ppm)	0.082 ± 0.004	0.079–0.085	2
Lead (ppm)	0.0785 ± 0.001	0.078–0.079	2
Mercury (ppm) ^a	0.0135 ± 0.002	0.012–0.015	2
Selenium (ppm)	0.4065 ± 0.110	0.329–0.484	2
Aflatoxins (ppb) ^a	5	–	2
Nitrate Nitrogen (ppm) ^b	19.55 ± 1.344	18.6–20.5	2
Nitrite Nitrogen (ppm) ^{a,b}	<0.61	–	2
BHA (ppm) ^{a,c}	<1.0	–	2
BHT (ppm) ^{a,c}	<1.0	–	2
Aerobic Plate Count (CFU/gm) ^a	<10	–	2
Coliform (MPN/gm) ^a	<3	–	2
<i>E. coli</i> (MPN/gm) ^a	<10	–	2
<i>Salmonella</i> (MPN/gm)	0	–	2
Total Nitrosamines (ppb) ^d	6.6 ± 1.70	5.4–7.8	2
N-Ndimethylamine (ppb) ^d	0	–	2
N-Npyrrolidine (ppb) ^d	6.6 ± 1.70	5.4–7.8	2
Pesticides (ppm)			
α-BHC ^a	<0.01	–	2
β-BHC ^a	<0.02	–	2
γ-BHC ^a	<0.01	–	2
δ-BHC ^a	<0.01	–	2
Heptachlor ^a	<0.01	–	2
Aldrin ^a	<0.01	–	2
Heptachlor Epoxide ^a	<0.01	–	2
DDE ^a	<0.01	–	2
DDD ^a	<0.01	–	2
DDT ^a	<0.01	–	2
HCB ^a	<0.01	–	2
Mirex ^a	<0.01	–	2
Methoxychlor ^a	<0.05	–	2
Dieldrin ^a	<0.01	–	2
Endrin ^a	<0.01	–	2

	Mean ± Standard Deviation	Range	Number of Samples
Telodrin ^a	<0.01	–	2
Chlordane ^a	<0.05	–	2
Toxaphene ^a	<0.10	–	2
Estimated PCBs ^a	<0.20	–	2
Ronnel ^a	<0.01	–	2
Ethion ^a	<0.02	–	2
Trithion ^a	<0.05	–	2
Diazinon ^a	<0.10	–	2
Methyl Chlorpyrifos	0.0391 ± 0.027	0.0200–0.0582	2
Methyl Parathion ^a	<0.02	–	2
Ethyl Parathion ^a	<0.02	–	2
Malathion ^a	<0.02	–	2
Endosulfan I ^a	<0.01	–	2
Endosulfan II ^a	<0.01	–	2
Endosulfane Sulfate ^a	<0.03	–	2

1 All samples were irradiated. BHA = butylated hydroxyanisole; BHT = butylated hydroxytoluene; CFU = colony-forming units;
2 MPN = most probable number; BHC = hexachlorocyclohexane or benzene hexachloride; DDE =
3 dichlorodiphenyldichloroethylene; DDD = dichlorodiphenyldichloroethane; DDT = dichlorodiphenyltrichloroethane; HCB =
4 hexachlorobenzene; PCB = polychlorinated biphenyl.

5 ^aAll values were below the detection limit. The detection limit is given as the mean.

6 ^bSources of contamination include alfalfa, grains, and fish meal.

7 ^cSources of contamination include soy oil and fish meal.

8 ^dAll values were corrected for percent recovery.

9 B.2. NTP-2000 Feed

10 **Table B-5. Ingredients of NTP-2000 Rat and Mouse Ration**

Ingredients	Percent by Weight
Ground Hard Winter Wheat	22.26
Ground #2 Yellow Shelled Corn	22.18
Wheat Middlings	15.0
Oat Hulls	8.5
Alfalfa Meal (Dehydrated, 17% Protein)	7.5
Purified Cellulose	5.5
Soybean Meal (49% Protein)	5.0
Fish Meal (60% Protein)	4.0
Corn Oil (without Preservatives)	3.0
Soy Oil (without Preservatives)	3.0
Dried Brewer's Yeast	1.0

Ingredients	Percent by Weight
Calcium Carbonate (USP)	0.9
Vitamin Premix ^a	0.5
Mineral Premix ^b	0.5
Calcium Phosphate, Dibasic (USP)	0.4
Sodium Chloride	0.3
Choline Chloride (70% Choline)	0.26
Methionine	0.2

1 USP = United States Pharmacopeia.

2 ^aWheat middlings as carrier.

3 ^bCalcium carbonate as carrier.

4 **Table B-6. Vitamins and Minerals in NTP-2000 Rat and Mouse Ration**

	Amount ^a	Source
Vitamins		
Vitamin A	4,000 IU	Stabilized vitamin A palmitate or acetate
Vitamin D	1,000 IU	D-activated animal sterol
Vitamin K	1.0 mg	Menadione sodium bisulfite complex
α -Tocopheryl Acetate	100 IU	–
Niacin	23 mg	–
Folic Acid	1.1 mg	–
d-Pantothenic Acid	10 mg	d-Calcium pantothenate
Riboflavin	3.3 mg	–
Thiamine	4 mg	Thiamine mononitrate
B ₁₂	52 μ g	–
Pyridoxine	6.3 mg	Pyridoxine hydrochloride
Biotin	0.2 mg	d-Biotin
Minerals		
Magnesium	514 mg	Magnesium oxide
Iron	35 mg	Iron sulfate
Zinc	12 mg	Zinc oxide
Manganese	10 mg	Manganese oxide
Copper	2.0 mg	Copper sulfate
Iodine	0.2 mg	Calcium iodate
Chromium	0.2 mg	Chromium acetate

5 ^aPer kg of finished product.

1 **Table B-7. Nutrient Composition of NTP-2000 Rat and Mouse Ration**

Nutrient	Mean \pm Standard Deviation	Range	Number of Samples
Protein (% by Weight)	14.61 \pm 0.537	13.9–16.5	32
Crude Fat (% by Weight)	8.39 \pm 0.376	7.7–9.2	32
Crude Fiber (% by Weight)	9.26 \pm 0.603	7.1–10.1	32
Ash (% by Weight)	4.93 \pm 0.138	4.66–5.2	32
Amino Acids (% of Total Diet)			
Arginine	0.805 \pm 0.075	0.67–0.97	29
Cystine	0.220 \pm 0.021	0.15–0.25	29
Glycine	0.702 \pm 0.038	0.62–0.8	29
Histidine	0.342 \pm 0.070	0.27–0.68	29
Isoleucine	0.549 \pm 0.040	0.43–0.66	29
Leucine	1.100 \pm 0.063	0.96–1.24	29
Lysine	0.700 \pm 0.104	0.31–0.86	29
Methionine	0.409 \pm 0.042	0.26–0.49	29
Phenylalanine	0.623 \pm 0.047	0.471–0.72	29
Threonine	0.513 \pm 0.041	0.43–0.61	29
Tryptophan	0.155 \pm 0.027	0.11–0.2	29
Tyrosine	0.422 \pm 0.066	0.28–0.54	29
Valine	0.666 \pm 0.040	0.55–0.73	29
Essential Fatty Acids (% of Total Diet)			
Linoleic	3.94 \pm 0.235	3.49–4.55	29
Linolenic	0.30 \pm 0.064	0.005–0.368	29
Vitamins			
Vitamin A (IU/kg)	3,757 \pm 70.50	2,520–5,450	32
Vitamin D (IU/kg) ^a	1,000	–	–
α -Tocopherol (ppm)	2,456 \pm 128.17	13.6–69,100	29
Thiamine (ppm) ^b	7.5 \pm 0.614	6.1–9.0	32
Riboflavin (ppm)	8.17 \pm 2.841	42–17.5	29
Niacin (ppm)	78.66 \pm 8.11	66.4–98.2	29
Pantothenic Acid (ppm)	26.42 \pm 11.05	17.4–81.0	29
Pyridoxine (ppm) ^b	9.75 \pm 2.045	6.44–14.3	29
Folic Acid (ppm)	1.58 \pm 0.43	1.15–3.27	29
Biotin (ppm)	0.323 \pm 0.093	0.2–0.704	29
B ₁₂ (ppb)	50.41 \pm 34.89	18.3–174	29
Choline (as Chloride) (ppm)	2,593 \pm 633.8	1,160–3,790	29

Nutrient	Mean \pm Standard Deviation	Range	Number of Samples
Minerals			
Calcium (%)	0.902 \pm 0.056	0.697–1.02	32
Phosphorus (%)	0.551 \pm 0.023	0.504–0.615	32
Potassium (%)	0.668 \pm 0.029	0.626–0.733	29
Chloride (%)	0.392 \pm 0.044	0.3–0.517	29
Sodium (%)	0.195 \pm 0.027	0.16–0.283	29
Magnesium (%)	0.217 \pm 0.054	0.185–0.49	29
Sulfur (%)	0.170 \pm 0.029	0.116–0.209	14
Iron (ppm)	191.6 \pm 36.18	135–311	29
Manganese (ppm)	50.11 \pm 9.42	21–73.1	29
Zinc (ppm)	57.3 \pm 25.54	43.3–184	29
Copper (ppm)	7.57 \pm 2.49	3.21–16.3	29
Iodine (ppm)	0.513 \pm 0.221	0–0.972	29
Chromium (ppm)	1.02 \pm 1.04	0.33–3.97	28
Cobalt (ppm)	0.222 \pm 0.152	0.0857–0.864	27

1 ^aFrom formulation.
2 ^bAs hydrochloride.

1 **Table B-8. Contaminant Levels in NTP-2000 Rat and Mouse Ration**

	Mean \pm Standard Deviation	Range	Number of Samples
Contaminants			
Arsenic (ppm)	0.199 \pm 0.045	0.143–0.307	32
Cadmium (ppm)	0.065 \pm 0.080	0.015–0.5	32
Lead (ppm)	0.16 \pm 0.202	0.059–1.19	32
Mercury (ppm) ^a	0.012 \pm 0.004	0.01–0.026	32
Selenium (ppm)	0.169 \pm 0.039	0.029–0.266	32
Aflatoxins (ppb) ^a	<5.0	–	32
Nitrate Nitrogen (ppm) ^b	18.88 \pm 9.5	10.0–45.9	32
Nitrite Nitrogen (ppm) ^{a,b}	0.61	–	32
BHA (ppm) ^{a,c}	<1.00	–	32
BHT (ppm) ^{a,c}	1.03 \pm 0.156	1.0–1.88	32
Aerobic Plate Count (CFU/gm)	14.84 \pm 18.56	10.0–110.0	32
Coliform (MPN/gm)	3.0	–	32
<i>E. coli</i> (MPN/gm) ^a	<10.0	–	32
<i>Salmonella</i> (MPN/gm)	Negative	–	32
Total Nitrosamines (ppb) ^d	9.8 \pm 5.12	0–19.9	32
N-Ndimethylamine (ppb) ^d	2.3 \pm 2.6	0–11.1	32
N-Npyrrolidine (ppb) ^d	7.8 \pm 4.24	0–18.6	32
Pesticides (ppm)			
α -BHC ^a	<0.01	–	32
β -BHC ^a	<0.02	–	32
γ -BHC ^a	<0.01	–	32
δ -BHC ^a	<0.01	–	32
Heptachlor ^a	<0.01	–	32
Aldrin ^a	<0.01	–	32
Heptachlor Epoxide ^a	<0.01	–	32
DDE ^a	<0.01	–	32
DDD ^a	<0.01	–	32
DDT ^a	<0.01	–	32
HCB ^a	<0.01	–	32
Mirex ^a	<0.01	–	32
Methoxychlor ^a	<0.05	–	32
Dieldrin ^a	<0.01	–	32
Endrin ^a	<0.01	–	32

	Mean \pm Standard Deviation	Range	Number of Samples
Telodrin ^a	<0.01	–	32
Chlordane ^a	<0.05	–	32
Toxaphene ^a	<0.10	–	32
Estimated PCBs ^a	<0.20	–	32
Ronnel ^a	<0.01	–	32
Ethion ^a	<0.02	–	32
Trithion ^a	<0.05	–	32
Diazinon ^a	<0.10	–	32
Methyl Chlorpyrifos	0.112 \pm 0.141	0.02–0.686	32
Methyl Parathion ^a	<0.02	–	32
Ethyl Parathion ^a	<0.02	–	32
Malathion	0.07 \pm 0.07	0.02–0.234	32
Endosulfan I ^a	<0.01	–	32
Endosulfan II ^a	<0.01	–	32
Endosulfane Sulfate ^a	<0.03	–	32

1 All samples were irradiated. BHA = butylated hydroxyanisole; BHT = butylated hydroxytoluene; CFU = colony-forming units;
2 MPN = most probable number; BHC = hexachlorocyclohexane or benzene hexachloride; DDE =
3 dichlorodiphenyldichloroethylene; DDD = dichlorodiphenyldichloroethane; DDT = dichlorodiphenyltrichloroethane; HCB =
4 hexachlorobenzene; PCB = polychlorinated biphenyl.

5 ^aAll values were below the detection limit. The detection limit is given as the mean.

6 ^bSources of contamination include alfalfa, grains, and fish meal.

7 ^cSources of contamination include soy oil and fish meal.

8 ^dAll values were corrected for percent recovery.

1 **Appendix C. Sentinel Animal Program**

2 **Table of Contents**

3 C.1. Methods.....C-2
4 C.2. Results.....C-2

5
6 **Tables**

7 Table C-1. Methods and Results for Sentinel Animal Testing in Male and Female RatsC-3
8 Table C-2. Methods and Results for Sentinel Animal Testing in Male and Female Mice.....C-4
9

1 **C.1. Methods**

2 Rodents used in the National Toxicology Program are produced in optimally clean facilities to
3 eliminate potential pathogens that might affect study results. The Sentinel Animal Program is
4 part of the periodic monitoring of animal health that occurs during the toxicological evaluation of
5 test compounds. Under this program, the disease state of the rodents is monitored via sera or
6 feces from extra (sentinel) or exposed animals in the study rooms. The sentinel animals and the
7 study animals are subject to identical environmental conditions. Furthermore, the sentinel
8 animals come from the same production source and weanling groups as the animals used for the
9 studies of test compounds.

10 In these toxicology and carcinogenesis studies, blood samples were collected from each sentinel
11 animal, allowed to clot, and the serum was separated. Additionally, fecal samples were collected
12 and tested for endoparasites and *Helicobacter* species. All samples were processed appropriately,
13 with serology and *Helicobacter* testing sent to IDEXX BioResearch (formerly Rodent Animal
14 Diagnostic Laboratory [RADIL], University of Missouri), Columbia, MO, for determination of
15 the presence of pathogens. Evaluation for endo- and ectoparasites was performed in-house by the
16 testing laboratory.

17 The laboratory methods and agents for which testing was performed are tabulated in Table C-1
18 and Table C-2 below; the times at which samples were collected during the studies are also
19 listed.

20 **C.2. Results**

21 Rats: Positive for endoparasites – pinworms (*Syphacia* spp.) for the 2-year study. All other test
22 results were negative.

23 Mice: All test results were negative.

24

1 **Table C-1. Methods and Results for Sentinel Animal Testing in Male and Female Rats**

Collection Time Points	Three-month Study			Two-year Study							
	Quarantine ^a	3.5 Weeks ^b	End of Study	Quarantine ^a	1 Month	6 Months	12 Months	16 Months ^c	17 Months ^c	18 Months	End of Study
Number Examined (Males/Females)	0/10	0/8	5/5	0/10	5/5	5/5	5/5	0/1	0/1	7/7	5/5
Method/Test											
Multiplex Fluorescent Immunoassay (MFI)											
Kilham rat virus (KRV)	-	-	-	-	-	-	-	-	-	-	-
<i>Mycoplasma pulmonis</i>	-	-	-	-	-	-	-	-	-	-	-
Parvo NS-1	-	-	-	NT	NT	NT	NT	NT	NT	NT	NT
Pneumonia virus of mice (PVM)	-	-	-	-	-	-	-	-	-	-	-
Rat coronavirus/sialodacryoadenitis virus (RCV/SDA)	-	-	-	-	-	-	-	-	-	-	-
Rat minute virus (RMV)	-	-	-	-	-	-	-	-	-	-	-
Rat parvo virus (RPV)	-	-	-	-	-	-	-	-	-	-	-
Rat theilovirus (RTV)	-	-	-	-	-	-	-	-	-	-	-
Sendai	-	-	-	-	-	-	-	-	-	-	-
Theiler's murine encephalomyelitis virus (TMEV)	-	-	-	NT	NT	NT	NT	NT	NT	NT	NT
Toolan's H1	-	-	-	-	-	-	-	-	-	-	-
Immunofluorescence Assay (IFA)											
<i>Pneumocystis carinii</i>	NT	NT	NT	NT	NT	-	NT	NT	NT	NT	NT
In-house Evaluation											
Endoparasite evaluation (evaluation of cecal content)	NT	NT	NT	+	+	NT	+	+	+	+	NT
Ectoparasite evaluation (evaluation of perianal surface)	NT	NT	NT	-	-	NT	-	-	-	-	NT

2 - = negative; + = positive; NT = not tested.

3 ^aAge-matched nonpregnant females.4 ^bTime-mated females that did not have a litter.5 ^cSingle sentinel rat tested at this time point.

1 **Table C-2. Methods and Results for Sentinel Animal Testing in Male and Female Mice**

Collection Time Points	Three-month Study			Two-year Study					
	1 Month	End of Study	Quarantine	1 Month	6 Months	12 Months	15 Months	18 Months	End of Study
Number Examined (Males/Females)	5/5	5/5	5/5	5/5	5/5	5/5	5/5	5/5	5/5
Method/Test									
Multiplex Fluorescent Immunoassay (MFI)									
Ectromelia virus	-	-	-	-	-	-	NT	-	-
Epizootic Diarrhea of Infant Mice	-	-	-	-	-	-	NT	-	-
Lymphocytic choriomeningitis virus (LCMV)	-	-	-	-	-	-	NT	-	-
<i>Mycoplasma pulmonis</i>	-	-	-	-	-	-	NT	-	-
Mouse hepatitis virus (MHV)	-	-	-	-	-	-	NT	-	-
Mouse norovirus (MNV)	-	-	-	-	-	-	NT	-	-
Parvo NS-1	-	-	-	-	-	-	NT	-	-
Mouse parvovirus (MPV)	-	-	-	-	-	-	NT	-	-
Minute virus of mice (MVM)	-	-	-	-	-	-	NT	-	-
Pneumonia virus of mice (PVM)	-	-	-	-	-	-	NT	-	-
Reovirus (REO3)	-	-	-	-	-	-	NT	-	-
Sendai	-	-	-	-	-	-	NT	-	-
Theiler's murine encephalomyelitis virus (TMEV) GDVII	-	-	-	-	-	-	NT	-	-
Immunofluorescence Assay (IFA)									
Epizootic diarrhea of infant mice	NT	NT	NT	NT	NT	NT	NT	-	NT
Mouse norovirus (MNV)	NT	NT	NT	NT	NT	NT	NT	-	NT
Polymerase Chain Reaction (PCR)									
<i>Helicobacter</i> species	NT	NT	NT	NT	NT	NT	NT	-	NT
In-house Evaluation									
Endoparasite evaluation (evaluation of cecal content)	NT	NT	-	-	-	-	-	-	NT
Ectoparasite evaluation (evaluation of perianal surface)	NT	NT	-	-	-	-	-	-	NT

2 - = negative; + = positive; NT = not tested.

1 **Appendix D. Genetic Toxicology**

2 **Table of Contents**

3 D.1. Bacterial Mutagenicity D-2
4 D.2. Micronucleus Assay D-4
5 D.3. Comet Assay D-7
6

7 **Tables**

8 Table D-1. Mutagenicity of Sodium Tungstate Dihydrate in Bacterial Tester Strains..... D-3
9 Table D-2. Frequency of Micronuclei in Peripheral Blood Erythrocytes of Male and
10 Female Rats in the Three-month Drinking Water Study of Sodium Tungstate
11 Dihydrate D-6
12 Table D-3. Frequency of Micronuclei in Peripheral Blood Erythrocytes of Male and
13 Female Mice in the Three-month Drinking Water Study of Sodium Tungstate
14 Dihydrate D-7
15 Table D-4. DNA Damage in Male and Female Rats Exposed to Sodium Tungstate
16 Dihydrate in Drinking Water for Three Months D-10
17 Table D-5. DNA Damage in Male and Female Mice Exposed to Sodium Tungstate
18 Dihydrate in Drinking Water for Three Months D-11
19

1 **D.1. Bacterial Mutagenicity**

2 **D.1.1. Bacterial Mutagenicity Test Protocol**

3 Testing procedures were modified from those originally reported by Zeiger et al. (1992). Coded
4 samples of sodium tungstate dihydrate (ST; the same chemical lot that was used in the 2-year
5 bioassays) were incubated with the *Salmonella typhimurium* (TA98, TA100) or *Escherichia coli*
6 (WP2 *uvrA*/pKM101) tester strains either in buffer or S9 mix (metabolic activation enzymes and
7 cofactors from phenobarbital/benzoflavone-induced male Sprague Dawley rat liver) for
8 20 minutes at 37°C. Top agar supplemented with *L*-histidine (or tryptophan for the *E. coli* strain)
9 and *d*-biotin was added, and the contents of the tubes were mixed and poured onto the surfaces of
10 minimal glucose agar plates. Histidine- or tryptophan-independent mutant colonies arising on
11 these plates were counted following incubation for two days at 37°C.

12 Each trial consisted of triplicate plates of concurrent positive and negative controls and of at least
13 five doses of ST. The highest concentration tested was limited by toxicity in strain TA100; the
14 other two strains were tested up to the assay limit dose of 6,000 µg/plate. All trials were
15 repeated.

16 In this assay, a positive response is defined as a reproducible, dose-related increase in
17 histidine-independent (revertant) colonies in any one strain/activation combination. An equivocal
18 response is defined as an increase in revertants that is not dose-related, not reproducible, or not
19 of sufficient magnitude to support a determination of mutagenicity. A negative response is
20 obtained when no increase in revertant colonies is observed following chemical treatment. No
21 minimum percentage or fold increase is required for a chemical to be judged positive or weakly
22 positive, although positive calls are typically reserved for increases in mutant colonies that are at
23 least twofold over background.

24 **D.1.2. Results**

25 ST (12.5 to 6,000 µg/plate) was not mutagenic in *Salmonella typhimurium* strains TA98 or
26 TA100, or in *Escherichia coli* WP2 *uvrA*/pKM101, when tested with or without exogenous
27 metabolic activation provided by phenobarbital/benzoflavone-induced rat S9 and cofactors
28 (Table D-1).

1 **Table D-1. Mutagenicity of Sodium Tungstate Dihydrate in Bacterial Tester Strains^a**

Strain	Concentration (µg/plate)	Without S9	Without S9	With 10% Rat S9	With 10% Rat S9
TA98					
	0	28 ± 2	21 ± 2	46 ± 2	28 ± 1
	12.5	25 ± 1	25 ± 5	33 ± 3	28 ± 3
	50	27 ± 2	30 ± 4	29 ± 2	30 ± 4
	125	23 ± 2	19 ± 3	32 ± 1	31 ± 4
	500	22 ± 6	23 ± 5	34 ± 5	24 ± 1
	1,500	19 ± 2	17 ± 3	34 ± 6	38 ± 4
	6,000	30 ± 2	23 ± 6	33 ± 4	29 ± 6
Trial Summary		Negative	Negative	Negative	Negative
Positive Control ^b		518 ± 31	465 ± 25	1,216 ± 77	1,017 ± 26
TA100					
	0	99 ± 3	123 ± 8	113 ± 2	117 ± 6
	12.5	111 ± 8	129 ± 8	112 ± 6	113 ± 5
	50	106 ± 7	100 ± 5	106 ± 5	108 ± 10
	125	118 ± 8	107 ± 9	116 ± 3	111 ± 1
	500	113 ± 3	97 ± 3	120 ± 2	105 ± 4
	1,500	100 ± 5	113 ± 7	117 ± 6	138 ± 10
	6,000	100 ± 6	110 ± 6	111 ± 11	117 ± 12
Trial Summary		Negative	Negative	Negative	Negative
Positive Control		850 ± 9	800 ± 43	2,640 ± 135	1,724 ± 18
<i>Escherichia coli</i> WP2 <i>uvrA</i>/pKM101					
	0	186 ± 8	191 ± 12	194 ± 11	219 ± 7
	12.5	159 ± 3	183 ± 7	219 ± 6	183 ± 11
	50	183 ± 7	161 ± 6	221 ± 21	179 ± 8
	125	166 ± 18	156 ± 6	194 ± 4	181 ± 12
	500	162 ± 6	171 ± 14	200 ± 14	206 ± 15
	1,500	171 ± 8	185 ± 10	213 ± 17	211 ± 16
	6,000	185 ± 12	165 ± 18	193 ± 14	197 ± 16
Trial Summary		Negative	Negative	Negative	Negative
Positive Control		1,105 ± 49	959 ± 21	1,234 ± 70	1,009 ± 36

2 ^aStudies performed at Integrated Laboratory Systems, LLC. Data are presented as revertants/plate (mean ± standard error) from
3 three plates; 0 µg/plate served as the solvent control.

4 ^bThe positive controls in the absence of metabolic activation were sodium azide (TA100), 2-nitrofluorene (TA98), and methyl
5 methanesulfonate (*E. coli*). The positive control for metabolic activation with all strains was 2-aminoanthracene.

1 **D.2. Micronucleus Assay**

2 **D.2.1. Peripheral Blood Micronucleus Test Protocol**

3 At termination of the 3-month toxicity studies of ST, blood samples (approximately 200 μ L)
4 were collected from male and female rats and mice, placed in ethylenediaminetetraacetic acid
5 (EDTA)-coated tubes, and shipped overnight to the testing laboratory. Upon arrival, blood
6 samples were fixed in ultracold methanol using a MicroFlowPLUS Kit (Litron Laboratories,
7 Rochester, NY) according to the manufacturer's instructions. Fixed samples were stored in a
8 -80°C freezer until analysis. Thawed blood samples were analyzed for frequency of
9 micronucleated immature erythrocytes (polychromatic erythrocytes [PCEs], reticulocytes) and
10 mature erythrocytes (normochromatic erythrocytes, NCEs) using a flow cytometer;⁹⁹ both the
11 mature and the immature erythrocyte populations can be analyzed separately by employing
12 special cell surface markers to differentiate the two cell types. Because the very young
13 reticulocyte subpopulation (CD71-positive cells) can be targeted using this technique, rat blood
14 samples can be analyzed for damage that occurred in the bone marrow within the past 24–
15 48 hours, before the rat spleen appreciably alters the percentage of micronucleated reticulocytes
16 in circulation.¹⁰⁰ In mice, both the immature and mature erythrocyte populations can be
17 evaluated for micronucleus frequency because the mouse spleen does not sequester and eliminate
18 damaged erythrocytes. Damaged erythrocytes achieve steady state in the peripheral blood of
19 mice following four weeks of continuous exposure. Approximately 20,000 reticulocytes and
20 1×10^6 erythrocytes were analyzed per animal for frequency of micronucleated cells, and the
21 percentage of immature erythrocytes (% PCE) was calculated as a measure of bone marrow
22 toxicity resulting from ST exposure.

23 Prior experience with the large number of cells scored using flow cytometric scoring
24 techniques¹⁰¹ suggests it is reasonable to assume that the proportion of micronucleated
25 reticulocytes is approximately normally distributed. The statistical tests selected for trend and for
26 pairwise comparisons with the control group depend on whether the variances among the groups
27 are equal. The Levene test at $\alpha = 0.05$ is used to test for equal variances. In the case of equal
28 variances, linear regression is used to test for a linear trend with exposure concentration and the
29 Williams test is used to test for pairwise differences between each exposed group and the control
30 group. In the case of unequal variances, the Jonckheere test is used to test for linear trend and the
31 Dunn test is used for pairwise comparisons of each exposed group with the control group. To
32 correct for multiple pairwise comparisons, the p value for each comparison with the control
33 group is multiplied by the number of comparisons made. In the event that this product is >1.00 , it
34 is replaced with 1.00. Trend tests and pairwise comparisons with the controls are considered
35 significant at $p \leq 0.025$.

36 In the micronucleus test, it is preferable to base a positive result on the presence of both a
37 significant trend as well as at least one significantly elevated exposure group compared with the
38 corresponding control group. In addition, historical control data are used to evaluate the
39 biological significance of any observed response. Both statistical significance and biological
40 significance are considered when arriving at a call. The presence of either a significant trend or a
41 single significant exposure group generally results in an equivocal call. The absence of both a
42 trend and a significant exposure group results in a negative call. Ultimately, the scientific staff
43 determines the final call after considering the results of statistical analyses, reproducibility of any
44 effects observed (in acute studies), and the magnitudes of those effects.

1 **D.2.2. Evaluation Protocol**

2 These are the basic guidelines for arriving at an overall assay result for assays performed by the
3 National Toxicology Program. Statistical as well as biological factors are considered. For an
4 individual assay, the statistical procedures for data analysis have been described in the preceding
5 protocols. There have been instances, however, in which multiple samples of a chemical were
6 tested in the same assay, and different results were obtained among these samples and/or among
7 laboratories. Results from more than one aliquot or from more than one laboratory are not simply
8 combined into an overall result. Rather, all the data are critically evaluated, particularly those
9 concerning pertinent protocol variations, in determining the weight of evidence for an overall
10 conclusion of chemical activity in an assay. In addition to multiple aliquots, the in vitro assays
11 are conducted with and without exogenous metabolic activation. Results obtained in the absence
12 of activation are not combined with results obtained in the presence of activation; each testing
13 condition is evaluated separately. The summary table in the Abstract of this Technical Report
14 presents a result that is a scientific judgment of the overall evidence for activity of ST in an
15 assay.

16 **D.2.3. Results**

17 At the end of the 3-month studies, peripheral blood samples were obtained from male and female
18 rats and mice and analyzed for the frequency of micronucleated reticulocytes and erythrocytes
19 (Table D-2, Table D-3). In male and female rats, the reticulocyte population (PCEs), which is the
20 only red blood cell population that can be accurately assessed for micronucleus frequency in
21 peripheral blood of rats due to efficient splenic scavenging of damaged erythrocytes, did not
22 show an increase in micronucleated cells following 3 months of exposure to ST via drinking
23 water (0, 125, 250, 500, 1,000, and 2,000 mg/L) (Table D-2). Significant increases in the percent
24 reticulocytes were seen in both male and female rats, suggesting that ST could have stimulated
25 erythropoiesis in the bone marrow; however, the absolute increases in the percentages were small
26 compared to the vehicle control animals.

27 In male and female mice, there were no significant increases in micronucleated reticulocytes or
28 in micronucleated erythrocytes in either sex following 3 months of exposure to ST via drinking
29 water (0, 125, 250, 500, 1,000, and 2,000 mg/L) (Table D-3). A significant increase in the
30 percent reticulocytes was seen in male mice suggesting that ST could have stimulated
31 erythropoiesis in the bone marrow; however, the absolute increase was small compared to the
32 vehicle control group.

1 **Table D-2. Frequency of Micronuclei in Peripheral Blood Erythrocytes of Male and Female Rats in**
 2 **the Three-month Drinking Water Study of Sodium Tungstate Dihydrate^a**

	Number of Rats with Erythrocytes Scored	Micronucleated PCEs/1,000 PCEs ^b	P Value ^c	Micronucleated NCEs/1,000 NCEs ^b	P Value ^c	PCEs (%) ^b	P Value ^c
Male							
Exposure Concentration (mg/L)							
0	5	0.379 ± 0.04		0.055 ± 0.01		1.048 ± 0.08	
125	5	0.447 ± 0.06	0.483	0.037 ± 0.01	0.934	1.138 ± 0.08	1.000
250	5	0.320 ± 0.10	0.563	0.034 ± 0.01	0.969	0.926 ± 0.04	1.000
500	5	0.450 ± 0.03	0.386	0.051 ± 0.01	0.978	1.036 ± 0.04	1.000
1,000	5	0.450 ± 0.06	0.400	0.033 ± 0.00	0.983	1.182 ± 0.08	0.222
2,000	5	0.390 ± 0.09	0.412	0.025 ± 0.00	0.985	1.318 ± 0.08	0.016
Trend ^d		p = 0.439		p = 0.988		p = 0.003	
Female							
Exposure Concentration (mg/L)							
0	5	0.520 ± 0.03		0.074 ± 0.02		0.891 ± 0.11	
125	5	0.560 ± 0.14	0.649	0.038 ± 0.00	0.861	0.799 ± 0.08	1.000
250	5	0.488 ± 0.05	0.735	0.064 ± 0.02	0.921	1.050 ± 0.10	0.278
500	5	0.520 ± 0.04	0.770	0.056 ± 0.01	0.939	1.085 ± 0.14	0.275
1,000	5	0.506 ± 0.09	0.787	0.050 ± 0.01	0.948	1.240 ± 0.07	0.042
2,000	5	0.280 ± 0.04	0.800	0.045 ± 0.01	0.954	1.191 ± 0.09	0.043
Trend		p = 0.993		p = 0.853		p = 0.012	

3 PCE = polychromatic erythrocyte; NCE = normochromatic erythrocyte.

4 ^aStudy was performed at Integrated Laboratory Systems, LLC.

5 ^bData presented as mean ± standard error.

6 ^cPairwise comparisons with the vehicle control group performed using the Williams or Dunn test ($p \leq 0.025$).

7 ^dExposure concentration-related trends evaluated by linear regression of the Jonckheere test ($p \leq 0.025$).

1 **Table D-3. Frequency of Micronuclei in Peripheral Blood Erythrocytes of Male and Female Mice in**
 2 **the Three-month Drinking Water Study of Sodium Tungstate Dihydrate^a**

	Number of Mice with Erythrocytes Scored	Micronucleated PCEs/1,000 PCEs ^b	P Value ^c	Micronucleated NCEs/1,000 NCEs ^b	P Value ^c	PCEs (%) ^b	P Value ^c
Male							
Exposure Concentration (mg/L)							
0	5	2.530 ± 0.22		1.433 ± 0.05		1.558 ± 0.04	
125	5	2.690 ± 0.11	0.299	1.467 ± 0.04	1.000	1.444 ± 0.06	1.000
250	5	2.830 ± 0.19	0.357	1.491 ± 0.04	1.000	1.552 ± 0.08	1.000
500	5	2.840 ± 0.30	0.381	1.559 ± 0.04	0.181	1.608 ± 0.04	0.939
1,000	5	2.380 ± 0.23	0.395	1.504 ± 0.07	1.000	1.547 ± 0.04	0.955
2,000	5	2.700 ± 0.12	0.383	1.470 ± 0.01	1.000	1.745 ± 0.03	0.021
Trend ^d		p = 0.580		p = 0.240		p = 0.002	
Female							
Exposure Concentration (mg/L)							
0	5	1.810 ± 0.18		0.909 ± 0.01		1.302 ± 0.13	
125	5	2.103 ± 0.15	0.339	0.952 ± 0.04	0.293	1.938 ± 0.20	0.311
250	5	1.940 ± 0.09	0.875	0.992 ± 0.02	0.350	1.568 ± 0.20	0.373
500	5	1.730 ± 0.13	1.000	0.890 ± 0.01	0.373	1.613 ± 0.22	0.400
1,000	5	1.960 ± 0.27	1.000	0.934 ± 0.04	0.387	1.187 ± 0.08	0.411
2,000	5	1.930 ± 0.08	0.922	0.874 ± 0.02	0.398	1.509 ± 0.13	0.418
Trend		p = 0.522		p = 0.968		p = 0.567	

3 PCE = polychromatic erythrocyte; NCE = normochromatic erythrocyte

4 ^aStudy was performed at Integrated Laboratory Systems, LLC.

5 ^bData presented as mean ± standard error.

6 ^cPairwise comparisons with the vehicle control group performed using the Williams or Dunn test ($p \leq 0.025$).

7 ^dExposure concentration-related trends evaluated by linear regression of the Jonckheere test ($p \leq 0.025$).

8 **D.3. Comet Assay**

9 **D.3.1. Comet Assay Protocol**

10 For preparation of samples for the comet assay, a 50 μ L sample of blood was transferred to a
 11 tube containing 1 mL of freshly prepared cold mincing buffer [Mg^{+2} , Ca^{+2} , and phenol-free
 12 Hank's Balanced Salt Solution (Life Technologies, Carlsbad, CA) with 20 mM EDTA, pH 7.3 to
 13 7.5, and 10% v/v fresh dimethyl sulfoxide (DMSO)]. The ileum, liver, and kidney were rinsed
 14 with cold mincing buffer to remove residual blood and were held on ice briefly (≤ 5 minutes)
 15 until processed. Small portions (3 to 4 mm) of the ileum, liver, and kidney were placed in tubes
 16 containing cold mincing solution and rapidly minced until finely dispersed. All samples prepared
 17 for the comet assay were immediately flash frozen in liquid nitrogen¹⁰² and subsequently
 18 transferred to a $-80^{\circ}C$ freezer for storage until shipment by overnight courier on dry ice to the

1 analytical laboratory. Upon receipt, all samples were immediately placed in a -80°C freezer for
2 storage until further processing.

3 Blood and tissue samples were thawed on ice and maintained on ice during slide preparation.
4 Just before use, each cell suspension was shaken gently to mix the cells and placed back on ice
5 for 15 to 30 seconds to allow clumps to settle. A portion of the supernatant was empirically
6 diluted with 0.5% low melting point agarose (Lonza, Walkersville, MD) dissolved in Dulbecco's
7 phosphate buffer (Ca^{+2} , Mg^{+2} , and phenol-free) at 37°C and layered onto each well of a 2-well
8 CometSlide™ (Trevigen, Gaithersburg, MD). Slides were immersed in cold lysing solution
9 [2.5 M NaCl, 100 mM Na_2EDTA , 10 mM tris(hydroxymethyl)aminomethane (Tris), pH 10,
10 containing freshly added 10% DMSO (Fisher Scientific, Pittsburgh, PA), and 1% Triton X-100]
11 overnight in a refrigerator, protected from light. The following day, the slides were rinsed in
12 0.4 M Trizma base (pH 7.5), randomly placed onto the platform of a horizontal electrophoresis
13 unit, and treated with cold alkali solution (300 mM NaOH, 1 mM Na_2EDTA , pH>13) for
14 20 minutes to allow DNA unwinding, then electrophoresed at 4°C to 9°C for 20 minutes at 25 V
15 (0.7 V/cm), with a current of approximately 300 mA. After electrophoresis, slides were
16 neutralized with 0.4 M Trizma base (pH 7.5) for 5 minutes and then dehydrated by immersion in
17 absolute ethanol (Pharmco-AAPER, Shelbyville, KY) for at least 5 minutes and allowed to air
18 dry. Slides were prepared in a laboratory with a relative humidity no more than 60% and stored
19 at room temperature in a desiccator with a relative humidity of no more than 60% until stained
20 and scored; stained slides were stored in a desiccator. NaCl, Na_2EDTA , Triton X-100, and
21 Trizma base were purchased from Sigma-Aldrich (St. Louis, MO); NaOH was purchased from
22 Fisher Scientific (Pittsburgh, PA).

23 After staining with SYBR® Gold (Molecular Probes, Life Technologies, Grand Island, NY), the
24 slides, independently coded to mask treatment, were scored using Comet Assay IV Imaging
25 Software, Version 4.3.1 (Perceptive Instruments, Ltd., Suffolk, UK), validated for Good
26 Laboratory Practice Part 11 compliance. In the alkaline (pH > 13) comet assay, damaged nuclear
27 DNA fragments undergo unidirectional migration through the agarose gel within an electrical
28 field, forming an image that resembles a comet, and the greater the amount of fragmentation, the
29 greater the amount of DNA migration that will occur. The image analysis software partitions the
30 intensity of the fluorescent signal of the DNA in the entire comet image into the percent that is
31 attributable to the comet head and the percent attributable to the tail. Manual adjustment of the
32 automated detection of head and tail features is sometimes required. To evaluate DNA damage
33 levels, the extent of DNA migration was characterized for 100 scorable comet figures per
34 animal/tissue as percent tail DNA (intensity of all tail pixels divided by the total intensity of all
35 pixels in the comet, expressed as a percentage).

36 **D.3.2. Results**

37 In addition to evaluating the potential for chromosomal damage, the potential for DNA damage
38 was assessed using the comet assay in the same animals in which micronucleus induction was
39 evaluated. DNA damage from exposure to ST was assessed in liver, ileum, and kidney cell
40 samples, and in blood leukocytes (Table D-4, Table D-5). Significant increases in DNA damage,
41 measured as percent tail DNA, were observed in liver cells from male and female rats. Increases
42 in DNA damage were not observed for peripheral blood leukocytes in male or female rats, or for
43 ileum cells in female rats. Although cells from kidney tissue were evaluated from male and
44 female rats, and ileum tissue was evaluated from male rats, results from these tissues were

1 considered invalid due to unusually high levels of percent tail DNA in the control group. In male
2 mice, significant increases in DNA damage were observed in liver and ileum cells, but not in
3 kidney cells or peripheral blood leukocytes. No increases in percent tail DNA were observed in
4 female mice for liver, kidney, ileum, or peripheral blood leukocyte cells.

1 **Table D-4. DNA Damage in Male and Female Rats Exposed to Sodium Tungstate Dihydrate in Drinking Water for Three Months^a**

Exposure Concentration (mg/L)	Blood		Ileum		Kidney		Liver	
	Percent Tail DNA ^b	P Value ^c	Percent Tail DNA ^b	P Value	Percent Tail DNA ^b	P Value	Percent Tail DNA ^b	P Value
Male								
0	1.32 ± 0.22		IT		IT		3.22 ± 0.12	
125	1.15 ± 0.15	0.615	IT		IT		11.00 ± 1.44	0.078
250	1.21 ± 0.14	0.702	IT		IT		4.87 ± 0.70	1.000
500	1.59 ± 0.11	0.313	IT		IT		16.06 ± 1.43	0.003
1,000	1.49 ± 0.21	0.323	IT		IT		14.74 ± 1.43	0.010
2,000	1.77 ± 0.39	0.109	IT		IT		20.02 ± 2.08	<0.001
Trend ^d	p = 0.021		–		–		p < 0.001	
Female								
0	1.48 ± 0.19		17.38 ± 4.29		IT		9.84 ± 0.61	
125	1.58 ± 0.17	0.679	24.02 ± 2.57	0.445	IT		13.53 ± 1.63	0.460
250	1.28 ± 0.36	0.764	16.00 ± 1.73	0.522	IT		17.02 ± 2.57	0.049
500	1.35 ± 0.13	0.798	15.58 ± 3.15	0.555	IT		14.51 ± 1.07	0.131
1,000	1.10 ± 0.20	0.814	17.07 ± 2.73 ^e	0.588	IT		14.32 ± 1.87	0.196
2,000	1.22 ± 0.18	0.827	16.95 ± 2.69	0.587	IT		22.47 ± 1.91	<0.001
Trend	p = 0.892		p = 0.741		–		p < 0.001	

2 IT = invalid test due to unusually high control percent tail DNA.

3 ^aStudy was performed at Integrated Laboratory Systems, LLC.4 ^bData presented as mean ± standard error; n = 5.5 ^cPairwise comparisons with the vehicle control group performed using the Williams or Dunn test (p ≤ 0.025).6 ^dExposure concentration-related trends evaluated by linear regression of the Jonckheere test (p ≤ 0.025).7 ^en = 4.

1 **Table D-5. DNA Damage in Male and Female Mice Exposed to Sodium Tungstate Dihydrate in Drinking Water for Three Months^a**

Exposure Concentration (mg/L)	Blood		Ileum		Kidney		Liver	
	Percent Tail DNA ^b	P Value ^c	Percent Tail DNA ^b	P Value	Percent Tail DNA ^b	P Value	Percent Tail DNA ^b	P Value
Male								
0	2.49 ± 0.43		11.69 ± 2.34		6.20 ± 0.57		4.14 ± 0.32 ^e	
125	3.90 ± 0.48	0.049	20.26 ± 1.91	0.019	5.48 ± 0.73	0.838	17.55 ± 0.50	<0.001
250	3.86 ± 0.54	0.058	13.27 ± 0.68	0.023	5.07 ± 0.70	0.904	17.51 ± 1.25	<0.001
500	3.29 ± 0.37	0.061	22.91 ± 1.17	0.023	4.84 ± 0.34	0.924	17.01 ± 1.03	<0.001
1,000	3.34 ± 0.54	0.062	9.35 ± 1.56	0.023	5.67 ± 0.54	0.867	19.86 ± 1.02	<0.001
2,000	3.61 ± 0.33	0.060	19.96 ± 0.91	< 0.001	6.63 ± 0.38	0.402	19.97 ± 1.25	<0.001
Trend ^d	p = 0.199		p = 0.212		p = 0.308		p <0.001	
Female								
0	0.93 ± 0.19		20.50 ± 1.80		6.81 ± 1.87		6.40 ± 0.52	
125	1.24 ± 0.24	0.830	21.85 ± 1.80	0.340	4.84 ± 0.47	1.000	5.92 ± 0.89	0.573
250	1.03 ± 0.11	1.000	21.07 ± 1.85	0.404	8.54 ± 3.13	1.000	10.63 ± 1.22	0.053
500	1.04 ± 0.23	1.000	24.65 ± 1.01	0.131	5.36 ± 0.59	1.000	7.99 ± 1.14	0.056
1,000	1.28 ± 0.29	1.000	23.91 ± 1.94	0.134	12.16 ± 3.75	0.403	8.90 ± 1.00	0.056
2,000	1.42 ± 0.09	0.167	21.88 ± 1.05	0.137	7.01 ± 1.02	1.000	7.50 ± 0.64	0.056
Trend	p = 0.081		p = 0.112		p = 0.057		p = 0.105	

2 ^aStudy was performed at Integrated Laboratory Systems, LLC.3 ^bData presented as mean ± standard error; n = 5.4 ^cPairwise comparisons with the vehicle control group performed using the Williams or Dunn test (p ≤ 0.025).5 ^dExposure concentration-related trends evaluated by linear regression of the Jonckheere test (p ≤ 0.025).6 ^en = 4.

1 **Appendix E. Tungsten Concentration Determination**

2 **Table of Contents**

3	E.1. Sample Collection	E-2
4	E.2. Sample Analysis	E-2
5	E.3. Analysis of Xanthine and Methionine	E-4

6

7 **Tables**

8	Table E-1. Analytical Method Validation and Stability Data for Tungsten in Blood and	
9	Urine for the Three-month Studies	E-5
10	Table E-2. Analytical Method Validation and Stability Data for Tungsten in Urine for the	
11	Two-year Studies	E-6
12	Table E-3. Analytical Method Validation and Stability Data for Tungsten in Plasma for	
13	the Two-year Studies	E-7
14	Table E-4. Analytical Method Validation and Stability Data for Tungsten in Kidney for	
15	the Two-year Studies	E-8
16	Table E-5. Analytical Method Validation and Stability Data for Xanthine and Methionine	
17	in Rat Urine.....	E-9

18

1 **E.1. Sample Collection**

2 **E.1.1. Three-month Studies**

3 ***E.1.1.1. Urine***

4 At week 12, all F₁ rats were removed from exposure and placed in individual metabolism cages
5 to allow for collection of urine. Rats were fasted during the collection period and had access to
6 untreated water ad libitum. Urine samples were collected on wet ice, overnight, for
7 approximately 16 hours. Urine samples (1 mL from each rat) were frozen at –20°C and shipped
8 to the analytical laboratory (Battelle Toxicology Northwest, Richland, WA).

9 ***E.1.1.2. Blood***

10 At study termination, rats and mice were anesthetized with a CO₂/O₂ mixture, and blood was
11 collected from the retroorbital plexus (rats) or sinus (mice) into heparinized centrifuge tubes
12 containing ethylenediaminetetraacetic acid (EDTA). Blood samples were frozen at –80°C and
13 shipped to Battelle Toxicology Northwest (Richland, WA).

14 **E.1.2. Two-year Studies (Interim Evaluations)**

15 At the beginning of the 2-year study, groups of 40 interim study male and female rats and mice
16 were randomly assigned to the tissue distribution study and exposed identically to the core study
17 groups.

18 At the 3-, 6-, 12-, and 18-month interim evaluations, urine, feces, blood, and tissues (liver,
19 kidneys, stomach, small intestine, and bone) were collected from up to 10 predesignated
20 F₁ rats/sex/exposure group and up to 10 predesignated mice/sex/exposure group. Early death
21 animals were not replaced. On the morning of the day before scheduled blood collection, animals
22 were moved to metabolism cages (one animal per cage); while in the metabolism cages, the
23 animals had ad libitum access to feed and their assigned concentration of dosed drinking water.
24 Urine and feces were collected over a 24-hour period. Blood was collected via cardiac puncture
25 into tubes containing K₃ EDTA, centrifuged, and the plasma harvested. Immediately after blood
26 collection, the animals were euthanized and the entire liver, both kidneys, stomach (separated
27 into glandular and non-glandular), small intestine, and both femurs were collected, weighed, and
28 maintained on dry ice until moved into storage. All samples were stored at –85°C to –60°C until
29 shipped to Battelle Toxicology Northwest (Richland, WA).

30 **E.2. Sample Analysis**

31 **E.2.1. Three-month Studies**

32 Tungsten concentrations in samples were quantified using validated analytical methods, and
33 method validation data are given in Table E-1. Blood and urine samples from the study were
34 stored at –70°C and –20°C, respectively, following collection. Samples were allowed to thaw at
35 room temperature and mixed well. A 0.1 mL aliquot of blood was transferred into a Teflon
36 digestion tube (CEM Corporation, Matthews, NC) and 0.15 mL of concentrated HNO₃, 1.5 mL
37 of deionized water, and 0.1 mL of 10 µg/mL bismuth (internal standard) in approximately 1%
38 HNO₃ (using 1 mg/mL procured from SPEX CertiPrep, Metuchen, NJ) was added. Urine
39 samples were prepared similarly to blood except that to 0.1 mL of urine, 0.375 mL of

1 concentrated HNO₃, 1 mL of deionized water, and 0.25 mL of 0.2 µg/mL bismuth were added.
2 The final acid strength of samples was approximately 1% HNO₃. Samples were capped, allowed
3 to stand for approximately 15 minutes, and digested at 1600 W and 200°C for 40 minutes for
4 blood or 20 minutes for urine using a MARS5 Microwave Digestion System (CEM Corporation,
5 Mathews, NC). Samples were cooled to room temperature and diluted to 10 mL with water.

6 Corresponding matrix calibration standards, blanks, and quality control (QC) standards were
7 prepared and analyzed similarly to study samples. Calibration curves were run on blood with six
8 calibration standards (blood, 0.1 to 100 µg tungsten/L; urine, 0.5 to 150 µg tungsten/L). Blood
9 QC standards were prepared at 0.5 and 50 µg tungsten/L, and urine QC standards were prepared
10 at 2 and 100 µg/L. Study samples with responses greater than the highest calibration standard
11 were diluted with 1% HNO₃ such that the final concentration in the sample was within the
12 validated range. All samples were analyzed for tungsten concentration using an inductively
13 coupled plasma-mass spectrometry (ICP-MS) method as described below.

14 **E.2.2. Two-year Studies (Interim Evaluations)**

15 Tungsten concentrations in samples were quantified using validated analytical methods and
16 corresponding validation data are given in Table E-2, Table E-3, and Table E-4 for urine,
17 plasma, and kidney, respectively. All study samples were stored at -70°C following collection.
18 Plasma and urine samples were prepared using a method similar to the 3-month studies. Kidney
19 samples were weighed and homogenized with approximately 5 volumes of water for 5 minutes
20 using a polytron homogenizer. The homogenate was sonicated for approximately 30 minutes and
21 then vortexed to mix. A 0.1-mL aliquot of the homogenate was digested similar to other matrices
22 as before.

23 Corresponding solvent calibration standards, blanks, and matrix QC standards were prepared and
24 analyzed similarly to study samples. Calibration curves were run with six calibration standards
25 (0.1 to 100 µg tungsten/mL to quantitate plasma; 0.5 to 150 µg tungsten/mL to quantitate urine;
26 60 to 2,500 ng/mL to quantitate kidney homogenate). QC samples were prepared in all three
27 matrices at 75 and 1,875 µg tungsten/mL. Study samples with responses greater than the highest
28 calibration standard were diluted with 1% HNO₃ such that final concentration in the sample was
29 within the validated range. All samples were analyzed for tungsten concentration using an
30 ICP-MS method as described below.

31 **E.2.3. Instrumentation and Quantitation**

32 Samples from the 3-month studies were analyzed using an Agilent 7500ce (Agilent, Palo Alto,
33 CA) ICP-MS. The detector mode was set to auto with a total acquisition time of 3.3 seconds. The
34 ions monitored were m/z 182 and 209 for tungsten and bismuth, respectively. Samples from the
35 2-year studies were analyzed using a Perkin Elmer NexION 300 (Waltham, MA) ICP-MS. The
36 detector mode was set to dual and total acquisition time of 4.008 seconds. The ions monitored
37 were m/z 181.948 and 208.980 for tungsten and bismuth, respectively.

38 The performance of the calibration curve was evaluated before the analysis of each sample set. A
39 successful calibration was indicated by the following: correlation coefficient (r) ≥ 0.98 ; relative
40 standard deviation (RSD) less than or equal to $\pm 15\%$ (except at the limit of quantitation (LOQ)
41 where RSD was less than or equal to $\pm 20\%$); relative error (RE) less than or equal to $\pm 15\%$
42 (except at LOQ where RE was less than or equal to $\pm 20\%$).

1 Calibration curves relating response ratio of analyte to internal standard (following correction for
2 the background concentrations for the 3-month studies only) and concentration of tungsten in
3 matrix were constructed using a 1/X weighted linear regression. The concentrations of tungsten
4 in samples were calculated using response ratio, the regression equation, initial sample weight or
5 volume, digestion volume, and dilution when applicable. The concentrations were reported as μg
6 tungsten/mL (μg tungsten/g in 3-month studies) for blood, plasma, and urine and μg tungsten/g
7 for kidney in the 2-year studies.

8 Data from study samples were considered valid if they were bracketed by valid QC sets. In
9 general, each sample set, method blanks, and controls were bracketed by two QC sets, which
10 consisted of a calibration blank and two concentrations of calibration standards (QC low and QC
11 high). A QC set passed when the measured concentration for QC standards were within 20% of
12 its nominal value. If the QC standard failed, it was necessary to reanalyze the bracketed samples.
13 All QC standards were within 20% of nominal concentrations.

14 **E.3. Analysis of Xanthine and Methionine**

15 Xanthine and methionine concentrations in rat urine were quantified using validated analytical
16 methods using a standard addition approach. Validation data are listed in Table E-5. Study
17 samples were stored at -20°C until used. Samples were allowed to thaw at room temperature and
18 mixed well. Five 50 μL aliquot samples were transferred to individual wells in a 96-well plate,
19 and 50 μL aliquots of the spiking standards of either methionine or xanthine (target
20 concentrations 0, 0.075, 0.15, 0.3, 0.6 $\mu\text{g}/\text{mL}$) were added. After the addition of internal standard
21 (400 μL of 1,500 ng/mL $^{15}\text{N}_2$ xanthine or 2 $\mu\text{g}/\text{mL}$ $^2\text{H}_3$ methionine), the plate was covered with a
22 pierceable sealing mat and the samples were mixed for approximately 10 seconds.

23 All samples were analyzed by liquid chromatography with tandem mass spectrometry using
24 either an Agilent (Santa Clara, CA) or Shimadzu liquid chromatograph coupled to a Sciex
25 (Toronto, Ontario, Canada) 4,000 or 5,000 mass spectrometer. For analysis of xanthine,
26 chromatography was performed using a Phenomenex (Torrance, CA) Synergi Hydro RP
27 (2×50 mm) column. Mobile phases A (water with 1 mM ammonium acetate, 0.1% formic acid)
28 and B (acetonitrile with 0.1% formic acid) were run at a flow rate of 0.275 mL/minute with a
29 linear gradient from 3% B to 95% B over 3 minutes followed by a 2.5-minute hold. The
30 turbospray ion source was operated in negative mode. Transitions monitored for xanthine and
31 $^{15}\text{N}_2$ xanthine were 151: >108 and 153: >109, respectively. For analysis of methionine,
32 chromatography was performed using a Phenomenex (Torrance, CA) Develosil C30 (50×2 mm)
33 column. Mobile phases were the same as for analysis of xanthine and were run at a flow rate of
34 0.275 mL/minute with a linear gradient from 3% B (3 minutes) to 95% B over 1 minute followed
35 by a 0.5-minute hold. The turbospray ion source was operated in negative mode. Transitions
36 monitored for methionine and $^2\text{H}_3$ methionine were 150: >104 and 153: >107, respectively.

37 Calibration curves relating response ratio of analyte to internal standard and concentration of
38 xanthine or methionine in matrix were constructed using linear regression. The xanthine
39 concentration in each was determined as the negative x-intercept of the standard addition curve.
40 The concentration was reported as $\mu\text{g}/\text{mL}$ of urine. Data from study samples were considered
41 valid if the QCs were within 30% of the nominal values.

1 **Table E-1. Analytical Method Validation and Stability Data for Tungsten in Blood and Urine for**
 2 **the Three-month Studies^a**

Validation Parameter	Rat Blood	Rat Urine	Mouse Blood
Matrix Concentration Range ($\mu\text{g/g}$)	0.01–10	0.002–3	–
LOQ ($\mu\text{g/g}$)	0.0068	0.0018	–
LOD ($\mu\text{g/g}$)	0.002	0.0054	–
Correlation Coefficient (r)	≥ 0.999	≥ 0.999	–
Recovery (%) ^b	101.3–117.2	57.1 – 127.5	–
Precision and Accuracy ^{c,d}			
Intra-day % RSD	≤ 7.9	≤ 4.6	≤ 6.8
Intra-day % RE	$\leq \pm 7.1$	$\leq \pm 4.4$	$\leq \pm 3.1$
Inter-day % RSD	≤ 5.9	≤ 4.0	–
Inter-day % RE	$\leq \pm 5.0$	$\leq \pm 1.9$	–
Dilution Verification			
	up to 50 $\mu\text{g/g}$	up to 30 $\mu\text{g/g}$	up to 100 $\mu\text{g/g}$
% RSD	4.4	1.3	2.8
% RE	–10.5	–14.0	–10.8
Extract Stability (Ambient Storage) (Average % RE) ^d			
	$\leq \pm 2.5$ up to 7 days	$\leq \pm 4.3$ up to 3 days	$\leq \pm 1.5$ up to 8 days
Matrix Stability (Average % RE) ^d			
Freeze-thaw (three cycles)	$\leq \pm 9.4$	$\leq \pm 6.4$	$\leq \pm 6.3$
Frozen matrix (-70°C for blood and -20°C for urine, up to approximately 60 days)	$\leq \pm 11.8$	$\leq \pm 10.0$	$\leq \pm 7.6$

3 LOQ = limit of quantitation; LOD = limit of detection; RSD = relative standard deviation; RE = relative error.

4 ^aMethod was fully validated in Sprague Dawley rat blood and urine and cross-validated in B6C3F1/N mouse blood using quality
 5 control (QC) samples prepared in mouse blood at three concentrations (0.05, 1, 5 μg tungsten/g) and analyzed using a rat blood
 6 calibration curve.

7 ^bEstimated by comparing response of matrix standards to solvent standards.

8 ^cPrecision was estimated as % RSD. Accuracy was estimated as average % RE.

9 ^dDetermined for six replicate QCs at three levels: 0.05, 1, and 5 μg tungsten/g for blood and 0.01, 0.2, and 2 μg tungsten/g for
 10 urine.

1 **Table E-2. Analytical Method Validation and Stability Data for Tungsten in Urine for the Two-year**
 2 **Studies^a**

Validation Parameter	Rat	Mouse
Solvent Standard Concentration Range (ng/mL)	60–2,500	60–2,500
LOQ (ng/mL)	60	60
LOD (ng/mL)	13.0	13.0
Correlation Coefficient (r)	≥0.998	≥0.999
Recovery (%) ^b	102–105	104–117
Precision and Accuracy ^{c,d,e}		
Intra-day % RSD	≤7.1	≤2.9
Intra-day % RE	≤ ± 11.9	≤ ± 29.3
Inter-day % RSD	≤5.8	–
Inter-day % RE	≤ ± 10.6	–
Dilution Verification – Water Predigestion		
	up to 100,000 ng/mL	up to 100,000 ng/mL
% RSD	1.4	3.3
% RE	7.9	–5.1
Dilution Verification – Acid Postdigestion		
	up to 100,000 ng/mL	up to 100,000 ng/mL
% RSD	2.9	3.4
% RE	4.0	4.0
Matrix Evaluation ^{d,e,f}		
Sprague Dawley rat % RSD	≤11.4	–
Sprague Dawley rat % RE	≤ ± 27.0	–
B6C3F1/N mouse % RSD	–	≤3.6
B6C3F1/N mouse % RE	–	≤ ± 17.1
Postpreparative Stability (ambient storage) (% RE) ^{e,g}	≤ ± 15.8 up to 16 days	≤ ± 9.3 up to 16 days
Matrix Stability (% RE) ^{d,e}		
Freeze-thaw (four cycles)	≤ ± 11.7	≤ ± 2.7
Frozen matrix (–70°C up to approximately 61 days)	≤ ± 15.0	≤ ± 12.8

3 LOQ = limit of quantitation; LOD = limit of detection; RSD = relative standard deviation; RE = relative error.

4 ^aMethod was fully validated in Sprague Dawley rat urine using a solvent standard curve and cross-validated in B6C3F1 mouse
 5 urine using quality control (QC) samples prepared in mouse urine at three concentrations (75, 375, 1,875 ng tungsten/mL) and
 6 analyzed using solvent curve.

7 ^bEstimated by comparing response of matrix QC samples to solvent QC samples.

8 ^cPrecision was estimated as % relative standard deviation (RSD). Accuracy was estimated as average % relative error (RE).

9 ^dDetermined for six replicate QCs at three concentrations: 75, 375, and 1,875 ng tungsten/mL for rat and mouse urine.

10 ^eCorrected for endogenous tungsten in urine.

11 ^fStudy samples matrices were assessed using six replicate QCs at three concentrations: 75,375,1875 ng tungsten/mL.

12 ^gDetermined for six replicate QCs at three concentrations: 75, 375, and 1,875 ng tungsten/mL for rat and mouse urine evaluated
 13 using freshly extracted standards and stored standard extracts.

1 **Table E-3. Analytical Method Validation and Stability Data for Tungsten in Plasma for the**
 2 **Two-year Studies^a**

Validation Parameter	Rat	Mouse
Solvent Concentration Range (ng/mL)	60–2,500	60–2,500
LOQ (ng/mL)	60	60
LOD (ng/mL)	13.0	13.0
Correlation Coefficient (r)	≥0.99	≥0.99
Recovery (%) ^b	103–117	90.5–101
Precision and Accuracy ^{c,d,e}		
Intra-day % RSD	≤5.1	≤2.9
Intra-day % RE ^e	≤ ± 12.9	≤ ± 14.6
Inter-day % RSD	–	–
Inter-day % RE	–	–
Dilution Verification – Water Predigestion		
	up to 25,000 ng/mL	up to 25,000 ng/mL
% RSD	2.5	2.2
% RE	5.2	5.2
Dilution Verification – Acid Postdigestion		
	up to 25,000 ng/mL	up to 25,000 ng/mL
% RSD	2.2	1.2
% RE	–1.6	10.4
Matrix Evaluation ^{d,e,f}		
Sprague Dawley rat % RSD	≤4.7	–
Sprague Dawley rat % RE	≤ ± 17.4	–
B6C3F1/N mouse % RSD	–	≤3.5
B6C3F1/N mouse % RE	–	≤ ± 19.1
Postpreparative Stability (Ambient Storage) (% RE) ^{e,g}	≤ ± 18.4 up to 14 days	≤ ± 18.2 up to 14 days
Matrix Stability (Average % RE) ^{d,e}		
Freeze-thaw (four cycles)	≤ ± 9.3	≤ ± 11.8
Frozen matrix (–70°C up to approximately 60 days)	≤ ± 16.2	≤ ± 18.3

3 LOQ = limit of quantitation; LOD = limit of detection; RSD = relative standard deviation; RE = relative error.

4 ^aMethod was fully validated in Sprague Dawley rat urine using a solvent curve and crossed-validated in rat plasma and B6C3F1
 5 mouse plasma using three concentrations (75, 375, 1,875 ng tungsten/mL) of quality control (QC) samples prepared in rat and
 6 mouse plasma and analyzed using solvent curve.

7 ^bEstimated by comparing response of matrix sample to solvent sample.

8 ^cPrecision was estimated as % relative standard deviation (RSD). Accuracy was estimated as average % relative error (RE).

9 ^dDetermined for six replicate QCs at three concentrations: 75, 375, and 1,875 ng tungsten/mL for rat and mouse plasma.

10 ^eCorrected for endogenous tungsten in plasma.

11 ^fStudy samples matrices were assessed using six replicate QCs at three concentrations: 75, 375, 1,875 ng tungsten/mL.

12 ^gDetermined for six replicate QCs at three concentrations: 75, 375, and 1,875 ng tungsten/mL for rat and mouse plasma evaluated
 13 using freshly extracted standards and stored standard extracts.

1 **Table E-4. Analytical Method Validation and Stability Data for Tungsten in Kidney for the**
 2 **Two-year Studies^a**

Validation Parameter	Rat	Mouse
Matrix Concentration Range (ng/mL)	60–2,500	60–2,500
LOQ (ng/mL)	60	60
LOD (ng/mL)	13.0	13.0
Correlation Coefficient (r)	≥0.99	≥0.99
Recovery (%) ^b	118–120	111–124
Precision and Accuracy ^{c,d,e}		
Intra-day % RSD	≤1.2	≤2.1
Intra-day % RE	≤ ± 14.1	≤ ± 11.5
Inter-day % RSD	–	–
Inter-day % RE	–	–
Dilution Verification – Water Predigestion		
	up to 125,000 ng/g	up to 125,000 ng/g
% RSD	2.1	2.6
% RE	5.6	7.2
Dilution Verification – Acid Postdigestion		
	up to 125,000 ng/g	up to 125,000 ng/g
% RSD	1.7	1.8
% RE	–3.2	–3.2
Matrix Evaluation ^{d,e,f}		
Sprague Dawley rat % RSD	≤4.3	–
Sprague Dawley rat % RE	≤ ± 3.5	–
B6C3F1/N mouse % RSD	–	≤4.6
B6C3F1/N mouse % RE	–	≤ ± 9.9
Postpreparative Stability (Ambient Storage) (% RE) ^{e,g}	≤ ± 11.6 up to 28 days	≤ ± 10.8 up to 28 days
Matrix Stability (% RE) ^{d,e}		
Freeze-thaw (six cycles)	≤ ± 12.5	≤ ± 5.8
Frozen matrix (–70°C up to approximately 55 days)	≤ ± 10.3	≤ ± 11.9

3 LOQ = limit of quantitation; LOD = limit of detection; RSD = relative standard deviation; RE = relative error.

4 ^aMethod was fully validated in rat urine and cross-validated in Sprague Dawley rat and B6C3F1 and mouse kidney homogenate
 5 using quality control (QC) samples prepared in rat and mouse kidney homogenates at three concentrations (375, 1,875, and
 6 9,375 ng tungsten/g) and analyzed using solvent curve.

7 ^bEstimated by comparing response of matrix samples to solvent samples.

8 ^cPrecision was estimated as % relative standard deviation (RSD). Accuracy was estimated as average % relative error (RE).

9 ^dDetermined for six replicate QCs at three concentrations: 375, 1,875, and 9,375 ng tungsten/g for kidney homogenate.

10 ^eCorrected for endogenous tungsten in kidney.

11 ^fStudy samples matrices were assessed using six replicates of QCs at three concentrations: 75, 375, and 1,875 ng tungsten/g.

12 ^gDetermined for six replicate QCs at three concentrations: 75, 375, and 1,875 ng tungsten/mL for rat and mouse kidney evaluated
 13 using freshly extracted standards and stored standard extracts.

1 **Table E-5. Analytical Method Validation and Stability Data for Xanthine and Methionine in Rat**
 2 **Urine**

Parameter	Xanthine	Methionine
Matrix Concentration Range ($\mu\text{g/mL}$)	0.075–0.6	0.075–0.6
LOQ ($\mu\text{g/mL}$)	0.075	0.075
Correlation Coefficient (r)	≥ 0.994	≥ 0.999
Precision and Accuracy ^{a,b}		
Intra-day % RSD	≤ 27.2	≤ 12.5
Intra-day % RE	$\leq \pm 25.4$	$\leq \pm 13.5$
Inter-day % RSD	23.1	11.9
Inter-day % RE	-13.1	4.4
Stability (% RE) ^b		
Freeze-thaw (three cycles)	-29.7	-11.8
Frozen matrix (-20°C up to approximately 190 days)	-15.0	2.2

3 LOQ = limit of quantitation; RSD = relative standard deviation; RE = relative error.

4 ^aPrecision was estimated as % relative standard deviation (RSD). Accuracy was estimated as average % relative error (RE).

5 ^{b,c}Determined for six replicate quality control samples at 3 $\mu\text{g/mL}$ for xanthine and 1.5 $\mu\text{g/mL}$ for methionine.

1 **Appendix F. Peer-review Report**

2 [The peer-review report will appear in a future draft of this report.]

1 **Appendix G. Supplemental Data**

2 Tables with supplemental data can be found here: [https://doi.org/10.22427/NTP-DATA-TR-](https://doi.org/10.22427/NTP-DATA-TR-599)
3 [599](https://doi.org/10.22427/NTP-DATA-TR-599).⁸⁶

4 **G.1. Perinatal and Three-month Study in Rats**

- 5 E03 – Growth Curves
- 6 E04 – Mean Body Weights and Survival Table
- 7 E05 – Clinical Observations Summary
- 8 E07 – Mean Water Consumption by Treatment Group
- 9 E08 – Water and Compound Consumption Table
- 10 Gestational Body Weights (grams)
- 11 Gestational Water Consumption (grams)
- 12 Gestational and Lactational Chemical Consumption
- 13 Lactational Body Weights (grams)
- 14 Lactational Water Consumption (grams)
- 15 Litter Data by Dam – PND1
- 16 Live Litter Size and Survival – PND 4 and 21
- 17 P03 – Incidence Rates of Non–Neoplastic Lesions by Anatomic Site
- 18 P04 – Neoplasms by Individual Animal
- 19 P05 – Incidence Rates of Neoplasms by Anatomic Site (Systemic Lesions Abridged)
- 20 P09 – Non–Neoplastic Lesions by Individual Animal
- 21 P10 – Statistical Analysis of Non–Neoplastic Lesions
- 22 P14 – Individual Animal Pathology Data
- 23 P18 – Incidence Rates of Non–Neoplastic Lesions by Anatomic Site with Average Severity
24 Grades
- 25 P40 – Survival Curves
- 26 PA06 – Organ Weights Summary
- 27 PA41 – Clinical Chemistry Summary
- 28 PA43 – Hematology Summary

- 1 PA44 – Urinalysis Summary
- 2 PA48 – Tissue Concentration Summary
- 3 PA48C – Tissue Concentration Curve
- 4 Pup Body Weights (grams)
- 5 R02 – Reproductive Performance Summary
- 6 R06 – Andrology Summary
- 7 R07 – Hormone Summary
- 8 **G.2. Perinatal and Three-month Study in Rats – Individual Animal Data**
- 9 Female Individual Animal Body Weight Data
- 10 Female Individual Animal Clinical Observations
- 11 Female Individual Animal Non–Neoplastic Pathology Data
- 12 Female Individual Animal Survival Data
- 13 Female Individual Animal Terminal Body Weight Data
- 14 Male Individual Animal Body Weight Data
- 15 Male Individual Animal Non–Neoplastic Pathology Data
- 16 Male Individual Animal Survival Data
- 17 Male Individual Animal Terminal Body Weight Data
- 18 Individual Animal Andrology Data
- 19 Individual Animal Clinical Chemistry Data
- 20 Individual Animal DamID and PupID Data
- 21 Individual Animal Hematology Data
- 22 Individual Animal Hormone Data
- 23 Individual Animal Organ Weight Data
- 24 Individual Animal Reproductive Performance Data
- 25 Individual Animal Tissue Concentration Data
- 26 Individual Animal Urinalysis Data

1 **G.3. Three-month Study in Mice**

- 2 E03 – Growth Curves
- 3 E04 – Mean Body Weights and Survival Table
- 4 E05 – Clinical Observations Summary
- 5 E07 – Mean Water Consumption by Treatment Group
- 6 E08 – Water and Compound Consumption Table
- 7 P03 – Incidence Rates of Non–Neoplastic Lesions by Anatomic Site
- 8 P04 – Neoplasms by Individual Animal
- 9 P05 – Incidence Rates of Neoplasms by Anatomic Site (Systemic Lesions Abridged)
- 10 P09 – Non–Neoplastic Lesions by Individual Animal
- 11 P10 – Statistical Analysis of Non–Neoplastic Lesions
- 12 P14 – Individual Animal Pathology Data
- 13 P18 – Incidence Rates of Non–Neoplastic Lesions by Anatomic Site with Average Severity
- 14 Grades
- 15 P40 – Survival Curves
- 16 PA06 – Organ Weights Summary
- 17 PA43 – Hematology Summary
- 18 PA48 – Tissue Concentration Summary
- 19 PA48C – Tissue Concentration Curve
- 20 R06 – Andrology Summary

21 **G.4. Three-month Study in Mice – Individual Animal Data**

- 22 Female Individual Animal Body Weight Data
- 23 Female Individual Animal Clinical Observations
- 24 Female Individual Animal Non–Neoplastic Pathology Data
- 25 Female Individual Animal Survival Data
- 26 Female Individual Animal Terminal Body Weight Data
- 27 Male Individual Animal Body Weight Data
- 28 Male Individual Animal Non–Neoplastic Pathology Data

- 1 Male Individual Animal Survival Data
- 2 Male Individual Animal Terminal Body Weight Data
- 3 Individual Animal Andrology Data
- 4 Individual Animal Hematology Data
- 5 Individual Animal Organ Weight Data
- 6 Individual Animal Tissue Concentration Data
- 7 **G.5. Perinatal and Two-year Study in Rats**
- 8 Analysis of PND 1 Litter Data
- 9 Analysis of PND 4 Live Litter Size and Survival
- 10 E01 – Animal Removal Summary by Treatment Group
- 11 E02 – Animals Removed from Experiment
- 12 E03 – Growth Curves (Litter based)
- 13 E04 – Mean Body Weights and Survival (Litter based)
- 14 E05 – Clinical Observations Summary
- 15 E07 – Mean Water Consumption by Treatment Group
- 16 E08 – Water and Compound Consumption Table
- 17 E12 – Animal History
- 18 I05 – Clinical Observations Summary
- 19 Litter Data Analysis of Gestational Body Weight
- 20 Litter Data Analysis of Gestational Water Consumption
- 21 Litter Data Analysis of Gestational/Lactational Chemical Consumption
- 22 Litter Data Analysis of Lactational Body Weight
- 23 Litter Data Analysis of Lactational Water Consumption
- 24 Litter Data Analysis of Pup Body Weights
- 25 Litter Data Analysis of Pup Body Weights (cont'd)
- 26 P02 – Incidence Rates of Neoplasms by Anatomic Site
- 27 P03 – Incidence Rates of Non–Neoplastic Lesions by Anatomic Site
- 28 P04 – Neoplasms by Individual Animal

- 1 P05 – Incidence Rates of Neoplasms by Anatomic Site (Systemic Lesions Abridged)
- 2 P08 – Litter Statistical Analysis of Primary Tumors
- 3 P09 – Non–Neoplastic Lesions by Individual Animal
- 4 P10 – Statistical Analysis of Non–Neoplastic Lesions
- 5 P11 – Statistical Analysis of Survival Data
- 6 P14 – Individual Animal Pathology Data
- 7 P17 – Neoplasms by Individual Animal (Systemic Lesions Abridged)
- 8 P18 – Incidence Rates of Non–Neoplastic Lesions by Anatomic Site with Average Severity
- 9 Grades
- 10 P40 – Survival Curves
- 11 PA06 – Organ Weights Summary
- 12 PA48 – Summary of Tissue Concentration
- 13 PA48C – Tissue Concentration Curve
- 14 R02 – Reproductive Performance Summary

15 **G.6. Perinatal and Two-year Study in Rats – Individual Animal Data**

- 16 Female Individual Animal Body Weight Data
- 17 Female Individual Animal Clinical Observations
- 18 Female Individual Animal Neoplastic Pathology Data
- 19 Female Individual Animal Non–Neoplastic Pathology Data
- 20 Female Individual Animal Survival Data
- 21 Female Individual Animal Terminal Body Weight Data
- 22 Female Pup Individual Animal Body Weight Data
- 23 Male Individual Animal Body Weight Data
- 24 Male Individual Animal Clinical Observations
- 25 Male Individual Animal Neoplastic Pathology Data
- 26 Male Individual Animal Non–Neoplastic Pathology Data
- 27 Male Individual Animal Survival Data
- 28 Male Individual Animal Terminal Body Weight Data

- 1 Male Pup Individual Animal Body Weight Data
- 2 Individual Animal Clinical Observations Data
- 3 Individual Animal DamID and PupID Data
- 4 Individual Animal Organ Weight Data
- 5 Individual Animal Reproductive Performance Data
- 6 Individual Animal Tissue Concentration Data
- 7 Individual Animal Urinalysis Data
- 8 Individual Pup Census and Litter Weight by Sex Data
- 9 **G.7. Two-year Study in Mice**
- 10 E01 – Animal Removal Summary by Treatment Group
- 11 E02 – Animals Removed from Experiment
- 12 E03 – Growth Curves
- 13 E04 – Mean Body Weights and Survival Table
- 14 E05 – Clinical Observations Summary
- 15 E07 – Mean Water Consumption by Treatment Group
- 16 E08 – Water and Compound Consumption Table
- 17 E12 – Animal History
- 18 P02 – Incidence Rates of Neoplasms by Anatomic Site
- 19 P03 – Incidence Rates of Non–Neoplastic Lesions by Anatomic Site
- 20 P04 – Neoplasms by Individual Animal
- 21 P05 – Incidence Rates of Neoplasms by Anatomic Site (Systemic Lesions Abridged)
- 22 P08 – Statistical Analysis of Primary Tumors
- 23 P09 – Non–Neoplastic Lesions by Individual Animal
- 24 P10 – Statistical Analysis of Non–Neoplastic Lesions
- 25 P11 – Statistical Analysis of Survival Data
- 26 P14 – Individual Animal Pathology Data
- 27 P17 – Neoplasms by Individual Animal (Systemic Lesions Abridged)

1 P18 – Incidence Rates of Non–Neoplastic Lesions by Anatomic Site with Average Severity
2 Grades

3 P40 – Survival Curves

4 PA06 – Organ Weight Summary

5 PA48 – Summary of Tissue Concentration

6 PA48C – Tissue Concentration Curve

7 **G.8. Two-year Study in Mice – Individual Animal Data**

8 Female Individual Animal Body Weight Data

9 Female Individual Animal Clinical Observations

10 Female Individual Animal Neoplastic Pathology Data

11 Female Individual Animal Non–Neoplastic Pathology Data

12 Female Individual Animal Survival Data

13 Female Individual Animal Terminal Body Weight Data

14 Male Individual Animal Body Weight Data

15 Male Individual Animal Clinical Observations

16 Male Individual Animal Neoplastic Pathology Data

17 Male Individual Animal Non–Neoplastic Pathology Data

18 Male Individual Animal Survival Data

19 Male Individual Animal Terminal Body Weight Data

20 Individual Animal Organ Weight Data

21 Individual Animal Tissue Concentration Data

22 Individual Animal Urinalysis Data

23 **G.9. Genetic Toxicology**

24 **G.9.1. In Vivo Peripheral Blood Micronucleus Study G03038B in Mice**

25 G03038B G04 In Vivo Micronucleus Summary Data

26 G03038B Individual Animal In Vivo Micronucleus Data

27 **G.9.2. In Vivo Peripheral Blood Micronucleus Study G03038C in Rats**

28 G03038C G04 In Vivo Micronucleus Summary Data

- 1 G03038C Individual Animal In Vivo Micronucleus Data
- 2 **G.9.3. *Salmonella/E.coli* Mutagenicity Test or Ames Test Study G03038D**
- 3 G03038D G06 Ames Summary Data
- 4 **G.9.4. DNA Damage Study G03038E in Mice**
- 5 G01 – In Vivo Alkaline Comet Summary Data
- 6 Individual Animal In Vivo Alkaline Comet Data
- 7 **G.9.5. DNA Damage Study G03038F in Rats**
- 8 G01 – In Vivo Alkaline Comet Summary Data
- 9 Individual Animal In Vivo Alkaline Comet Data