








NTP Nonneoplastic Lesion Atlas

Lung – Metaplasia, Osseous






NTP Nonneoplastic Lesion Atlas

Lung – Metaplasia, Osseous

Figure Legend: **Figure 1** Lung - Metaplasia, Osseous in a F344/N rat from a chronic study. A focus of unmineralized bone is present in the lung parenchyma. **Figure 2** Lung - Metaplasia, Osseous in a F344/N rat from a chronic study (higher magnification of Figure 1). The eosinophilic material in the lung is unmineralized bone. **Figure 3** Lung - Metaplasia, Osseous in a male F344/N rat from a subchronic study. A focus of mineralized bone is present in the lung parenchyma. **Figure 4** Lung - Metaplasia, Osseous in a female B6C3F1/N mouse from a chronic study. There is a subpleural focus of mineralized bone within a focus of unmineralized bone. **Figure 5** Lung - Metaplasia, Osseous from a male Wistar Han rat in a chronic study. Though this focus of osseous metaplasia (arrow) is within an area of inflammation, it is likely not treatment related. **Figure 6** Lung - Metaplasia, Osseous in a female F344/N rat from a chronic study. A focus of mineralized bone is present in the peripheral lung.

Comment: Osseous metaplasia (Figure 1, Figure 2, Figure 3, Figure 4, Figure 5, and Figure 6) is frequently seen as a background lung lesion in NTP studies. No association with test article administration has been shown. It may occur anywhere in the lung, including the pleura (Figure 4). Osseous metaplasia is seen as foci of eosinophilic osteoid or woven bone or basophilic mineralized bone that contain interspersed lacunae. Rarely, bone is formed that contains hematopoietic elements (bone marrow). Cartilaginous metaplasia is also seen in the lung, comprising amphophilic to basophilic foci of chondroid matrix that contain interspersed lacunae. In laboratory rodents, osseous metaplasia is most common in the lungs of B6C3F1 mice. Cartilaginous and osseous metaplasia may occur as a response to injury or inflammation. These lesions occur when fibroblastic tissue is stimulated or induced to form nonneoplastic bone and/or cartilage. New bone forms with or without a cartilage template and can be bordered by fibroblastic tissue. Foci of osseous or cartilaginous metaplasia must be distinguished from metastatic osteosarcoma or chondrosarcoma, respectively.

Recommendation: Lung - Metaplasia, Osseous is generally considered a background lesion but should be diagnosed for completeness. Similarly, cartilaginous metaplasia should also be diagnosed when present. These lesions need not be graded.



NTP Nonneoplastic Lesion Atlas

Lung – Metaplasia, Osseous

References:

Dixon D, Herbert RA, Sills RC, Boorman GA. 1999. Lungs, pleura, and mediastinum. In: Pathology of the Mouse: Reference and Atlas (Maronpot RR, Boorman GA, Gaul BW, eds). Cache River Press, Vienna, IL, 293-332.

Long PH, Leininger JR, Nold JB, Lieuallen WG. 1993. Proliferative lesions of bone, cartilage, tooth, and synovium in rats. MST-2. In: Guides for Toxicologic Pathology. STP/ARP/AFIP, Washington, DC.

Authors:

Mark F. Cesta, DVM, PhD, DACVP
Staff Scientist/NTP Pathologist
NTP Pathology Group
National Toxicology Program
National Institute of Environmental Health Sciences
Research Triangle Park, NC

Darlene Dixon, DVM, PhD, DACVP
Group Leader
Molecular Pathogenesis Group
National Toxicology Program
National Institute of Environmental Health Sciences
Research Triangle Park, NC

Ronald A. Herbert, DVM, PhD
Group Leader/NTP Pathologist
Pathology Support Group
National Toxicology Program
National Institute of Environmental Health Sciences
Research Triangle Park, NC

Lauren M. Staska, DVM, PhD, DACVP
Senior Pathologist
WIL Research
Hillsborough, NC