





NTP Nonneoplastic Lesion Atlas

Nose – Hyperplasia, Goblet Cell






NTP Nonneoplastic Lesion Atlas

Nose – Hyperplasia, Goblet Cell

Figure Legend: **Figure 1** Nose, Respiratory epithelium - Hyperplasia, Goblet cell in a male F344/N rat from a chronic study. Goblet cells are increased in number in the epithelium lining the nasal septum. **Figure 2** Nose, Respiratory epithelium - Hyperplasia, Goblet cell in a male F344/N rat from a chronic study. Goblet cells are increased in number in the epithelium on one side of the nasal septum (top); the opposite side is lined by metaplastic squamous epithelium. **Figure 3** Nose, Respiratory epithelium - Hyperplasia, Goblet cell (right) and normal respiratory epithelium (left) of the nasopharyngeal duct in a male F344/N rat from a subchronic study. Image provided courtesy of Dr. R. Miller. **Figure 4** Nose, Respiratory epithelium - Hyperplasia, Goblet cell in the nasopharyngeal duct in a male F344/N rat from a subchronic study. Hematoxylin and eosin staining of goblet cell hyperplasia is on the left, and periodic acid–Schiff staining is on the right. Image provided courtesy of Dr. R. Miller. **Figure 5** Nose, Transitional epithelium - Hyperplasia, Goblet cell in a male F344/N rat from a chronic study. Increased numbers of goblet cells are present in the epithelium lining the turbinate, with coalescence of goblet cells on one side of the turbinate (arrow).

Comment: Goblet (mucous) cell hyperplasia (Figure 1, Figure 2, Figure 3, Figure 4, and Figure 5) is seen after exposure to some irritants and may be the only change noted in response to short-term exposure to a mild irritant. It is most often seen in areas of the nasal cavity that are lined by transitional and respiratory epithelium (levels I and II). The respiratory epithelium normally contains a few goblet cells; the term “metaplasia” may be more correct when the lesion is present in the transitional epithelium (which normally lacks goblet cells). However, to maintain consistency, the NTP prefers the term “hyperplasia” for this lesion, regardless of the epithelial type. The lesion is characterized by the presence of (in transitional epithelium) or increase in the number of (in respiratory epithelium) goblet cells in the epithelial lining. The goblet cells may form intraepithelial glands, or they may coalesce to form mucous cysts (Figure 5, arrow). Goblet cell hyperplasia has been seen in both short- and long-term studies. It is thought to be an adaptive response to irritant exposure and is not considered a preneoplastic change.

Recommendation: Goblet cell hyperplasia should be diagnosed whenever it is present and assigned a severity grade. The type of epithelium affected should be indicated in the diagnosis by a site modifier (e.g., respiratory epithelium or transitional epithelium). The goblet cells may be enlarged or



NTP Nonneoplastic Lesion Atlas

Nose – Hyperplasia, Goblet Cell


hypertrophied, but goblet cell hypertrophy should not be concurrently diagnosed; rather, the hypertrophy should be described in the pathology narrative. Associated lesions, such as inflammation or epithelial hyperplasia, should be diagnosed separately.

References:

- Boorman GA, Morgan KT, Uraih LC. 1990. Nose, larynx, and trachea. In: Pathology of the Fischer Rat: Reference and Atlas (Boorman GA, Eustis SL, Elwell MR, eds). Academic Press, San Diego, 315-337.
- Harkema JR, Hotchkiss JA, Henderson RF. 1989. Effects of 0.12 and 0.80 ppm ozone on rat nasal and nasopharyngeal epithelial mucosubstances: Quantitative histochemistry. *Toxicol Pathol* 17:525-535.
Abstract: <http://www.ncbi.nlm.nih.gov/pubmed/2814228>
- Harkema JR, Morgan KT, Gross EA, Catalano PJ, Griffith WC. 1994. Consequences of Prolonged Inhalation of Ozone on F344/N Rats: Collaborative Studies. Part VII: Effects on the Nasal Mucociliary Apparatus. Research Report No. 65. Health Effects Institute, Cambridge, MA.
Abstract: <http://www.ncbi.nlm.nih.gov/pubmed/7888110>
- Herbert RA, Leninger JR. 1999. Nose, larynx, and trachea. In: Pathology of the Mouse: Reference and Atlas (Maronpot RR, ed). Cache River Press, Vienna, IL, 259-292.
- Monticello TM, Morgan KT, Uraih LC. 1990. Nonneoplastic nasal lesions in rats and mice. *Environ Health Perspect* 85:249-274.
Full Text: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC1568333/>
- National Toxicology Program. 1997. NTP TR-461. Toxicology and Carcinogenesis Studies of Nitromethane (CAS No. 75-52-5) in F344/N Rats and B6C3F1 Mice (Inhalation Studies). NTP, Research Triangle Park, NC.
Abstract: <http://ntp.niehs.nih.gov/go/6066>
- National Toxicology Program. 1999. NTP TR-490. Toxicology and Carcinogenesis Studies of Glutaraldehyde (CAS No. 111-30-8) in F344/N Rats and B6C3F1 Mice (Inhalation Studies). NTP, Research Triangle Park, NC.
Abstract: <http://ntp.niehs.nih.gov/go/10170>
- Renne R, Brix A, Harkema J, Kittel B, Lewis D, March T, Nagano K, Pino M, Rittinghausen S, Rosenbruch M, Tellier P, Wohrmann T. 2009. Proliferative and nonproliferative lesions of the rat and mouse respiratory tract. *Toxicol Pathol* 37(7 suppl):5S-73S.
Abstract: <http://www.ncbi.nlm.nih.gov/pubmed/20032296>
- Wagner JG, Hotchkiss JA, Harkema JR. 2001. Effects of ozone and endotoxin coexposure on rat airway epithelium: Potentiation of toxicant-induced alterations. *Environ Health Perspect* 109:591-598.
Abstract: <http://www.ncbi.nlm.nih.gov/pubmed/11544169>



National Toxicology Program
U.S. Department of Health and Human Services



NTP Nonneoplastic Lesion Atlas

Nose – Hyperplasia, Goblet Cell

Authors:

Rodney A. Miller, DVM, PhD, DACVP
NC Pathology Group Manager
Senior Pathologist
Experimental Pathology Laboratories, Inc.
Research Triangle Park, NC

Mark F. Cesta, DVM, PhD, DACVP
Staff Scientist, NTP Pathologist
Cellular and Molecular Pathology Branch
Division of the National Toxicology Program
National Institute of Environmental Health Sciences
Research Triangle Park, NC