TDMS No. 20011 - 06 Test Type: CHRONIC P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

1-Bromopropane

CAS Number: 106-94-5

F1_M3

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

Lab: BNW

Species/Strain: MICE/B6C3F1

C Number: C20011

Route: RESPIRATORY EXPOSURE WHOLE BODY

Lock Date: 06/15/2006

Cage Range: ALL

Date Range: ALL

Reasons For Removal: ALL

Removal Date Range: ALL

Treatment Groups: Include ALL

Study Gender: Both

TDMSE Version: 2.0.0

TDMS No. 20011 - 06 Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

1-Bromopropane

CAS Number: 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

| B6C3F1 MICE MALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM |
|--|--------------------------------------|--|-------------------------------|---------------------------------------|
| Disposition Summary | | | | |
| Animals Initially in Study Early Deaths | 50 | 50 | 50 | 50 |
| Moribund Sacrifice Natural Death | 6 7 | 13 4 | 9 9 | 9 5 |
| Survivors Terminal Sacrifice Animals Examined Microscopically | 37 50 | 33 50 | 32 50 | 36 50 |
| ALIMENTARY SYSTEM | | | | |
| Gallbladder Degeneration, Hyaline Hemorrhage | (40) 2 (5%) | (40) | (34) | (38) 1 (3%) 1 (3%) |
| Hyperplasia Infiltration Cellular, Polymorphonuclear | 1 (3%) | | 1 (3%) | |
| Intestine Large, Rectum Inflammation, Suppurative Ulcer | (45) | (48) | (43) | (48) 1 (2%) 1 (2%) |
| Artery, Inflammation, Chronic Active Intestine Small, Duodenum Inflammation, Acute | 1 (2%) (44) | (48) 1 (2%) | (43) | (46) |
| Intestine Small, Jejunum Liver | (44) (50) | (48) (50) | (43) (50) | (46) (50) |
| Angiectasis Basophilic Focus Clear Cell Focus Eosinophilic Focus | 1 (2%) 9 (18%) 20 (40%) | 3 (6%) 2 (4%) 14 (28%) 16 (32%) | 1 (2%) 16 (32%) 8 (16%) | 3 (6%) 12 (24%) 9 (18%) |
| Fatty Change, Focal Inflammation, Chronic Mineralization | 1 (2%) | 3 (6%) | 1 (2%) | 5 (10%) 1 (2%) 1 (2%) |
| Mixed Cell Focus Necrosis Tension Lipidosis Vacuolization Cytoplasmic | 2 (4%) 5 (10%) 4 (8%) | 7 (14%) 4 (8%) 3 (6%) | 1 (2%) 3 (6%) 6 (12%) | 1 (2%) 9 (18%) 2 (4%) 1 (2%) |
| Bile Duct, Hyperplasia Mesentery Artery, Inflammation | (9) | (11) | (6) | 1 (2%) (7) 1 (14%) |
| Artery, Mineralization Fat, Necrosis Pancreas Atrophy | 1 (11%) 7 (78%) (48) 1 (2%) | 11 (100%) (50) | 5 (83%) (48) | 5 (71%) (50) 1 (2%) |

a - Number of animals examined microscopically at site and number of animals with lesion

TDMS No. 20011 - 06 Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

1-Bromopropane

CAS Number: 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

| B6C3F1 MICE MALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM | |
|--------------------------------------|-----------------|-----------------|----------------|----------------|--|
| | | | | | |
| Basophilic Focus | 1 (2%) | | 1 (2%) | | |
| Cyst | 1 (2%) | | | | |
| Inflammation, Chronic Active | 1 (2%) | (50) | (40) | (50) | |
| Salivary Glands | (49) | (50) | (49) | (50) | |
| Atrophy Stomach, Forestomach | 1 (2%) (47) | (50) | 1 (2%) (48) | (50) | |
| Hyperplasia, Squamous | (47) 5 (11%) | (50) 7 (14%) | 5 (10%) | (50) 3 (6%) | |
| Inflammation | 2 (4%) | 2 (4%) | 1 (2%) | 3 (0 %) | |
| Ulcer | 2 (4%) | 4 (8%) | 3 (6%) | 2 (4%) | |
| Artery, Inflammation, Chronic Active | 1 (2%) | 4 (070) | 3 (070) | 2 (470) | |
| Stomach, Glandular | (47) | (49) | (47) | (49) | |
| Mineralization | 2 (4%) | (10) | (, | (10) | |
| Artery, Inflammation, Chronic Active | 1 (2%) | | | | |
| Tooth | (8) | (8) | (9) | (11) | |
| Dysplasia | 7 (88%) | 8 (100%) | 9 (100%) | 11 (100%) | |
| Malformation | 1 (13%) | , , | , , | , , | |
| ARDIOVASCULAR SYSTEM | | | | | |
| Blood Vessel | (3) | (1) | (0) | (0) | |
| Inflammation, Chronic Active | 1 (33%) | 4 (4000() | | | |
| Thrombosis | 0 (070/) | 1 (100%) | | | |
| Aorta, Mineralization Heart | 2 (67%) (50) | (50) | (50) | (50) | |
| Cardiomyopathy | 12 (24%) | 10 (20%) | 14 (28%) | 9 (18%) | |
| Hemorrhage | 12 (24 %) | 10 (20%) | 14 (20 %) | 1 (2%) | |
| Inflammation, Acute | 1 (2%) | | | 1 (270) | |
| Mineralization | 2 (4%) | | | | |
| Thrombosis | 1 (2%) | 2 (4%) | 1 (2%) | | |
| Artery, Inflammation, Chronic Active | 1 (2%) | 1 (2%) | , | | |
| Epicardium, Inflammation, Chronic | | 1 (2%) | | | |
| IDOCRINE SYSTEM | | | | | |
| Adrenal Cortex | (49) | (50) | (49) | (50) | |
| Hyperplasia | 16 (33%) | 18 (36%) | 14 (29%) | 12 (24%) | |
| Hypertrophy | 23 (47%) | 19 (38%) | 22 (45%) | 28 (56%) | |
| Subcapsular, Hyperplasia | (==) | 1 (2%) | //5 | (50) | |
| Adrenal Medulla | (50) | (49) | (49) | (50) | |
| Hyperplasia | 1 (2%) | 2 (4%) | 1 (2%) | 2 (4%) | |
| Islets, Pancreatic | (48) | (50) | (48) | (50) | |
| Hyperplasia Pituitary Gland | 1 (2%) (46) | (48) | (46) | (48) | |
| Cyst | (40) | (40) | 1 (2%) | 1 (2%) | |
| Cyst | | | 1 (2/0) | 1 (2/0) | |

a - Number of animals examined microscopically at site and number of animals with lesion

TDMS No. 20011 - 06 Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

1-Bromopropane CAS Number: 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

| B6C3F1 MICE MALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM | |
|---|-----------------|--------------|----------------|--------------|--|
| Pars Distalis, Hyperplasia | 2 (4%) | 2 (4%) | 1 (2%) | 1 (2%) | |
| Pars Intermedia, Hyperplasia | (40) | (50) | 1 (2%) | (50) | |
| Thyroid Gland Follicular Cell, Hyperplasia | (49) 1 (2%) | (50) | (48) | (50) | |
| GENERAL BODY SYSTEM | | | | | |
| Peritoneum | (0) | (0) | (1) | (0) | |
| GENITAL SYSTEM | | | | | |
| Coagulating Gland Inflammation, Suppurative | (1) 1 (100%) | (0) | (0) | (0) | |
| Epididymis | (50) | (50) | (50) | (50) | |
| Granuloma Sperm | 1 (2%) | 1 (2%) | 1 (2%) | | |
| Necrosis | (40) | (40) | (50) | 2 (4%) | |
| Preputial Gland Inflammation, Suppurative | (49) | (49) | (50) 1 (2%) | (50) | |
| Inflammation, Granulomatous | | | 1 (2%) | 1 (2%) | |
| Prostate | (49) | (50) | (48) | (50) | |
| Inflammation, Suppurative | 2 (4%) | (| 2 (4%) | 2 (4%) | |
| Inflammation, Chronic Active | | | 1 (2%) | 1 (2%) | |
| Artery, Inflammation, Chronic Active Seminal Vesicle | 1 (2%) | (50) | (40) | (50) | |
| Amyloid Deposition | (48) 1 (2%) | (50) | (49) | (50) | |
| Inflammation, Suppurative | 1 (2%) | | | | |
| Inflammation, Chronic Active | | | 1 (2%) | | |
| Testes | (50) | (50) | (50) | (50) | |
| Atrophy | 1 (2%) | 1 (2%) | 1 (2%) | 3 (6%) | |
| HEMATOPOIETIC SYSTEM | | | | | |
| Bone Marrow | (48) | (48) | (48) | (50) | |
| Lymph Node | (1) | (1) | (3) | (0) | |
| Lymph Node, Bronchial | (23) | (30) | (32) | (31) | |
| Infiltration Cellular, Mixed Cell Lymph Node, Mandibular | (27) | 1 (3%) | (24) | (26) | |
| Lymph Node, Mediastinal | (27) (35) | (28) (32) | (21) (35) | (26) (34) | |
| Infiltration Cellular, Mixed Cell | (30) | 1 (3%) | (00) | (04) | |
| Lymph Node, Mesenteric | (48) | (46) | (44) | (48) | |
| Angiectasis | 1 (2%) | 4 | 4> | | |
| Spleen | (48) | (50) | (49) | (50) | |
| Angiectasis | | 1 (2%) | | | |

a - Number of animals examined microscopically at site and number of animals with lesion

TDMS No. 20011 - 06 Test Type: CHRONIC

Species/Strain: MICE/B6C3F1

Route: RESPIRATORY EXPOSURE WHOLE BODY

1-Bromopropane **CAS Number:** 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

| B6C3F1 MICE MALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM | |
|--|---|---|---|--|--|
| Atrophy Congestion Hematopoietic Cell Proliferation Hyperplasia, Lymphoid Infarct | 7 (15%) 4 (8%) 20 (42%) 5 (10%) 1 (2%) | 8 (16%) 4 (8%) 19 (38%) 2 (4%) | 2 (4%) 1 (2%) 14 (29%) 5 (10%) | 3 (6%) 3 (6%) 16 (32%) 2 (4%) | |
| Necrosis Necrosis, Lymphoid Thymus Atrophy Cyst Necrosis, Lymphoid Medulla, Hyperplasia, Lymphoid | 1 (2%) 1 (2%) (38) 13 (34%) 5 (13%) 1 (3%) 16 (42%) | (35) 12 (34%) 1 (3%) 9 (26%) | 1 (2%) (37) 19 (51%) 7 (19%) 1 (3%) 14 (38%) | 2 (4%) (32) 12 (38%) 4 (13%) 3 (9%) 6 (19%) | |
| INTEGUMENTARY SYSTEM | | | | | |
| Skin Cyst Epithelial Inclusion Inflammation, Chronic Active Ulcer | (50) 5 (10%) 1 (2%) | (50) 5 (10%) | (50) 3 (6%) | (50) 1 (2%) 8 (16%) | |
| MUSCULOSKELETAL SYSTEM | | | | | |
| Bone Hyperostosis Necrosis Skeletal Muscle | (50) (0) | (49) 2 (4%) (2) | (50) (1) | (50) 1 (2%) 1 (2%) (2) | |
| | (0) | (2) | (1) | (2) | |
| NERVOUS SYSTEM | | | | | |
| Brain | (50) | (50) | (49) | (50) | |
| RESPIRATORY SYSTEM | | | | | |
| Larynx Vacuolization Cytoplasmic Lung Congestion, Chronic Hemorrhage Inflammation, Chronic Active Mineralization Pigmentation Thrombosis | (48) (50) 1 (2%) 2 (4%) 1 (2%) 1 (2%) | (50) 5 (10%) (50) 2 (4%) 1 (2%) | (48) 10 (21%) (49) 1 (2%) 2 (4%) | (50) 11 (22%) (49) 1 (2%) | |

a - Number of animals examined microscopically at site and number of animals with lesion

TDMS No. 20011 - 06 Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

1-Bromopropane

CAS Number: 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

| BF1 MICE MALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM | |
|---|-----------|-----------|------------------|-----------|--|
| | 0 (40) | 0 (40() | 4 (00() | E (400() | |
| eolar Epithelium, Hyperplasia | 2 (4%) | 2 (4%) | 1 (2%) | 5 (10%) | |
| eolus, Infiltration Cellular, Histiocyte | | 3 (6%) | 1 (2%) | 1 (2%) | |
| nchiole, Hyperplasia | | 3 (6%) | | 1 (2%) | |
| nchiole, Necrosis | 4 (00/) | 44 (000() | 20 (700() | 1 (2%) | |
| nchiole, Regeneration | 1 (2%) | 44 (88%) | 38 (78%) | 47 (96%) | |
| nchiole, Vacuolization Cytoplasmic | (50) | 18 (36%) | 19 (39%) | 17 (35%) | |
| eign Body | (50) | (50) | (50) | (50) | |
| eign Body ammation, Suppurative | 2 (40/) | 1 (2%) | 2 (40/) | 3 (60() | |
| yp, Inflammatory | 2 (4%) | 3 (6%) | 2 (4%) | 3 (6%) | |
| yp, mnammatory ınds, Hyperplasia | | 1 (2%) | | 3 (6%) | |
| actory Epithelium, Atrophy | 2 (4%) | 4 (8%) | 7 (14%) | 4 (8%) | |
| actory Epithelium, Attoprty actory Epithelium, Metaplasia, Respiratory | 2 (4%) | 7 (14%) | 6 (12%) | 3 (6%) | |
| spiratory Epithelium, Hyperplasia | 16 (32%) | 29 (58%) | 23 (46%) | 26 (52%) | |
| spiratory Epithelium, Vacuolization | 10 (32 %) | 12 (24%) | 19 (38%) | 20 (40%) | |
| cytoplasmic | | 12 (2470) | 19 (30 %) | 20 (4078) | |
| a | (0) | (0) | (1) | (1) | |
| nea | (49) | (50) | (47) | (50) | |
| cuolization Cytoplasmic | (10) | 15 (30%) | 24 (51%) | 24 (48%) | |
| AL SENSES SYSTEM | | | | | |
| | (45) | (47) | (44) | (47) | |
| aract | 1 (2%) | 1 (2%) | 1 (2%) | | |
| generation | 1 (2%) | 1 (2%) | | 1 (2%) | |
| nea, Inflammation, Acute | | | 1 (2%) | | |
| nea, Inflammation, Chronic Active | 2 (4%) | | 1 (2%) | | |
| ina, Atrophy | 1 (2%) | (40) | 1 (2%) | (40) | |
| erian Gland | (46) | (49) | (49) | (48) | |
| perplasia | 3 (7%) | 4 (8%) | 1 (2%) | 4 (8%) | |
| RY SYSTEM | | | | | |
| ey | (49) | (50) | (50) | (49) | |
| yloid Deposition | , , | , , | , , | 1 (2%) | |
| st | | 1 (2%) | | | |
| dronephrosis | 1 (2%) | | | 1 (2%) | |
| arct | 1 (2%) | 2 (4%) | 3 (6%) | 1 (2%) | |
| ammation, Suppurative | 1 (2%) | | 2 (4%) | 2 (4%) | |
| taplasia, Osseous | 2 (4%) | 1 (2%) | | | |
| neralization | 2 (4%) | | | | |
| phropathy | 44 (90%) | 46 (92%) | 47 (94%) | 41 (84%) | |
| pilla, Necrosis | - 4 | 1 (2%) | 1 (2%) 2 (4%) | | |
| nal Tubule, Hyperplasia | 2 (4%) | 4 (00/) | | | |

a - Number of animals examined microscopically at site and number of animals with lesion

TDMS No. 20011 - 06

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Test Type: CHRONIC

1-Bromopropane Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

CAS Number: 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

Lab: BNW

| B6C3F1 MICE MALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM |
|--|------------------|----------|---------|---------------|
| | | | | . (22() |
| Renal Tubule, Necrosis Urethra | (0) | (0) | (0) | 1 (2%) (1) |
| Bulbourethral Gland, Angiectasis | • • | | | 1 (100%) |
| Urinary Bladder Inflammation, Chronic Active | (47) | (50) | (47) | (48) |
| Mineralization | 1 (2%) 1 (2%) | | 1 (2%) | 1 (2%) |

*** END OF MALE ***

TDMS No. 20011 - 06
Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

1-Bromopropane **CAS Number:** 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

| B6C3F1 MICE FEMALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM |
|---|------------------|------------------|------------------|---------|
| | | | | |
| Disposition Summary | | | | |
| Animals Initially in Study | 50 | 50 | 50 | 50 |
| Early Deaths | 40 | _ | • | |
| Moribund Sacrifice Natural Death | 10 4 | 7 3 | 8 5 | 8 |
| Survivors | 7 | 9 | 3 | |
| Moribund Sacrifice | 1 | | | 1 |
| Terminal Sacrifice | 35 | 40 | 37 | 41 |
| Animals Examined Microscopically | 50 | 50 | 50 | 50 |
| | | | | |
| ALIMENTARY SYSTEM | | | | |
| Esophagus | (50) | (50) | (50) | (50) |
| Gallbladder | (43) | (36) | (36) | (44) |
| Inflammation, Chronic Active | | 1 (3%) | | |
| Intestine Large, Cecum | (48) | (49) | (46) | (50) |
| Hemorrhage Inflammation, Suppurative | 1 (2%) | 1 (2%) | | |
| Intestine Large, Colon | (48) | (49) | (46) | (50) |
| Necrosis | 1 (2%) | (30) | (-10) | (00) |
| Intestine Large, Rectum | (47) | (49) | (46) | (50) |
| Artery, Inflammation, Chronic Active | | | | 1 (2%) |
| Intestine Small, Duodenum | (46) | (49) | (46) | (50) |
| Intestine Small, Ileum | (48) | (49) | (46) | (50) |
| Epithelium, Hyperplasia Intestine Small, Jejunum | (46) | (49) | 1 (2%) (46) | (50) |
| Epithelium, Hyperplasia | (40) | (43) | 1 (2%) | (30) |
| Liver | (50) | (50) | (50) | (50) |
| Angiectasis | | 1 (2%) | 1 (2%) | 2 (4%) |
| Basophilic Focus | 2 (4%) | 3 (6%) | 2 (4%) | 3 (6%) |
| Clear Cell Focus | 2 (4%) | 7 (4 (2)) | 1 (2%) | 2 (4%) |
| Eosinophilic Focus | 6 (12%) | 7 (14%) | 7 (14%) | 5 (10%) |
| Fatty Change Fatty Change, Focal | 1 (2%) 2 (4%) | 1 (2%) | | 2 (4%) |
| Hematopoietic Cell Proliferation | 2 (470) | | 1 (2%) | |
| Infarct | | 1 (2%) | . (= /0) | |
| Inflammation, Acute | 1 (2%) | | | |
| Inflammation, Chronic | | 1 (2%) | 1 (2%) | |
| Malformation, Lobular | | | 1 (2%) | |
| Mineralization | 2 (49/) | 2 (60/) | 1 (2%) | |
| Mixed Cell Focus Necrosis | 2 (4%) 1 (2%) | 3 (6%) 1 (2%) | 3 (6%) 2 (4%) | 1 (2%) |
| 140010313 | 1 (2/0) | 1 (2/0) | 2 (470) | 1 (2/0) |

a - Number of animals examined microscopically at site and number of animals with lesion

TDMS No. 20011 - 06 Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

1-Bromopropane **CAS Number:** 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

| B6C3F1 MICE FEMALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM | |
|---|---------|----------|---------|----------|--|
| | | | | | |
| Tension Lipidosis | 3 (6%) | 10 (20%) | 4 (8%) | 1 (2%) | |
| Thrombosis | | | 1 (2%) | | |
| Vacuolization Cytoplasmic | | 1 (2%) | | | |
| Bile Duct, Hyperplasia | | 1 (2%) | | | |
| Hepatocyte, Mitotic Alteration | 1 (2%) | 4-1 | | (1.5) | |
| Mesentery | (11) | (9) | (11) | (13) | |
| Artery, Inflammation | 1 (9%) | | . () | | |
| Fat, Hemorrhage | 2 (222) | 1 (11%) | 1 (9%) | 10 (000) | |
| Fat, Necrosis | 9 (82%) | 9 (100%) | 9 (82%) | 12 (92%) | |
| Pancreas | (50) | (49) | (50) | (50) | |
| Atrophy | 7 (14%) | 4 (8%) | 5 (10%) | 2 (4%) | |
| Basophilic Focus | 1 (2%) | 2 (4%) | | 4 (00() | |
| Inflammation, Chronic Active | (50) | 1 (2%) | (50) | 1 (2%) | |
| Salivary Glands | (50) | (50) | (50) | (50) | |
| Stomach, Forestomach | (50) | (50) | (49) | (50) | |
| Hyperplasia, Squamous | 3 (6%) | 2 (4%) | 2 (4%) | 1 (2%) | |
| Inflammation | 1 (2%) | - / | | - () | |
| Ulcer | 2 (4%) | 2 (4%) | | 2 (4%) | |
| Artery, Inflammation, Chronic Active | 4> | | () | 1 (2%) | |
| Stomach, Glandular | (50) | (49) | (48) | (50) | |
| Infiltration Cellular, Mixed Cell | 4 (22() | 1 (2%) | | | |
| Infiltration Cellular, Mononuclear Cell | 1 (2%) | | . (224) | . (00.1) | |
| Mineralization | | . (22() | 1 (2%) | 1 (2%) | |
| Necrosis | | 1 (2%) | 1 (2%) | . (00.1) | |
| Artery, Inflammation, Chronic Active | (2) | (2) | 440 | 1 (2%) | |
| Tongue | (0) | (0) | (1) | (0) | |
| Tooth | (0) | (1) | (0) | (0) | |
| Dysplasia | | 1 (100%) | | | |
| ARDIOVASCULAR SYSTEM | | | | | |
| Heart | (50) | (50) | (50) | (50) | |
| Cardiomyopathy | 6 (12%) | 12 (24%) | 8 (16%) | 8 (16%) | |
| Inflammation, Suppurative | | | 1 (2%) | | |
| Mineralization | 3 (6%) | | 2 (4%) | | |
| Necrosis | | 1 (2%) | _ , . | | |
| Thrombosis | | | 2 (4%) | | |
| Artery, Inflammation, Chronic Active | 1 (2%) | 1 (2%) | | | |
| NDOCRINE SYSTEM | | | | | |
| Adrenal Cortex | (50) | (50) | (50) | (50) | |
| Hyperplasia | 3 (6%) | 10 (20%) | 4 (8%) | 6 (12%) | |
| Hypertrophy | 4 (8%) | 3 (6%) | 1 (2%) | 9 (18%) | |

a - Number of animals examined microscopically at site and number of animals with lesion

TDMS No. 20011 - 06 Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

1-Bromopropane

CAS Number: 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

| B6C3F1 MICE FEMALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM | |
|--|----------|--------------------|------------------|-----------|--|
| | | | | | |
| Inflammation, Suppurative | | | | 1 (2%) | |
| Subcapsular, Hyperplasia | | 1 (2%) | 1 (2%) | | |
| Adrenal Medulla | (50) | (50) | (49) | (50) | |
| Hyperplasia | | 5 (10%) | 1 (2%) | 4 (8%) | |
| Islets, Pancreatic | (50) | (49) | (49) | (49) | |
| Hyperplasia | (10) | 2 (4%) | 1 (2%) | (10) | |
| Pituitary Gland | (46) | (49) | (47) | (48) | |
| Pars Distalis, Angiectasis | 1 (2%) | 1 (2%) | 2 (4%) | 15 (240/) | |
| Pars Distalis, Hyperplasia Pars Intermedia, Hyperplasia | 15 (33%) | 14 (29%) 1 (2%) | 11 (23%) | 15 (31%) | |
| Pars Intermedia, Hypertrophy | 1 (2%) | 1 (2%) | 2 (4%) | | |
| Thyroid Gland | (49) | (50) | (46) | (50) | |
| Follicular Cell, Hyperplasia | 1 (2%) | 1 (2%) | 1 (2%) | 1 (2%) | |
| Tomodia Con, Hypotpiaola | . (278) | . (270) | . (278) | (270) | |
| SENERAL BODY SYSTEM | | | | | |
| Peritoneum | (0) | (0) | (0) | (1) | |
| ENITAL SYSTEM | | | | | |
| Ovary | (50) | (50) | (50) | (49) | |
| Angiectasis | 1 (2%) | 2 (4%) | | 2 (4%) | |
| Congestion, Chronic | | 1 (2%) | | | |
| Cyst | 12 (24%) | 8 (16%) | 12 (24%) | 2 (4%) | |
| Hyperplasia, Tubular | | | | 1 (2%) | |
| Thrombosis | 2 (4%) | 2 (4%) | (==) | 1 (2%) | |
| Uterus | (50) | (50) | (50) | (50) | |
| Angiectasis | 2 (4%) | 3 (6%) | 6 (12%) | 3 (6%) | |
| Fibrosis Inflammation, Suppurative | | | 1 (2%) 1 (2%) | | |
| Inflammation, Suppurative Inflammation, Acute | 1 (2%) | | 1 (2%) | | |
| Thrombosis | 1 (2%) | 1 (2%) | 2 (4%) | 1 (2%) | |
| Endometrium, Hyperplasia, Cystic | 39 (78%) | 43 (86%) | 43 (86%) | 41 (82%) | |
| | (, | - () | | () | |
| EMATOPOIETIC SYSTEM | | | | | |
| Bone Marrow | (50) | (50) | (49) | (50) | |
| Angiectasis | 2 (4%) | . , | . , | | |
| Atrophy | ` ' | | | 1 (2%) | |
| Lymph Node | (6) | (5) | (3) | (8) | |
| Iliac, Angiectasis | 2 (33%) | 1 (20%) | | | |
| Iliac, Hyperplasia, Lymphoid | 1 (17%) | | | | |
| Lumbar, Angiectasis | | 1 (20%) | 1 (33%) | 1 (13%) | |
| | | | | | |

a - Number of animals examined microscopically at site and number of animals with lesion

TDMS No. 20011 - 06 Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

1-Bromopropane CAS Number: 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

| Renal, Angiectasis Renal, Hyperplasia, Lymphoid Lymph Node, Bronchial Lymph Node, Mandibular Hyperplasia, Lymphoid Lymph Node, Mediastinal Hyperplasia, Lymphoid | 1 (17%) (37) | 1 (20%) | | | |
|--|-----------------|--------------|----------|----------|--|
| Renal, Hyperplasia, Lymphoid Lymph Node, Bronchial Lymph Node, Mandibular Hyperplasia, Lymphoid Lymph Node, Mediastinal Hyperplasia, Lymphoid | (37) | 1 (20%) | | | |
| Lymph Node, Bronchial Lymph Node, Mandibular Hyperplasia, Lymphoid Lymph Node, Mediastinal Hyperplasia, Lymphoid | (37) | 1 (20%) | | 1 (13%) | |
| Lymph Node, Mandibular Hyperplasia, Lymphoid Lymph Node, Mediastinal Hyperplasia, Lymphoid | | (00) | (22) | (0.1) | |
| Hyperplasia, Lymphoid Lymph Node, Mediastinal Hyperplasia, Lymphoid | | (32) | (39) | (34) | |
| Lymph Node, Mediastinal Hyperplasia, Lymphoid | (45) | (30) | (30) | (37) | |
| Hyperplasia, Lymphoid | (45) | 1 (3%) | (44) | (07) | |
| Hyperpiasia, Lympnoid | (45) | (39) | (41) | (37) | |
| | 1 (2%) | (47) | (47) | (40) | |
| Lymph Node, Mesenteric | (48) | (47) | (47) | (49) | |
| Angiectasis | 2 (4%) | 2 (4%) | . (55.) | 1 (2%) | |
| Hyperplasia, Lymphoid | (==) | (40) | 1 (2%) | (=0) | |
| Spleen | (50) | (49) | (50) | (50) | |
| Atrophy | 4 (8%) | 1 (2%) | 2 (4%) | 5 (10%) | |
| Congestion | 10 (000) | 4.0 (0.00) | 1 (2%) | 2 (4%) | |
| Hematopoietic Cell Proliferation | 13 (26%) | 10 (20%) | 13 (26%) | 16 (32%) | |
| Hemorrhage | | | | 1 (2%) | |
| Hyperplasia, Histiocytic | 2 (4%) | _ ,,,,,, | | - 4 | |
| Hyperplasia, Lymphoid | 7 (14%) | 7 (14%) | 5 (10%) | 3 (6%) | |
| Hyperplasia, Plasma Cell | | | | 1 (2%) | |
| Necrosis, Lymphoid | | 1 (2%) | 1 (2%) | | |
| Pigmentation, Hemosiderin | 1 (2%) | 2 (4%) | | | |
| Thymus | (49) | (45) | (45) | (45) | |
| Atrophy | 14 (29%) | 13 (29%) | 17 (38%) | 16 (36%) | |
| Cyst | 5 (10%) | 7 (16%) | 6 (13%) | 9 (20%) | |
| Necrosis, Lymphoid | | 2 (4%) | 4 (9%) | | |
| Medulla, Hyperplasia, Lymphoid | 25 (51%) | 29 (64%) | 30 (67%) | 26 (58%) | |
| FEGUMENTARY SYSTEM | | | | | |
| Mammary Gland | (50) | (50) | (50) | (50) | |
| Hyperplasia | 1 (2%) | 1 (2%) | 1 (2%) | (33) | |
| Skin | (50) | (50) | (50) | (50) | |
| Cyst | 1 (2%) | () | () | () | |
| Inflammation, Chronic Active | () | 4 (8%) | 2 (4%) | 1 (2%) | |
| JSCULOSKELETAL SYSTEM | | | | | |
| Done | (50) | (50) | (FO) | (50) | |
| Bone | (50) | (50) | (50) | (50) | |
| Cyst Skeletal Muscle | (0) | 1 (2%) | (0) | (1) | |
| Skeletal iviuscie | (0) | (0) | (0) | (1) | |
| RVOUS SYSTEM | | | | | |
| 5 | | /= -> | /=-> | (=0) | |
| Brain | (50) | (50) | (50) | (50) | |

a - Number of animals examined microscopically at site and number of animals with lesion

TDMS No. 20011 - 06 Test Type: CHRONIC

Route: RESPIRATORY EXPOSURE WHOLE BODY

Species/Strain: MICE/B6C3F1

1-Bromopropane

CAS Number: 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

| B6C3F1 MICE FEMALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM | |
|---|----------------|-----------------|----------------|----------------------|--|
| Necrosis | | 1 (2%) | 1 (2%) | | |
| Artery, Inflammation | 1 (2%) | . (= /0) | | | |
| Peripheral Nerve | (1) | (1) | (1) | (1) | |
| Degeneration Spinal Cord | (4) | 1 (100%) | (4) | (4) | |
| Inflammation, Chronic Active | (1) | (1) 1 (100%) | (1) | (1) | |
| ESPIRATORY SYSTEM | | | | | |
| Larynx | (50) | (50) | (50) | (50) | |
| Vacuolization Cytoplasmic | | 3 (6%) | 2 (4%) | 2 (4%) | |
| Artery, Inflammation, Chronic Active | (50) | (50) | (50) | 1 (2%) | |
| Lung Inflammation, Chronic Active | (50) | (50) 1 (2%) | (50) | (50) | |
| Alveolar Epithelium, Hyperplasia | 1 (2%) | 1 (2%) | | 3 (6%) | |
| Alveolus, Infiltration Cellular, Histiocyte | 1 (270) | 1 (270) | 4 (8%) | 1 (2%) | |
| Bronchiole, Hyperplasia | 1 (2%) | | (===) | 1 (2%) | |
| Bronchiole, Necrosis | , , | | 1 (2%) | | |
| Bronchiole, Regeneration | | 45 (90%) | 43 (86%) | 49 (98%) | |
| Bronchiole, Vacuolization Cytoplasmic | (50) | 3 (6%) | 4 (8%) | 3 (6%) | |
| Nose Inflammation, Suppurative | (50) | (50) | (50) | (50) 1 (2%) | |
| Olfactory Epithelium, Atrophy | | | | 6 (12%) | |
| Olfactory Epithelium, Metaplasia, Respiratory | | 4 (8%) | 5 (10%) | 14 (28%) | |
| Respiratory Epithelium, Hyperplasia | 11 (22%) | 25 (50%) | 28 (56%) | 27 (54%) | |
| Respiratory Epithelium, Vacuolization | , | 3 (6%) | 5 (Ì0%) | 8 (Ì6%) [°] | |
| Cytoplasmic | | | | | |
| Pleura | (0) | (0) | (1) | (0) | |
| Trachea | (50) | (49) | (50) | (50) | |
| Vacuolization Cytoplasmic | | 8 (16%) | 7 (14%) | 4 (8%) | |
| PECIAL SENSES SYSTEM | | | | | |
| Eye | (47) | (49) | (45) | (50) | |
| Cataract | 3 (6%) | 1 (2%) | | 2 (4%) | |
| Cornea, Hyperplasia, Squamous | 1 (2%) | | 0 (404) | 4 (00) | |
| Cornea, Inflammation, Chronic Active Cornea, Mineralization | 1 (2%) | | 2 (4%) | 1 (2%) | |
| Cornea, Mineralization Harderian Gland | 1 (2%) (49) | (49) | 2 (4%) (48) | 1 (2%) (50) | |
| Hyperplasia | 3 (6%) | 1 (2%) | 2 (4%) | 6 (12%) | |
| Inflammation, Chronic Active | 3 (370) | . (= /0) | - (170) | 1 (2%) | |
| Zymbal's Gland | (0) | (1) | (0) | (0) | |
| • | . , | . , | . , | • • | |

a - Number of animals examined microscopically at site and number of animals with lesion

TDMS No. 20011 - 06

P03: INCIDENCE RATES OF NON-NEOPLASTIC LESIONS BY ANATOMIC SITE(a)

Test Type: CHRONIC
Route: RESPIRATORY EXPOSURE WHOLE BODY

1-Bromopropane

Species/Strain: MICE/B6C3F1

CAS Number: 106-94-5

Date Report Requested: 10/01/2008 Time Report Requested: 09:42:38 First Dose M/F: 07/21/03 / 07/21/03

Lab: BNW

| B6C3F1 MICE FEMALE | CONTROL | 62.5 PPM | 125 PPM | 250 PPM |
|---|--------------------|--|--------------------------|--------------------|
| URINARY SYSTEM | | | | |
| Kidney Amyloid Deposition Cyst Inflammation, Suppurative | (50) | (49) | (49) 1 (2%) 1 (2%) | (50) 1 (2%) |
| Inflammation, Chronic Active Metaplasia, Osseous Nephropathy Renal Tubule, Hyperplasia Transitional Epithelium, Hyperplasia | 3 (6%) 33 (66%) | 1 (2%) 4 (8%) 37 (76%) 1 (2%) | 40 (82%) 1 (2%) | 2 (4%) 39 (78%) |
| Urinary Bladder Inflammation, Chronic Active Artery, Inflammation, Chronic Active | (50) | (49) | (48) 1 (2%) | (50) 2 (4%) |

*** END OF REPORT ***

a - Number of animals examined microscopically at site and number of animals with lesion