



National Toxicology Program

U.S. Department of Health and Human Services

Revised Draft Report on Carcinogens Monograph on Antimony Trioxide: Appendices

August 15, 2018

Office of the Report on Carcinogens
Division of the National Toxicology Program
National Institute of Environmental Health Sciences
U.S. Department of Health and Human Services

This revised Report on Carcinogens monograph has not been formally distributed by the National Toxicology Program. It does not represent and should not be construed to represent any final NTP determination or policy.

Table of Contents

Appendix A: Literature Search Strategy	A-1
A.1 General approach	A-1
A.2 Search strategies.....	A-3
A.3 Exclusion of treatment for leishmaniasis from human cancer searches	A-3
A.4 Updating the literature search	A-3
A.5 Review of citations using web-based systematic review software	A-4
Appendix B: ADME Tables.....	B-1
Appendix C: Human Studies Tables.....	C-1
Appendix D: Animal Study Quality Tables.....	D-1
Appendix E: Mechanistic and Other Relevant Information	E-1
E.1 Tox21/ToxCast high-throughput screening	E-1
E.2 Effects of antioxidants and inhibitors of oxidative stress related enzymes on cells exposed to compounds containing trivalent antimony	E-2
E.3 Genotoxicity tables	E-4
E.4 Studies related to cell proliferation.....	E-20
E.5 Transcriptomic of antimony(III) potassium tartrate trihydrate in HepG2 cells	E-21
E.6: Immune effects from compounds containing pentavalent antimony.....	E-28
Appendix F: Other Relevant Information.....	F-1
F.1 Studies of antimony(III) potassium tartrate carcinogenicity in experimental animals	F-1
References.....	1

List of Tables

Table A-1. Major topics searched.....	A-1
Table B-1. Antimony(III) trioxide levels ^a (µg/g) in red blood cells during a 1-year chronic inhalation exposure (after 6 months and 12 months of exposure) and a 1-year observation period (6 months and 12 months after exposure) in Fischer 344 male and female rats.....	B-1
Table B-2. Blood antimony concentrations (µg/g blood) in female rats and mice exposed to antimony trioxide (N = 5 except where indicated).....	B-1
Table B-3. Tissue distribution of antimony (µg antimony/g tissue) in rats after oral exposure to antimony(III) trioxide by gavage or in the diet.....	B-2
Table C-1. Evaluation of selection bias in human cancer studies.....	C-1
Table C-2. Evaluation of exposure assessment methods in human cancer studies.	C-1
Table C-3. Evaluation of outcome assessment in human cancer studies.....	C-2
Table C-4. Evaluation of study sensitivity in human cancer studies.	C-2
Table C-5. Evaluation of potential for confounding bias for human cancer studies.	C-3
Table C-6. Evaluation of analysis and selective reporting for human cancer studies.	C-3
Table D-1. Schroeder <i>et al.</i> (1970) study of male rats exposed to antimony potassium tartrate in the drinking water.....	D-1

Table D-2. Schroeder <i>et al.</i> (1970) study of female rats exposed to antimony potassium tartrate in the drinking water	D-2
Table D-3. Kanisawa and Schroeder (1969) and Schroeder <i>et al.</i> (1968) study of male and female (combined) mice exposed to antimony potassium tartrate in drinking water for the lifespan of the animals	D-3
Table D-4. NTP (2017b) study of male rats exposed to antimony trioxide by inhalation for 105 weeks	D-4
Table D-5. NTP (2017b) study of female rats exposed to antimony trioxide by inhalation for 105 weeks	D-5
Table D-6. NTP (2017b) study of male mice exposed to antimony trioxide by inhalation for 105 weeks	D-6
Table D-7. NTP (2017b) study of female mice exposed to antimony trioxide by inhalation for 105 weeks	D-7
Table D-8. Groth <i>et al.</i> (1986) study of male rats exposed to antimony trioxide by inhalation for 53 weeks followed by post-exposure observation for 71 to 73 weeks	D-8
Table D-9. Groth <i>et al.</i> (1986) study of female rats exposed to antimony trioxide by inhalation for 53 weeks followed by post-exposure observation for 71 to 73 weeks	D-9
Table D-10. Newton <i>et al.</i> (1994) study of male rats exposed to antimony trioxide by inhalation for 12 months followed by post-exposure observation for 24 months	D-10
Table D-11. Newton <i>et al.</i> (1994) study of female rats exposed to antimony trioxide by inhalation for 12 months followed by post-exposure observation for 24 months	D-11
Table D-12. Watt (1983) study of female rats exposed to antimony trioxide by inhalation for 1 year followed by post-exposure observation for 2 years.....	D-12
Table E.2-1. Effects of antioxidants and inhibitors of oxidative stress related enzymes on cells exposed to compounds containing trivalent antimony.....	E-2
Table E.3-1. Genotoxicity of antimony compounds: Mutations ^{a,b}	E-4
Table E.3-2. Genotoxic DNA damaging effects of antimony compounds	E-6
Table E.3-3. Genotoxicity of antimony compounds – chromosomal aberrations, micronucleus, and sister chromatic exchange ^{a, b, c, d}	E-12
Table E.4-1. Mutations in the lung of mice and rats after two-year inhalation exposure to antimony trioxide (NTP 2017b).	E-20
Table E.5-1. Top 10 upstream regulators for antimony.....	E-22
Table E.5-2. Top ten canonical pathways affected by 6-hour exposure to 20 µM antimony(III) potassium tartrate trihydrate	E-27
Table E.6-1. Effects of compounds containing pentavalent antimony on immunity.....	E-28
Table F.1-1. Neoplasms induced in experimental animal carcinogenicity studies by drinking water studies of antimony potassium tartrate	F-1
Table F.1-2. Cancer studies in experimental animals exposed to antimony(III) potassium tartrate.....	F-1
Table F.1-3. Schroeder <i>et al.</i> (1970) study of male rats and female rats exposed to antimony(III) potassium tartrate in drinking water for the life span of the animals.....	F-3
Table F.1-4. Kanisawa and Schroeder (1969) study of male and female (combined) mice	

exposed to antimony potassium tartrate in drinking water for the lifespan of the animals..... F-4

List of Figures

Figure A-1. Literature search strategy and review..... A-2

Appendix A: Literature Search Strategy

Introduction

The objective of the literature search approach is to identify published literature that is relevant for evaluating the potential carcinogenicity of antimony trioxide (https://ntp.niehs.nih.gov/ntp/about_ntp/bsc/2016/december/meetingmaterials/draftantimonytrioxide_508.pdf). The literature search strategy was used to identify publications in the following areas:

- Properties and human exposure (focusing on the U.S. population)
- Disposition (ADME) and toxicokinetics
- Human cancer studies
- Studies of cancer in experimental Animals
- Mechanistic data and other relevant effects
 - Genetic and related effects
 - Mechanistic considerations


A.1 General approach

Database searching encompasses selecting databases and search terms and conducting the searches. Searches of several citation databases are generally conducted using search terms for antimony, combined with search terms for cancer and/or specific topics, including epidemiological and mechanistic studies. A critical step in the process involves consultation with an information specialist to develop relevant search terms. These terms are used to search bibliographic databases. Table A-1 highlights the general concepts searched and databases consulted. To review all the terms used, please refer to the full search strings in Antimony: RoC Protocol (https://ntp.niehs.nih.gov/ntp/roc/protocols/antimonytrioxide_508.pdf).

Table A-1. Major topics searched

Topic	Search Method	Databases searched
Exposure	Antimony String AND occur*[tiab]	PubMed
Human Studies	Antimony String AND ORoC Epidemiological (Human) Studies Search AND ORoC Cancer Search	PubMed, Scopus, Web of Science
Animal Studies	Antimony String AND Experimental Animals Studies Search AND ORoC Cancer Search	PubMed, Scopus, Web of Science
Mechanism and Genotoxicity	Antimony String AND ORoC Characteristics of Carcinogens Search	PubMed, Scopus, Web of Science

Figure A-1. Literature search strategy and review



A.2 Search strategies

Relevant literature is identified using search terms, data sources, and strategies as discussed below.

General data search: This search covers a broad range of general data sources for information relevant to many or all of the wide range of monograph topics pertaining to antimony.

Exposure-related data search: This search covers a broad range of potential sources for exposure-related information and physical-chemical properties.

Database searches in PubMed, Scopus, and Web of Science: The majority of the primary literature used to draft the antimony monograph was identified from searches of these three extensive databases available through the NIEHS Library. Searches for antimony were combined with the search terms for each of the monograph topics listed above to create the specific literature searches.

Searches for human cancer studies are somewhat unique because they involve the identification of search terms for exposure scenarios that might result in exposure of people to antimony. For antimony, these exposure-related search terms were based on uses of antimony identified from the EPA's TRI database and the Chemical Data Report rule website.

QUOSA library of occupational case-control studies search of the QUOSA-based library of more than 6,000 occupational case-control studies, approximately 95% of which are currently available as searchable full-text pdfs, was conducted using the term "antimony."

Secondary sources: Citations identified from authoritative reviews or from primary references located by literature search, together with publications citing key papers identified using the Web of Science, "Cited Reference Search," were also added.

A.3 Exclusion of treatment for leishmaniasis from human cancer searches

The use of antimony for the treatment of leishmaniasis is considered an intentional medical exposure and out of the scope of this monograph. The large corpus of literature related to leishmaniasis treatment was excluded when identifying human studies. Unlike other parts of the monograph, in which leishmaniasis related content was excluded via search terms, the mechanisms section literature search did not exclude leishmaniasis via the use of search terms. The studies on the *Leishmania* parasite itself were excluded at levels 1 and 2 by reviewers, and studies on the host or cells not infected by leishmaniasis were included for information related to mechanism.

A.4 Updating the literature search

The literature searches will be updated prior to submitting the draft monograph for peer review and prior to finalizing the monograph. Monthly search alerts for antimony searches were created in PubMed, Scopus, and Web of Science, and the results of these searches from the closing date of the initial search will be downloaded for review.

A.5 Review of citations using web-based systematic review software

Citations retrieved from literature searches were uploaded to web-based systematic review software and screened using inclusion and exclusion criteria. Multi-level reviews of the literature were conducted, with initial reviews (Level 1) based on titles and abstracts only to identify citations that could be excluded and to assign the included literature to one or more monograph topics; subsequent reviews (Level 2) for literature assigned to the various monograph topics (Exposure, ADME & TK, Human cancer studies, etc.) were based on full-text (i.e., PDFs) of the papers and were carried out by the writer and scientific reviewer for each monograph section. Two reviewers, at least one of whom is a member of the ORoC at NIEHS, participated at each level of review.

Appendix B: ADME Tables

Table B-1. Antimony(III) trioxide levels^a (µg/g) in red blood cells during a 1-year chronic inhalation exposure (after 6 months and 12 months of exposure) and a 1-year observation period (6 months and 12 months after exposure) in Fischer 344 male and female rats

Group	6 mo	12 mo	18 mo (12 mo exposure and 6 mo obs)	24 mo (12 mo exposure and 12 mo obs)
Males				
I- Control	BDL	BDL	0.17 ± 0.39	BDL
II- 0.055 mg/m ³	0.53 ± 0.31	1.09 ± 0.21	0.86 ± 0.68	BDL
III- 0.51 mg/m ³	5.07 ± 0.29	7.55 ± 0.60	3.93 ± 0.25	2.53 ± 0.27
IV- 4.5 mg/m ³	34.50 ± 3.8	70.70 ± 6.3	38.60 ± 4.8	30.50 ± 7.5
Females				
I- Control	BDL	BDL	BDL	BDL
II- 0.055 mg/m ³	0.74 ± 0.06	1.48 ± 0.10	0.81 ± 0.30	BDL
III- 0.51 mg/m ³	5.69 ± 0.62	9.94 ± 1.32	6.53 ± 0.90	3.39 ± 0.28
IV- 4.5 mg/m ³	75.60 ± 8.4	121.00 ± 10.6	74.60 ± 18.3	36.60 ± 15.5

Source: Newton *et al.* (1994).

mo = month; Below detection limit = BDL (lowest limit of detection = 0.02 µg of antimony/mL, i.e., 0.024 µg of antimony(III) trioxide/mL); obs = observation

^aTotal antimony in red blood cells was reported as total antimony(III) trioxide using the relationship 1 mole Sb₂O₃ = 1.197 mole Sb₂.

Table B-2. Blood antimony concentrations (µg/g blood) in female rats and mice exposed to antimony trioxide (N = 5 except where indicated)

	Day 61	Day 124	Day 269	Day 369	Day 551
Female Mice					
Controls	0.001 ± 0.000	0.001 ± 0.001	0.001 ± 0.000	0.001 ± 0.000	0.001 ± 0.000
3 mg/m ³	0.043 ± 0.002**	0.058 ± 0.001**	0.053 ± 0.006**	0.052 ± 0.003**	0.061 ± 0.010**
10 mg/m ³	0.083 ± 0.002**	0.089 ± 0.002**	0.091 ± 0.002**	0.088 ± 0.003**	0.087 ± 0.004**
30 mg/m ³	0.141 ± 0.003**	0.148 ± 0.005**	0.163 ± 0.008** ^a	0.137 ± 0.007**	0.163 ± 0.006** ^a
Female Rats					
Controls	0.139 ± 0.012	0.050 ± 0.002	0.077 ± 0.002	0.084 ± 0.008	0.066 ± 0.005
3 mg/m ³	7.352 ± 0.375**	16.135 ± 0.995**	39.590 ± 3.915**	50.917 ± 2.296**	63.297 ± 3.906**
10 mg/m ³	18.079 ± 0.793**	40.350 ± 1.543**	88.833 ± 2.210**	102.083 ± 2.738**	149.192 ± 8.472** ^a
30 mg/m ³	43.574 ± 1.741**	96.082 ± 3.940**	175.437 ± 6.471**	200.239 ± 10.302**	231.934 ± 8.681**

Source: NTP (2017b).

**Significantly different ($P < 0.01$) from the chamber control group by Shirley's test.

^aN = 4.

Table B-3. Tissue distribution of antimony (μg antimony/g tissue) in rats after oral exposure to antimony(III) trioxide by gavage or in the diet

Tissue	Controls (M/F) ^a	1000 mg/kg Sb ₂ O ₃ suspension p.o. for 1 day (M/F) ^a	1000 mg/kg Sb ₂ O ₃ suspension p.o. for 14 days (M/F) ^a	2% Sb ₂ O ₃ in diet* for 49 days ^b	2% Sb ₂ O ₃ in diet* for 8 months ^c
Thyroid	0.098/0.195	1.507/2.103	2.639/2.280	88.9	156.0
Adrenal	NR	NR	NR	67.8	NR
Lung	0.004/0.002	0.041/0.061	0.746/0.882	14.0	3.7
Spleen	0.010/0.032	0.197/0.113	1.485/1.386	18.9	8.1
Heart	0.004/0.003	0.042/0.041	0.643/0.356	7.6	5.1
Kidney	0.003/0.002	0.012/0.023	0.323/0.261	6.7	6.0
Liver	0.004/0.003	0.041/0.064	0.823/0.675	8.9	15.5
Bone marrow	0.080/0.142	1.192/1.996	2.486/3.517	NR	NR
Bone or femur	0.019/0.010	0.048/0.032	0.254/0.265	NR	2.5
Muscle	0.003/0.003	0.005/0.005	0.039/0.044	NR	0.3
Whole blood	0.003/0.003	0.708/0.640	8.278/6.886	NR	NR

Sources: ^a TNO Quality of Life 2005 as cited by EU 2008; ^b Westrick 1953; ^c Gross *et al.* 1955 as cited by EU 2008.

F = female; M = male; NR = not reported; p.o. = per os (by mouth).

*Based on consumption of 5 g of food per day per 100 g body weight (Johns Hopkins University 2017), rats exposed to 2% Sb₂O₃ in the diet or by gavage at 1,000 mg/kg body weight would be exposed to approximately 0.1 g per 100 g body weight.

Appendix C: Human Studies Tables

Table C-1. Evaluation of selection bias in human cancer studies.

Study	Selection bias
Jones 1994	<i>Rating: ++; Direction: ↓</i> <i>Rationale:</i> Only an external analysis was conducted. Although the impact of healthy worker survivor effect (HWSE) is mitigated by stratification by time-since-exposure, HWSE is still possible and may bias results toward the null.
Schnorr <i>et al.</i> 1995	<i>Rating: ++; ↓</i> <i>Rationale:</i> Only an external analysis was conducted. HWSE was not accounted for in this analysis, which may result in an underestimating of the risk estimates.
Jones <i>et al.</i> 2007	<i>Rating: ++; ↓</i> <i>Rationale:</i> Missing death information for 5.7% of untraced individuals would slightly bias results if they experienced the outcome. HWSE was not accounted for in the analyses, however, the impact of the smelter closing during follow-up would reduce the residual survival advantage.
Wingren and Axelson 1993	<i>Rating: +++; ↔</i> <i>Rationale:</i> Cases and controls were selected from the same parishes. No evidence suggests that the selection of subjects was related to both antimony exposure and disease.

↑ = Results bias away from the null; ↓ = Results bias toward the null; ↔ = Unknown direction of bias

Table C-2. Evaluation of exposure assessment methods in human cancer studies.

Study	Exposure assessment rating
Jones 1994	<i>Rating: ++/+++; Direction: ↔</i> <i>Rationale:</i> Exposure assessment methods have decent sensitivity and specificity, leading to reliable classification with respect to ever-exposure to antimony and exposure duration. Antimony exposure is assumed based on job description at smelter site.
Schnorr <i>et al.</i> 1995	<i>Rating: ++; ↓</i> <i>Rationale:</i> Exposure was reliably characterized as ever-exposure to antimony and duration of antimony exposure, but not with respect to concentration of exposure. Based on the reported environmental sampling data, antimony air exposure varied by plant location and year sampled; however, exposure is not captured at the individual level due to lack of information on job duties, and therefore, may be subject to misclassification.
Jones <i>et al.</i> 2007	<i>Rating: ++; ↓</i> <i>Rationale:</i> Given the modeling efforts used to account for the uncertainty in early air contamination levels, and because air sampling concentrations are likely an underestimate of true individual antimony exposure, exposure levels and timing may not represent true antimony concentrations and worker exposure prior to 1972. Authors mention changes in plant processes before NIOSH collected exposure estimates. The 3 scenarios for back-extrapolation (1. twice as high air concentrations in 1937, 2. average concentration from 1937 to 1972, and 3. a doubling in concentrations from 1937-1960 then a decrease to 1972 levels) are assumptions based on little empirical data.
Wingren and Axelson 1993	<i>Rating: +; ↑</i> <i>Rationale:</i> Exposure to antimony was based on reported job title at death. Those classified as unexposed who may have worked in a glass producing facility or had other antimony occupational exposure over a lifetime may have misclassified exposure. Reported level of antimony used by surveyed glass working facilities may not represent individual-level

Study	Exposure assessment rating
	exposure to employees. Facility surveys of antimony use was taken at one time point; unknown if antimony use patterns were consistent.

↑ = Results bias away from the null; ↓ = Results bias toward the null; ↔ = Unknown direction of bias

Table C-3. Evaluation of outcome assessment in human cancer studies.

Study	Outcome assessment rating
Jones 1994	<i>Rating:</i> +++; <i>Direction:</i> ↔ <i>Rationale:</i> Outcome methods distinguish between diseased and non-diseased subjects. Follow-up and diagnoses are conducted independent of exposure status.
Schnorr <i>et al.</i> 1995	<i>Rating:</i> +++; ↔ <i>Rationale:</i> Outcome methods distinguish between diseased and non-diseased subjects. Follow-up and diagnoses are conducted independent of exposure status.
Jones <i>et al.</i> 2007	<i>Rating:</i> +++; ↔ <i>Rationale:</i> Outcome methods distinguish between diseased and non-diseased subjects. Follow-up and diagnoses are conducted independent of exposure status.
Wingren and Axelson 1993	<i>Rating:</i> ++; ↑ <i>Rationale:</i> Outcome methods distinguish between diseased and non-diseased subjects. Occupational title (i.e. exposure status) was collected from the death and burial register, which noted mortality status. Given the lack of information on the blinding status, diagnostic bias cannot be ruled out. If coder identified diseased subjects as being exposed, it would bias the results away from the null.

↑ = Results bias away from the null; ↓ = Results bias toward the null; ↔ = Unknown direction of bias

Table C-4. Evaluation of study sensitivity in human cancer studies.

Study	Sensitivity rating
Jones 1994	<i>Rating:</i> ++; <i>Direction:</i> ↔ <i>Rationale:</i> Study has few exposed cases but a substantial duration of exposure with a long range for follow-up. Stratification by exposure duration and years increase sensitivity.
Schnorr <i>et al.</i> 1995	<i>Rating:</i> ++; ↔ <i>Rationale:</i> Study had a small-to-moderate number of exposed cases. There was adequate duration for follow-up, with a substantial duration of exposure. Duration and ever-exposure were measured, but not the range of antimony concentrations.
Jones <i>et al.</i> 2007	<i>Rating:</i> +; ↔ <i>Rationale:</i> Adequate number of potentially-exposed subjects but a small number of exposed cases. Exposure characterized by job-exposure matrix and detailed work histories. Exposure was modeled with a substantial range and level of exposure with adequate duration for latency. However, exposure was not at an individual level and exposure was extrapolated based on assumptions.
Wingren and Axelson 1993	<i>Rating:</i> +; ↔ <i>Rationale:</i> Study captures variability in antimony use by parish where cases and controls died. However, given the unknown number of exposed subjects, exposed cases, the unknown number of controls, and the unknown individual-level exposure to antimony in glass workers, this study has poor sensitivity.

↑ = Results bias away from the null; ↓ = Results bias toward the null; ↔ = Unknown direction of bias

Table C-5. Evaluation of potential for confounding bias for human cancer studies.

Study	Confounding rating
Jones 1994	<i>Rating: +; Direction: ↑</i> <i>Rationale:</i> No control for smoking or occupational co-exposures in statistical analysis. Likely co-exposure to arsenic and PAHs (lung carcinogens) based on smelting source materials. Smoking not controlled for, despite high prevalence in the study population.
Schnorr <i>et al.</i> 1995	<i>Rating: +++; ↔</i> <i>Rationale:</i> No control for smoking or occupational co-exposures in statistical analysis. Confounding from occupational co-exposures to arsenic and lead are minimal based on source information and environmental testing. Smoking prevalence rates were assumed to be low in this particular population.
Jones <i>et al.</i> 2007	<i>Rating: ++; ↑</i> <i>Rationale:</i> No attempt to statistically account for measured occupational co-exposures in analysis. High level of correlation between antimony, lead, and arsenic air concentrations suggests likely occupational co-exposure. Minimal concern for smoking, but not controlled for in analysis.
Wingren and Axelson 1993	<i>Rating: +; ↑</i> <i>Rationale:</i> Smoking and occupational co-exposures lead and asbestos were not statistically controlled for in analysis. Lead and antimony use patterns were highly correlated, and lead was associated with an increased risk of stomach cancer mortality in this population; therefore, risk of confounding bias is high.

↑ = Results bias away from the null; ↓ = Results bias toward the null; ↔ = Unknown direction of bias

Table C-6. Evaluation of analysis and selective reporting for human cancer studies.

Study	Analysis rating	Reporting rating
Jones 1994	<i>Rating: +++</i> <i>Rationale:</i> The study used relevant data and appropriate assumptions and methods of analysis.	<i>Rating: +++</i> <i>Rationale:</i> No evidence that reporting of the data or analyses were limited to only a subset of data that were collected.
Schnorr <i>et al.</i> 1995	<i>Rating: ++</i> <i>Rationale:</i> 95% CI are generally favorable to 90% CI regardless of <i>a priori</i> outcome status.	<i>Rating: +++</i> <i>Rationale:</i> No evidence that reporting of the data or analyses were limited to only a subset of the data that were collected.
Jones <i>et al.</i> 2007	<i>Rating: ++</i> <i>Rationale:</i> 95% CI are generally favorable to 90% CI regardless of <i>a priori</i> outcome status.	<i>Rating: +++</i> <i>Rationale:</i> No evidence that reporting of the data or analyses were limited to only a subset of the data that were collected.
Wingren and Axelson 1993	<i>Rating: ++</i> <i>Rationale:</i> 95% CI are generally favorable to 90% CI regardless of <i>a priori</i> outcome status.	<i>Rating: ++</i> <i>Rationale:</i> It is unknown whether reporting was done on only a subset on data. Sample size for cases, controls, and exposure groups were not reported.

Appendix D: Animal Study Quality Tables

Table D-1. Schroeder *et al.* (1970) study of male rats exposed to antimony potassium tartrate in the drinking water

Utility question	Rating	Rationale
Study design		
Randomization	NR	Randomization and initial body weights were not reported.
Controls	+++	Concurrent control group, exposed to untreated drinking water, had the same number of animals as exposure group did.
Historical data	No	
Animal model	+++	Both sexes of a random bred strain, which increases external validity.
Statistical power	+++	Large numbers of animals (51 males, 59 females) per concentration group were used.
Exposure		
Chemical characterization	NR	Not reported, not even purity.
Dosing regimen	++	The maximally tolerated dose level was not reached, because the treated group did not show decreased body weight compared to the control group, although the median life spans and longevity (mean age of the last surviving 10%) for both sexes were decreased by the treatment. The dose might not have been high enough to detect neoplastic effects.
Exposure duration	+++	Though exposure duration was never clearly stated, this study appears to use a life time exposure.
Dose-response	+	Only one dose level was tested and no basis for that level was reported. All other elements were administered at the same level, except for lead.
Outcome		
Pathology	+	Only grossly visible tumors were reported. The methods stated that gross tumors were fixed, but did not state that they were stained or microscopically examined. Consequently, small tumors might have been missed.
Consistency between groups	++	A pneumonia epidemic killed many rats and the death rates varied among the groups.
Study duration	+++	Life time study, because the animals were observed until their nature death (as compared to scheduled euthanization after a predetermined exposure period).
Confounding		
Confounding	++	Pneumonia killed various numbers of animals per group, before penicillin treatment controlled the disease. It is unclear that all rats, or only visibly sick rats, received penicillin. Furthermore, the disease might in effect select stronger/healthier animals (than the general population) to complete the study. Additionally, test substance purity was unknown.
Reporting and analysis		
Reporting data and statistics	++	The statistical methods and results of survival measures were reported, but statistical analysis of tumor incidences were not reported.
Combining lesions	+	Tumors were counted based on gross observation, not histological analysis occurred.
Overall utility: +. The study has low utility because of many limitations, including only reporting grossly visible tumors without organ site or tumor type.		

Table D-2. Schroeder *et al.* (1970) study of female rats exposed to antimony potassium tartrate in the rinking water

Utility question	Rating	Rationale
Study design		
Randomization	NR	Randomization and initial body weights were not reported.
Controls	+++	Concurrent control group, exposed to untreated drinking water, had the same number of animals as exposure group did.
Historical data	No	
Animal model	+++	Both sexes of a random bred strain, which increases external validity.
Statistical power	+++	Large numbers of animals (51 males, 59 females) per concentration group were used.
Exposure		
Chemical characterization	NR	Not reported, not even purity.
Dosing regimen	++	The maximally tolerated dose level was not reached, because the treated group did not show decreased body weight compared to the control group, although the median life spans and longevity (mean age of the last surviving 10%) for both sexes were decreased by the treatment. The dose might not have been high enough to detect neoplastic effects.
Exposure duration	+++	Though exposure duration was never clearly stated, this study appears to use a life time exposure.
Dose-response	+	Only one dose level was tested and no basis for that level was reported.
Outcome		
Pathology	+	Only grossly visible tumors were reported. The methods stated that gross tumors were fixed, but did not state that they were stained or microscopically examined. Consequently, small tumors might have been missed.
Consistency between groups	++	A pneumonia epidemic killed many rats and the death rates varied among the groups.
Study duration	+++	Life time study, because the animals were observed until their nature death (as compared to scheduled euthanization after a predetermined exposure period).
Confounding		
Confounding	++	Pneumonia killed various numbers of animals per group, before penicillin treatment controlled the disease. It is unclear that all rats, or only visibly sick rats, received penicillin. Furthermore, the disease might in effect select stronger/healthier animals (than the general population) to complete the study. Additionally, test substance purity was unknown.
Reporting and analysis		
Reporting data and statistics	++	The statistical methods and results of survival measures were reported, but statistical analysis of tumor incidences were not reported.
Combining lesions	+	Tumors were counted based on gross observation, not histological analysis occurred.
Overall utility: +. The study has low utility because of many limitations, including only reporting grossly visible tumors without organ site or tumor type.		

Table D-3. Kanisawa and Schroeder (1969) and Schroeder *et al.* (1968) study of male and female (combined) mice exposed to antimony potassium tartrate in drinking water for the lifespan of the animals

Utility question	Rating	Rationale
Study design		
Randomization	NR	Not reported.
Controls	+++	Concurrent control group, exposed to doubly deionized water with added essential trace elements, had the same number of animals as the exposure group did.
Historical data		No
Animal model	+++	Both sexes of random bred mice were used, giving a high level of external validity.
Statistical power	+++	A large number (54) of mice per sex per group were used.
Exposure		
Chemical characterization	NR	No chemical characterization was reported, not even purity.
Dosing regimen	+	The maximally tolerated dose level was not reached, because the treated group did not show decreased body weight compared to the control group. The dose might not have been high enough to detect neoplastic effects.
Exposure duration	+++	Mice were exposed for their lifetimes.
Dose-response	+	Only one concentration was tested and no rationale for the dose selection was reported. Dose response relationships cannot be evaluated due to only one dose level.
Outcome		
Pathology	++	Only gross lesions were microscopically evaluated.
Consistency between groups	++	The treated and control groups were treated the same while mice were alive. The examination of organs/tissues varied, because only gross lesions (not all major organs) were examined microscopically.
Study duration	+++	The study duration was lifetime, up to the animals' natural death.
Confounding		
Confounding	+++	Testing substance purity and supplier were unknown. Exposure to antimony via other sources (feed, housing) was negligible because the feed was antimony free and metal exposure via housing was minimized.
Reporting and analysis		
Reporting data and statistics	++	Although tumor incidents were not statistically analyzed in the study, the data were reported and enabled us to conduct statistical analysis. Statistical methods were described as "numerical data were treated by Chi-square analysis and by Student's t test", but the reported probability in tables did not specify the result was from which method.
Combining lesions	++	Tumor incidence was reported for two sexes combined only, instead of male and female separately. Site specific (lung, liver, mammary gland, and other) information was limited to tumor incidence, with no subtype. Tumors were also grouped by origin (epithelial, non-epithelial) along with being benign or malignant. Overall information did not allow detecting of specific tumor type increase in either sex.

Overall utility: +. Due to many limitations, including only one tested concentration (below maximally tolerated dose), unknown test substance purity, tumor incidences only reported in combined sexes with no histologic information, and lack of site specific information (except incidences of three sites in sexes combined), this study is of low utility. Data lack sufficient details to allow us determine whether any specific type of tumor had increased.

Table D-4. NTP (2017b) study of male rats exposed to antimony trioxide by inhalation for 105 weeks

Utility question	Rating	Rationale
Study design		
Randomization	+++	Animals were randomly assigned to groups.
Controls	+++	Concurrent chamber control was used. Data was also compared with historical control.
Historical data		Yes
Animal model	+++	Standard model.
Statistical power	+++	A large number, 50/sex/group, of animals were used.
Exposure		
Chemical characterization	+++	The chemical and exposure chamber were well characterized, showing high purity, stability, and homogeneity. Concentration inside the exposure chamber was measured in real-time and alarmed if readings were not within limits of acceptable concentrations. Aerosol size, measured monthly, was also consistently less than 4 μm (MMAD = 1-1.4 μm , GSD 1.8-2.2. Stability in the generation and exposure system was tested before the test, during the test (at 4 weeks for rats), and the end of two-year study.
Dosing regimen	+++	Consistent and very close to target concentrations. The highest exposure level was near maximally tolerated levels as evidenced by the trend of decreased survival and a significant decrease in body weight. Neoplasms were significantly increased.
Exposure duration	+++	Exposure duration was near life-span
Dose-response	+++	Three dose levels spanning a range of 30 fold were used.
Outcome		
Pathology	+++	Detailed and covering all tissues. Full necropsy with histological exam of all major organs was conducted and verified by an independent quality control pathologist.
Consistency between groups	+++	Nothing was reported to suggest that animals from different groups were treated differently.
Study duration	+++	Near life-span study duration was used.
Confounding		
Confounding	+++	No concerns of confounding were reported.
Reporting and analysis		
Reporting data and statistics	+++	Statistical analysis was clearly reported.
Combining lesions	+++	No indication of concern. Detailed groupings were provided.

Overall utility: +++. There were no concerns of confounding as the chemical was pure, stable, the exposure was well characterized, and all groups were treated the same. The study had a high level of sensitivity to detect neoplasms as it used large numbers of both sexes of rats, exposed at three dose levels, which reached the maximally tolerated level, for a near life-span duration. However, the stock of rat used was new to NTP and so little historical control data existed compared to the previously used Fischer 344 rat stock. Complete necropsies with histological examination of most organs was performed, so the ability to detect neoplasms was high.

Table D-5. NTP (2017b) study of female rats exposed to antimony trioxide by inhalation for 105 weeks

Utility question	Rating	Rationale
Study design		
Randomization	+++	Animals were randomly assigned to groups
Controls	+++	Concurrent chamber control was used. Data was also compared with historical control.
Historical data		No
Animal model	+++	Standard model.
Statistical power	+++	A large number, 50/sex/group, of animals were used.
Exposure		
Chemical characterization	+++	The chemical and exposure chamber were well characterized, showing high purity, stability, and homogeneity. Concentration inside the exposure chamber was measured in real-time and alarmed if readings were not within limits of acceptable concentrations. Aerosol size, measured monthly, was also consistently less than 4 μm (MMAD = 0.9-1.5 μm , GSD = 1.7-2.1). Stability in the generation and exposure system was tested before the test, during the test (at 4 weeks for rats), and the end of two-year study.
Dosing regimen	+++	Consistent, and very close to target, concentrations. The highest exposure level was near maximally tolerated levels as evidenced by the trend of decreased survival and a significant decrease in body weight. Neoplasms were significantly increased.
Exposure duration	+++	Exposure duration was near life-span.
Dose-response	+++	Three dose levels spanning a range of 30 fold were used.
Outcome		
Pathology	+++	Detailed and covering all tissues. Full necropsy with histological exam of all major organs was conducted and verified by an independent quality control pathologist.
Consistency between groups	+++	Nothing was reported to suggest that animals from different groups were treated differently.
Study duration	+++	Near life-span study duration was used.
Confounding		
Confounding	+++	No concerns of confounding were reported.
Reporting and analysis		
Reporting data and statistics	+++	Statistical analysis was clearly reported.
Combining lesions	+++	No indication of concern. Detailed groupings were provided.

Overall utility: +++. There were no concerns of confounding as the chemical was pure, stable, the exposure was well characterized, and all groups were treated the same. The study had a high level of sensitivity to detect neoplasms as it used large numbers of both sexes of rats, exposed at three dose levels, which reached the maximally tolerated level, for a near life-span duration. However, the stock of rat used was new to NTP and so little historical control data existed compared to the previously used Fisher 344 rat stock. Complete necropsies with histological examination of most organs was performed, so the ability to detect neoplasms was high.

Table D-6. NTP (2017b) study of male mice exposed to antimony trioxide by inhalation for 105 weeks

Utility question	Rating	Rationale
Study design		
Randomization	+++	Animals were randomly assigned to groups.
Controls	+++	Concurrent chamber control was used. Data was also compared with historical control.
Historical data		Yes
Animal model	+++	Standard model.
Statistical power	+++	A large number, 50/sex/group, of animals were used.
Exposure		
Chemical characterization	+++	The chemical and exposure chamber were well characterized, showing high purity, stability, and homogeneity. Concentration inside the exposure chamber was measured in real-time and alarmed if readings were not within limits of acceptable concentrations. Aerosol size, measured monthly, was also consistently less than 4 μm (MMAD = 0.9-1.5 μm , GSD 1.7-2.1). Stability in the generation and exposure system was tested before the test, during the test (at 4 weeks for rats), and the end of two-year study.
Dosing regimen	+++	Consistent, and very close to target, concentrations. The highest exposure level was near maximally tolerated levels as evidenced by the trend of decreased survival and a significant decrease in body weight. Neoplasms were significantly increased.
Exposure duration	+++	Exposure duration was near life-span.
Dose-response	+++	Three dose levels spanning a range of 30 fold were used.
Outcome		
Pathology	+++	Detailed and covering all tissues. Full necropsy with histological exam of all major organs was conducted and verified by an independent quality control pathologist.
Consistency between groups	+++	Nothing was reported to suggest that animals from different groups were treated differently.
Study duration	+++	Near life-span study duration was used.
Confounding		
Confounding	+++	No concerns of confounding were reported.
Reporting and analysis		
Reporting data and statistics	+++	Statistical analysis was clearly reported.
Combining lesions	+++	No indication of concern. Detailed groupings were provided.

Overall utility: +++. There were no concerns of confounding as the chemical was pure, stable, the exposure was well characterized, and all groups were treated the same. The study had a high level of sensitivity to detect neoplasms as it used large numbers of both sexes of mice, exposed at three dose levels, which reached the maximally tolerated level, for a near life-span duration. Complete necropsies with histological examination of most organs was performed, so the ability to detect neoplasms was high.

Table D-7. NTP (2017b) study of female mice exposed to antimony trioxide by inhalation for 105 weeks

Utility question	Rating	Rationale
Study design		
Randomization	+++	Animals were randomly assigned to groups.
Controls	+++	Concurrent chamber control was used. Data were also compared with historical control.
Historical data		No
Animal model	+++	Standard model.
Statistical power	+++	A large number, 50/sex/group, of animals were used.
Exposure		
Chemical characterization	+++	The chemical and exposure chamber were well characterized, showing high purity, stability, and homogeneity. Concentration inside the exposure chamber was measured in real-time and alarmed if readings were not within limits of acceptable concentrations. Aerosol size, measured monthly, was also consistently less than 4 μm (MMAD = 0.9 - 1.5 μm , GSD = 1.7 - 2.1). Stability in the generation and exposure system was tested before the test, during the test (at 4 weeks for rats), and the end of two-year study.
Dosing regimen	+++	Consistent, and very close to target, concentrations. The highest exposure level was near maximally tolerated levels as evidenced by the trend of decreased survival and a significant decrease in body weight. Neoplasms were significantly increased.
Exposure duration	+++	Exposure duration was near life-span.
Dose-response	+++	Three dose levels spanning a range of 30 folds were used.
Outcome		
Pathology	+++	Detailed and covering all tissues. Full necropsy with histological exam of all major organs was conducted and verified by an independent quality control pathologist.
Consistency between groups	+++	Nothing was reported to suggest that animals from different groups were treated differently.
Study duration	+++	Near life-span study duration was used.
Confounding		
Confounding	+++	No concerns of confounding were reported.
Reporting and analysis		
Reporting data and statistics	+++	Statistical analysis was clearly reported.
Combining lesions	+++	No indication of concern. Detailed groupings were provided.

Overall utility: +++. There were no concerns of confounding as the chemical was pure, stable, the exposure was well characterized, and all groups were treated the same. The study had a high level of sensitivity to detect neoplasms as it used large numbers of both sexes of mice, exposed at three dose levels, which reached the maximally tolerated level, for a near life-span duration. Complete necropsies with histological examination of most organs was performed, so the ability to detect neoplasms was high.

Table D-8. Groth *et al.* (1986) study of male rats exposed to antimony trioxide by inhalation for 53 weeks followed by post-exposure observation for 71 to 73 weeks

Utility question	Rating	Rationale
Study design		
Randomization	NR	Not reported.
Controls	+++	Concurrent untreated chamber controls (filtered air) were used.
Historical data		No
Animal model	+++	Male and female inbred rats were used.
Statistical power	+++	A large number of rats (90/sex/group) were used.
Exposure		
Chemical characterization	++	The purity was estimated to be 95.8%, however the authors did not explicitly report the purity of antimony trioxide. The purity of elemental antimony was reported as 80% and NIH calculated this to mean a purity of antimony trioxide of 95.8%, assuming all of the antimony present was as antimony trioxide. Trace levels of contamination by carcinogens (arsenic and lead) and others (tin, cesium aluminum, and bromine) were not considered to have significantly contributed to carcinogenic effects.. The aerosol concentrations didn't reach target levels of 50 mg/m ³ until after 5 months of adjustment and modifications on the exposure equipment. MMAD of aerosol of 2.80 µm was fine, but the GSD was not reported. Aerosol size appeared only measured once at 6 month of exposure.
Dosing regimen	++	The exposure level was based on the middle of the concentration range that workers are exposed to, so it would not be expected to be at the maximally tolerated level. The concentrations fluctuated dramatically, up to 191.1 mg/m ³ for daily TWAs, while a mean daily TWA was 45, 46 mg/m ³ (two chambers) and the target concentration was 50 mg/m ³ . It is not clear whether the chamber air was humidified. Survival and body weights were similar to controls, but total neoplasms in the lung were significantly increased over untreated controls.
Exposure duration	++	Exposure duration was 53 weeks.
Dose-response	+	Only one exposure level was used and it was based on the middle of the concentration range that workers are exposed to (i.e., well below animals' maximal tolerated dose). The actual exposure concentration fluctuated greatly until about 5 months into the study when the target concentration was reached.
Outcome		
Pathology	+++	Most organs were histologically examined.
Consistency between groups	++	No indication of differential treatments.
Study duration	++	The study duration was 71 to 73 weeks long, roughly 1.4 years, with only 5 months of observation after the end of 53 week-long exposure. These lengths were likely limited because the rats were 8 months old at the beginning of the study.
Confounding		
Confounding	+	The chemical was only 80% antimony, with various other metal contaminants, such as tin, lead, cesium, aluminum, arsenic, and bromine. Lead and arsenic are carcinogenic.
Reporting and analysis		
Reporting data and statistics	+++	Statistical analysis was reported for body weights, tissue levels of antimony. Neoplasms were not reported as they were stated to have not occurred.
Combining lesions	+++	Neoplasms were combined by site. While no numbers of each pathological type were provided, the tumor combining is fine

Overall utility: ++. The chemical was not fully characterized, but was estimated to be 95.8% pure, with trace levels of lead and arsenic as contaminants, that were not considered to have significantly contributed to carcinogenic effects. The sensitivity of the study to detect neoplasms was low as only one dose level was used and it was based on the level of exposure to workers and not

the maximally tolerated dose. Further, the exposure concentration varied widely until 5 months into the study when the target concentration was reached. The exposure duration was more than a year and full necropsies with histological examinations were performed. Neoplasms were reported with statistical analysis as total neoplasms combined per organ site.

Table D-9. Groth *et al.* (1986) study of female rats exposed to antimony trioxide by inhalation for 53 weeks followed by post-exposure observation for 71 to 73 weeks

Utility question	Rating	Rationale
Study design		
Randomization	NR	Not reported.
Controls	+++	Concurrent untreated chamber controls (filtered air) were used.
Historical data		No
Animal model	+++	Male and female inbred rats were used.
Statistical power	+++	A large number of rats (90/sex/group) were used.
Exposure		
Chemical characterization	++	The purity was estimated to be 95.8%, however the authors did not explicitly report the purity of antimony trioxide. The purity of elemental antimony was reported as 80% and NIH calculated this to mean a purity of antimony trioxide of 95.8%, assuming all of the antimony present was as antimony trioxide. Trace levels of contamination by carcinogens (arsenic and lead) and others (tin, cesium aluminum, and bromine) were not considered to have significantly contributed to carcinogenic effects. The aerosol concentrations didn't reach target levels of 50 mg/m ³ until after 5 months of adjustment and modifications on the exposure equipment. MMAD of aerosol of 2.80 µm was fine, but the GSD was not reported. Aerosol size appeared only measured once at 6 month of exposure.
Dosing regimen	++	The exposure level was based on the middle of the concentration range that workers are exposed to, so it would not be expected to be at the maximally tolerated level. The concentrations fluctuated dramatically, up to 191.1 mg/m ³ for daily TWAs, while a mean daily TWA was 45, 46 mg/m ³ (two chambers) and the target concentration was 50 mg/m ³ . It is not clear whether the chamber air was humidified. Survival and body weights were similar to controls, but total neoplasms in the lung were significantly increased over untreated controls.
Exposure duration	++	Exposure duration was 53 weeks.
Dose-response	+	Only one exposure level was used and it was based on the middle of the concentration range that workers are exposed to (i.e., well below animals' maximal tolerated dose). The actual exposure concentration fluctuated greatly until about 5 months into the study when the target concentration was reached.
Outcome		
Pathology	+++	Most organs were histologically examined.
Consistency between groups	++	No indication of differential treatments.
Study duration	++	The study duration was 71 to 73 weeks long, roughly 1.4 years, with only 5 months of observation after the end of 53 week-long exposure. These lengths were likely limited because the rats were 8 months old at the beginning of the study.
Confounding		
Confounding	+	The chemical was only 80% antimony, with various other metal contaminants, such as tin, lead, cesium, aluminum, arsenic, and bromine. Lead and arsenic are carcinogenic
Reporting and analysis		
Reporting data and statistics	+++	Statistical analysis was reported for body weights, tissue levels of antimony. Statistical significance was not reported for neoplasm incidences, but was calculated by NTP.

Utility question	Rating	Rationale
Combining lesions	+++	Neoplasms were combined by site. While no numbers of each pathological type were provided, the tumor combining is fine.

Overall utility: ++. The chemical was not fully characterized, but was estimated to be 95.8% pure, with trace levels of lead and arsenic as contaminants, that were not considered to have significantly contributed to carcinogenic effects. The sensitivity of the study to detect neoplasms was low as only one dose level was used and it was based on the level of exposure to workers and not the maximally tolerated dose. Further, the exposure concentration varied widely until 5 months into the study when the target concentration was reached. The exposure duration was more than a year and full necropsies with histological examinations were performed. Neoplasms were reported with statistical analysis as total neoplasms combined per organ site.

Table D-10. Newton *et al.* (1994) study of male rats exposed to antimony trioxide by inhalation for 12 months followed by post-exposure observation for 24 months

Utility question	Rating	Rationale
Study design		
Randomization	+++	Used a computer program to randomly sort animals so that mean group weights were comparable.
Controls	+++	Use concurrent control at the same number of animals as exposure groups.
Historical data		No
Animal model	+++	Normal Fischer rats, which are often used in carcinogenicity studies.
Statistical power	+++	A large number of rats (65/sex/group) were used.
Exposure		
Chemical characterization	+++	A blend of lots from 9 producers of antimony trioxide. Highly pure material. Particle size was characterized as having a mass median aerodynamic diameter (MMAD) of 3.76 +/- 0.84 µm and a geometric standard deviation (GSD) of 1.79 +/- 0.32. Exposure concentration was analyzed four times a day and particle sizes were analyzed before the study and every three months. Homogeneity of the exposure chamber was verified by measuring 10 different locations.
Dosing regimen	++	There were not differences in body weight, survival, or neoplasm incidence, suggesting the dose was not at the maximally tolerated dose.
Exposure duration	+++	12-month exposure.
Dose-response	+++	Three exposed levels were used, which covered a 100-fold range.
Outcome		
Pathology	++	Only heart, airway, and peribronchial lymph nodes were histologically examined.
Consistency between groups	+++	Consistent treatment and evaluation of groups.
Study duration	+++	The study duration was 2 years, with 12 months of exposure and 12 months of observation.
Confounding		
Confounding	+++	Material of high purity. Animal husbandry reported in detail. No significant body weight loss in the treated groups, compared to the control.
Reporting and analysis		
Reporting data and statistics	+++	Since neoplasm incidences were not reported, as they were negative, statistical analysis wasn't reported.
Combining lesions	+++	No tumor combining, as only three cases [2 males (including one from control), 1 female] were seen.

Overall utility: ++. There was little concern for confounding as the chemical was pure, exposure conditions were well characterized, and groups were treated consistently with animals randomly assigned to exposure groups. The sensitivity of detecting neoplasms was good as high numbers of both sexes were tested. Exposure were at three concentrations for about half a life-span duration (1 year), though observations (1 year) continued to a near life-span duration. However, the highest exposure level did not reach the maximally tolerated level. Most organs were histologically examined, so most neoplasms had the ability of being detected. Although aerosol size was not ideal (slightly over the current upper limit of test guidelines), this paper did show Sb₂O₃ accumulation and decreased clearance in the lung (by 80% in the 4.5 mg/m³ group). For inert particle, such overload would/could cause lung tumor. The overload was observed at relatively low exposure concentrations (compared to inert particles, such as TiO₂) and Sb₂O₃ toxicity was suspected. It appears conditions that could potentially lead to cancer did persist (Table 9, post-exposure, chronic inflammation in most animals, although hyperplasia was in only very few animals).

Table D-11. Newton *et al.* (1994) study of female rats exposed to antimony trioxide by inhalation for 12 months followed by post-exposure observation for 24 months

Utility question	Rating	Rationale
Study design		
Randomization	+++	Used a computer program to randomly sort animals so that mean group weights were comparable.
Controls	+++	Use concurrent control at the same number of animals as exposure groups.
Historical data		No
Animal model	+++	Normal Fischer rats, which are often used in carcinogenicity studies.
Statistical power	+++	A large number of rats (65/sex/group) were used.
Exposure		
Chemical characterization	+++	A blend of lots from 9 producers of antimony trioxide. Highly pure material. Particle size was characterized as having a mass median aerodynamic diameter (MMAD) of 3.76 +/- 0.84 µm and a geometric standard deviation (GSD) of 1.79 +/- 0.32. Exposure concentration was analyzed four times a day and particle sizes were analyzed before the study and every three months. Homogeneity of the exposure chamber was verified by measuring 10 different locations.
Dosing regimen	++	There were not differences in body weight, survival, or neoplasm incidence, suggesting the dose was not at the maximally tolerated dose.
Exposure duration	+++	12-month exposure.
Dose-response	+++	Three exposed levels were used, which covered a 100-fold range.
Outcome		
Pathology	++	Only heart, airway, and peribronchial lymph nodes were histologically examined.
Consistency between groups	+++	Consistent treatment and evaluation of groups.
Study duration	+++	The study duration was 2 years, with 12 months of exposure and 12 months of observation.
Confounding		
Confounding	+++	Material of high purity. Animal husbandry reported in detail. No significant body weight loss in the treated groups, compared to the control.
Reporting and analysis		
Reporting data and statistics	+++	Since neoplasm incidences were not reported, as they were negative, statistical analysis wasn't reported.
Combining lesions	+++	No tumor combining, as only three cases [2 males (including one from control), 1 female] were seen.

Overall utility: ++. There was little concern for confounding as the chemical was pure, exposure conditions were well characterized, and groups were treated consistently with animals randomly assigned to exposure groups. The sensitivity of detecting neoplasms was good as high numbers of both sexes were tested. Exposure were at three concentrations for about half a life-span duration (1 year), though observations (1 year) continued to a near life-span duration. However, the highest exposure level did not reach the maximally tolerated level. Most organs were histologically examined, so most neoplasms had the ability of being detected. Although aerosol size was not ideal (slightly over the current upper limit of test guidelines), this paper did show Sb₂O₃ accumulation and decreased clearance in the lung (by 80% in the 4.5 mg/m³ group). For inert particle, such overload would/could cause lung tumor. The overload was observed at relatively low exposure concentrations (compared to inert particles, such as TiO₂) and Sb₂O₃ toxicity was suspected. It appears conditions that could potentially lead to cancer did persist (Table 9, post-exposure, chronic inflammation in most animals, although hyperplasia was in only very few animals).

Table D-12. Watt (1983) study of female rats exposed to antimony trioxide by inhalation for 1 year followed by post-exposure observation for 2 years

Utility question	Rating	Rationale
Study design		
Randomization	NR	Not reported.
Controls	+++	Concurrent controls were used, although animals were housed in different rooms (housing chambers separated to control, low concentration, and high concentration). Otherwise, treatments were the same.
Historical data		No
Animal model	++	Only female rats were used
Statistical power	+	Small number of animals were used. 13-18 animals per group sacrificed at the end of exposure. Less than 10 per group sacrificed between 2 and 12 months post exposure. Less than 20 per group sacrificed 12-months post exposure.
Exposure		
Chemical characterization	+++	Detailed chemical analysis verified that Sb ₂ O ₃ was of high purity. Small amounts of arsenic (0.02%) and lead (0.2%) were found as contaminants. Dust size (measured by SEM) was reported as Feret diameter. Presumably this is average from the same particle with rotation. Aerosol concentration in the exposure chamber. The equipment generated aerosols of MMAD less than 15 µm, but aerosol sizes were not measured. Based on conversion done in Newton et al 1994 paper Table 2, the MMAD is 5.06 µm, which is above the ideal range of rat inhalation study (no more than 4 µm).
Dosing regimen	+++	Another potential concern is the use of pine shaving in the exposure chamber. The rats were not in direct contact with shaving, but metabolism change from pine cannot be excluded. This does not affect the interpretation of this study as all groups were treated the same, but has been suggested by Newton et al 1994 as a factor affecting outcome even though it is based on concerns of increased particulates (rather than rat metabolism). Survival was not reported, but body weight gain was greater than controls, indicating the dose is not close to maximal tolerant dose. Significant increases in neoplasia occurred, indicating the dose level was high enough to cause carcinogenesis.
Exposure duration	+++	Exposure occurred for up to 1 year, with intermediate sacrifices at 3, 6, 9 months.
Dose-response	++	Only two dose levels, ranging over 2.5 folds, were used, limiting the examination of a dose response curve.
Outcome		
Pathology	++	Major organs were examined microscopically.
Consistency between groups	+++	Consistent treatment among groups, except housed in different rooms.

Utility question	Rating	Rationale
Study duration	+++	The study duration was 2 years, with 1 year of exposure and 1 year of observation.
Confounding		
Confounding	++	Animals in high dose group were heavier than low dose group at the beginning, suggesting slightly different development level. Not all organs appear to have been examined during necropsy.
Reporting and analysis		
Reporting data and statistics	++	While statistic methods were not specified, the data were reported with raw numbers and therefore enables statistical analysis
Combining lesions	+++	Tumor types were not combined. Scirrhou carcinoma, a pathologically distinctive lung cancer, alone, was significantly increased compared to negative controls.

Overall utility: ++. The chemical purity was high and exposure was characterized, though the particle size (converted by Newton et al (1994) to be around MMAD 5 μm) was over the recommended (1-4 μm). Only female rats were used, which eliminates the ability to detect sex differences. The sensitivity to detect neoplasms was low as a small number of rats were used at only two dose levels, though the exposure was near life-span duration. The ability to detect neoplasms, if they exist, was moderate as the organs examined during necropsy were not fully reported. The statistical methods used were not reported. The use of large exposure chamber with pigs inside and pine shaving also increased the chance of exposure to non-Sb₂O₃ particles (and possible metabolism alternation due to pine shaving and therefore affecting susceptibility).

Appendix E: Mechanistic and Other Relevant Information

This appendix lists Tox21/ToxCast high-throughput screening (Appendix E.1), effects of antioxidant and inhibitors of enzymes on antimony effects (Appendix E.2), genotoxicities of antimony compounds (Appendix E.3), effects related to cell proliferation (Appendix E.4), transcriptomic of antimony(III) potassium tartrate trihydrate in HepG2 cells (Appendix E.5), and immune effects of compounds containing pentavalent antimony (Appendix E.6).

E.1 Tox21/ToxCast high-throughput screening

A total of six antimony compounds, not including antimony(III) trioxide, were found in the Tox21 (Tice *et al.* 2013) and ToxCast (Kavlock and Dix 2010, Kavlock *et al.* 2012) results from the Tox21 Toolbox (NTP 2017a) and iCSS Dashboard (EPA 2017): (1) acetic acid, antimony(III) salt, (2) antimony potassium(III) tartrate trihydrate, (3) antimony(III) trichloride, (4) antimony(V) sulfide, (5) antimony(III) potassium tartrate hydrate, and (6) triphenylstibine(III).

All of the above antimony compounds except acetic acid, antimony(III) salt and antimony potassium(III) tartrate trihydrate were screened in some of the Tox21 assays, although the assays varied. Among the antimony compounds screened in Tox21, triphenylstibine(III) was also screened in ToxCast in only some of the assays in the Attagene (ATG), CeeTox, and NovaScreen (NVS) platforms. In addition, antimony(III) trichloride was also screened in the ATG platform and three estrogen receptor assays in the NVS platform in ToxCast.

The data are reviewed for antimony compounds screened in the subset of assays (Chiu *et al.* 2017, IARC 2017) that relate to the 10 key characteristics of human carcinogens (Smith *et al.* 2016). For the purpose of comparing different antimony compounds, only the responses from Tox21 assays, in which several antimony compounds were tested, were compared. The half maximal effective concentration (EC₅₀) and weighted area under the curve were obtained from the Tox21 Toolbox Activity Profiler. Assay results exhibiting the following characteristics were excluded from the analysis: observed cytotoxicity, autofluorescence, insufficient reporter gene activity readout support, suboptimal National Center for Advancing Translational Sciences fits, or substantial variation between sources. Assays that assessed only cell viability were not included. All effective EC₅₀s were within an order of magnitude. Please note that analysis via different criteria, such as dose-response fit threshold, will result in different hits, and therefore the results shown here might be different from others.

The only pentavalent antimony compound, antimony(V) sulfide, showed no activity in Tox21 assays. Antimony(III) potassium tartrate hydrate was active only in one androgen receptor antagonist assay, which was also activated by antimony(III) potassium tartrate trihydrate. Triphenylstibine was not active in any assays linked with the 10 key characteristics of carcinogens, but was active in assays associated with nuclear receptors, including constitutive androstane receptor, pregnane X receptor, and retinoic acid-related orphan receptors γ .

Antimony(III) trichloride and antimony(III) potassium tartrate trihydrate had hits in more assays than other screened antimony compounds. Observed hits by both were related to oxidative stress or antagonism of nuclear receptors, including the androgen receptor, farnesoid X receptor, and peroxisome proliferator-activated receptor delta. Antimony(III) potassium tartrate trihydrate was also active in an estrogen receptor antagonist assay. One of the common characteristics of

nuclear receptors is DNA-binding domain or zinc finger structure. Antimony(III) ions have been reported to displace Zn(II) in zinc finger domains (Nielson *et al.* 1985, Grosskopf *et al.* 2010), providing a possible link to the observed antagonist activity of nuclear receptors.

In summary, the activities of antimony compounds in Tox21 assays were mostly antagonistic to nuclear receptors, possibly because of displacement of Zn(II) in the zinc finger structures of these receptors by antimony(III) ions. These assays also indicated an oxidative stress response. Because only one antimony(V) compound was screened, and some of the trivalent compounds had very little activity in the Tox21 assays, it is unclear whether antimony(III) compounds are in general more active than antimony(V) compounds.

E.2 Effects of antioxidants and inhibitors of oxidative stress related enzymes on cells exposed to compounds containing trivalent antimony

Table E.2-1. Effects of antioxidants and inhibitors of oxidative stress related enzymes on cells exposed to compounds containing trivalent antimony

Cell types	Additional treatment (besides antimony exposure)	Oxidative stress and damage	MMP and cell death	Comparison group (cells)	Reference
Antimony (III) trioxide					
LOUCY, CCRF-CEM, HL-60, K-562	BSO, an inhibitor of γ -glutamylcysteine synthetase	↓ GSH	↓ MMP ^a ↑ cell death	exposed to Sb ₂ O ₃ alone	Lösler <i>et al.</i> 2009
HL-60, K-562	Mercaptosuccinic acid, an inhibitor of glutathione peroxidase		↑ cell death	exposed to Sb ₂ O ₃ alone	Lösler <i>et al.</i> 2009
K-562	3-amino-1,2,4-azole, an inhibitor of catalase		↑ cell death	exposed to Sb ₂ O ₃ alone	Lösler <i>et al.</i> 2009
CCRF-CEM, K-562	Sodium ascorbate, an antioxidant, but able to act as an oxidant under oxidative stress		↑ cell death	exposed to Sb ₂ O ₃ alone	Lösler <i>et al.</i> 2009
NB4	None	↑ ROS	↑ cell death	negative control	Mann <i>et al.</i> 2006
NB4-M-AsR3	None	↑ GSH	↓ cell death	parental NB4 cells	Mann <i>et al.</i> 2006
NB4	BSO, an inhibitor of γ -glutamylcysteine synthetase	↓ GSH ↑ ROS	↑ cell death	cells not treated with BSO	Mann <i>et al.</i> 2006
NB4-M-AsR3	BSO, an inhibitor of γ -glutamylcysteine synthetase	↓ GSH	↑ cell death	cells not treated with BSO	Mann <i>et al.</i> 2006
Antimony (III) trichloride					
Primary rat hepatocytes	none	↑ ROS ↑ lipid peroxidation	↓ MMP ↑ cell death		Hashemzaei <i>et al.</i> 2015
Primary rat hepatocytes	<i>n</i> -bromoheptane, a GSH-depleting agent	↓ GSH ↑ ROS	↓ MMP ↑ cell death	exposed to SbCl ₃ alone	Hashemzaei <i>et al.</i> 2015

Cell types	Additional treatment (besides antimony exposure)	Oxidative stress and damage	MMP and cell death	Comparison group (cells)	Reference
		↑ lipid peroxidation			
Primary rat hepatocytes	Dimethyl sulfoxide, a ROS scavenger	↓ ROS ↓ lipid peroxidation	↑ MMP ↑ cell death	exposed to SbCl ₃ alone	Hashemzaei <i>et al.</i> 2015
Primary rat hepatocytes	Mannitol, a ROS scavenger	↓ ROS ↓ lipid peroxidation	↑ MMP ↓ cell death	exposed to SbCl ₃ alone	Hashemzaei <i>et al.</i> 2015
Primary rat hepatocytes	Trifluoperazine, a mitochondria permeability transition pore sealing agent	↓ ROS ↓ lipid peroxidation	↑ MMP ↑ cell death	exposed to SbCl ₃ alone	Hashemzaei <i>et al.</i> 2015
Primary rat hepatocytes	Carnitine, a mitochondria permeability transition pore sealing agent	↓ ROS ↓ lipid peroxidation	↑ MMP ↓ cell death	exposed to SbCl ₃ alone	Hashemzaei <i>et al.</i> 2015
Primary rat hepatocytes	L-Glutamine, an adenosine triphosphate (ATP) generating agent	↓ ROS ↓ lipid peroxidation	↑ MMP ↑ cell death	exposed to SbCl ₃ alone	Hashemzaei <i>et al.</i> 2015
Antimony (III) potassium tartrate					
HL-60	none	↑ ROS	↓ MMP ↑ cell death	negative control	Lecureur <i>et al.</i> 2002
HL-60	BSO		↑ cell death	exposed to antimony alone	Lecureur <i>et al.</i> 2002
HL-60	N-acetylcysteine		↓ cell death	exposed to antimony alone	Lecureur <i>et al.</i> 2002

↑ = Increased.

↓ = Decreased.

NB4-M-AsR3 = Arsenic resistant subclone of parental NB4 due to increased GSH levels.

BSO = DL-buthionine-[S,R]-sulfoximine.

CCRF-CEM = a cell line derived from acute lymphoblastic leukemia cells.

HL-60 = a cell line derived from human promyelocytic leukemia.

K-562 = chronic myelogenous leukemia cells.

LOUCY = a cell line derived from T-cell acute lymphoblastic leukemia .

MMP = mitochondrial membrane potential.

NB4 = a cell line derived from human acute promyelocytic leukemia cells.

NB4-M-AsR3 cells = Arsenic-resistant APL cells (derived in Miller lab).

E.3 Genotoxicity tables

The genotoxic tables are organized by endpoints: mutations (Table E.3-1), DNA damage (Table E.3-2), chromosomal aberrations (Table E.3-3).

Table E.3-1. Genotoxicity of antimony compounds: Mutations^{a b}

Mutation studies are listed hierarchically according to the following criteria:

- 1 By genotoxicity endpoints;
- 2 By domain of target species (eukaryote and then prokaryote);
- 3 By testing system (e.g., *E. coli* strains and then *Salmonella* strains); and
- 4 By compound in the order of antimony(III) trioxide (bold) and then antimony(III) trichloride. Other forms of antimony, such as elemental antimony (Asakura *et al.* 2009) were not included in the table.

Genotoxicity endpoint	Antimony form	Testing system/exposure duration	Assay endpoint	Comments	Reference
Mammalian cells					
Point mutations and chromosome deletions	Antimony trioxide	L5178Y mouse lymphoma cell line (+/-S9, 2 experiments) 4-hour exposure duration	Negative (concentrations tested: 6–50 µg/mL)	Precipitate formed at top dose level; authors report no significant toxicity at these doses	Elliott <i>et al.</i> 1998
Bacteria					
A/T base pair substitutions	Antimony trioxide	<i>E. coli</i> B/r WP2 <i>try</i> and WP2 <i>hcr try</i> (-S9, spot test method)	Negative (concentrations tested: 0.05–0.5 M)	Microbial toxicity not reported	Kanematsu <i>et al.</i> 1980
A/T base pair substitution	Antimony trioxide	<i>E. coli</i> WP2P (+/-S9; plate incorporation and pre-incubation protocols) 3-day exposure duration	Negative (concentrations tested: 100–5000 µg/plate)	Microbial toxicity not reported	Elliott <i>et al.</i> 1998
A/T base pair substitution	Antimony trioxide	<i>E. coli</i> WP2PuvrA (+/-S9; plate incorporation and pre-incubation protocols) 3-day exposure duration	Negative (concentrations tested: 100–5000 µg/plate)		Elliott <i>et al.</i> 1998
A/T base pair substitutions	Antimony trichloride	<i>E. coli</i> B/r WP2 <i>try</i> and WP2 <i>hcr try</i> (-S9, spot test method)	Negative (concentrations tested: 0.05–0.5 M)	Microbial toxicity not reported	Kanematsu <i>et al.</i> 1980

Genotoxicity endpoint	Antimony form	Testing system/exposure duration	Assay endpoint	Comments	Reference
G/C base pair substitutions	Antimony trioxide	<i>S. typhimurium</i> TA 1535, TA 1537, TA100, TA98 (+/-S9; plate incorporation and pre-incubation protocols) 3-day exposure duration	Negative (concentrations tested: 100–5000 µg/plate)	Microbial toxicity not reported	Elliott <i>et al.</i> 1998
Frameshift mutations	Antimony trioxide	<i>S. typhimurium</i> TA 1537 and 98 (+/-S9; plate incorporation and 60 min pre-incubation protocols) 3-day exposure duration	Negative (concentrations tested: 100–5000 µg/plate)	Microbial toxicity not reported	Elliott <i>et al.</i> 1998
Base pair substitution and frameshift mutations	Antimony trioxide	<i>S. typhimurium</i> TA98, TA100, TA1535, TA1537, TA1568 (-S9, spot test method)	Negative (concentrations tested: 0.05–0.5 M)	Duration of chemical exposure for spot test assay not reported; microbial toxicity not reported	Kanematsu <i>et al.</i> 1980
Base pair substitution and frameshift mutations	Antimony trioxide	<i>S. typhimurium</i> TA100, TA98 (+/-S9; 20 min pre-incubation modification)	Negative in 3 experiments (concentrations tested: 0.43–1.71 µg/plate)	Survival after pre-incubation step reported	Kuroda <i>et al.</i> 1991
Base pair substitution and frameshift mutations	Antimony trichloride	<i>S. typhimurium</i> TA98, TA100, TA1535, TA1537, TA1568 (-S9, spot test method)	Negative (concentrations tested: 0.05–0.5 M)	Duration of chemical exposure for spot test assay not reported; microbial toxicity not reported	Kanematsu <i>et al.</i> 1980
Base pair substitution and frameshift mutations	Antimony trichloride	<i>S. typhimurium</i> TA100, TA98 (+/-S9; 20 min pre-incubation modification)	Negative in 3 experiments (concentrations tested: 625–5000 µg/plate)	Survival after pre-incubation step reported	Kuroda <i>et al.</i> 1991

^aAll data in prokaryotes were derived bacterial reverse mutation assays. The single eukaryotic study data was derived from the mouse lymphoma TK gene mutation assay.

^bLevels of significance are designated as follows: * $P < 0.05$; ** $P < 0.01$.

Table E.3-2. Genotoxic DNA damaging effects of antimony compounds

Listing order of the studies are as follows:

- 1 Assay, in the order of metaphase analysis, micronucleus assay, and sister chromatid exchange assay;
- 2 Target system, in the order of studies in human cells, animal studies, *in vitro* studies, and biochemical studies;
- 3 Compound, in the order of antimony(III) trioxide (bold), antimony(III) trichloride, and other antimony(III) compounds.

Genotoxicity endpoint	Antimony form	Assay name	Testing system	Assay endpoint ^a			Comments	References
DNA Damage (epidemiological studies)^b								
DNA strand breaks, alkali-labile sites, oxidized purines	Occupational antimony trioxide	Alkaline FPG-modified comet assay	Blood lymphocytes from occupationally exposed workers (-S9)	Frequency of subjects with oxidative DNA damage			Sb ₂ O ₃ levels for direct and indirect exposure groups lower than OSHA/NIOSH PEL and REL for workplace. Moderate oxidative DNA damage observed in direct exposure group (0.12 ± 0.11 µg/m ³); potential concomitant exposures not addressed.	Cavallo <i>et al.</i> 2002
				Conc. (µg/m³)		# with oxidative damage/total		
				0		3/23		
				0.120 ± 0.110		11/17		
				0.052 ± 0.038		1/6		
				Relative risk of DNA damage				
				Conc. (µg/m³)	Adjusted relative risk	95% CI		
				0	1	n/a		
				0.120 ± 0.110	14.2**	2.7–73.4		
				0.052 ± 0.038	1.7	0.1–22.5		
				Tail moment values for FPG-treated Cells				
				Conc. (µg/m³)		Mean ± SD		
				0		24.4 ± 9.51		
				0.120 ± 0.110		32.4 ± 16.3		
				0.052 ± 0.038		28.8 ± 5.61		
Tail moment values for untreated cells								
Conc. (µg/m³)		Mean ± SD						
0		16.3 ± 6.59						
0.120 ± 0.110		14.6 ± 8.29						

Genotoxicity endpoint	Antimony form	Assay name	Testing system	Assay endpoint ^a		Comments	References	
				0.052 ± 0.038	18.3 ± 8.78			
DNA strand breaks, alkali-labile sites, oxidized purines	Occupational antimony trioxide	AP sites quantified using ELISA technique	Blood lymphocytes from occupationally exposed workers (-S9)	<p>The quantity of DNA damage (determined by the number of AP sites/1×10^5 bp) among the studied workers was significantly ($p = 0.004$) higher compared to that recorded for the control group and a significant positive correlation was found between the quantity of DNA damage (in the form of increased AP sites) and urinary antimony level among workers ($r = 0.873$, $P < 0.001$). Total oxidative capacity (also measured by ELISA) was not different between workers and controls.</p>		The number of measured abasic sites ranged from 17.22 (control group) to 26.88 (exposed workers)/ 1×10^5 bp. This range is higher than expected.	El Shanawany <i>et al.</i> 2017	
DNA damage (<i>in vitro</i> studies in human cells)								
DNA strand breaks, alkali-labile sites, DNA-protein crosslinks	Antimony trichloride (concentrations tested: 1–50 μ M)	Alkaline comet assay +/- proteinase K	Human whole blood or human lymphocytes exposed ex vivo (-S9)	Mean tail moment in human whole blood in comet assay without proteinase K			Significance tested by Kruskal-Wallis one-way ANOVA on ranks.	Schaumlöffel and Gebel 1998
				Conc. (μM)	Time. (hrs)	Mean \pm SD		
				0	2.5	1.28 \pm 0.10		
				1	2.5	1.26 \pm 0.01		
				5	2.5	1.32 \pm 0.08		
				10	2.5	1.32 \pm 0.04		
				25	2.5	1.47 \pm 0.07		
				50	2.5	1.75 \pm 0.08*		
				Mean tail moment in human lymphocytes in comet assay without proteinase K				
				Conc. (μM)	Time (hrs)	Mean \pm SD		
0	2.5	1.00 \pm 0.02						
1	2.5	1.23 \pm 0.28						
5	2.5	1.39 \pm 0.19*						

Genotoxicity endpoint	Antimony form	Assay name	Testing system	Assay endpoint ^a			Comments	References
				10	2.5	1.56 ± 0.04*		
				25	2.5	1.64 ± 0.03***		
				50	2.5	2.14 ± 0.01***		
				Mean tail moment in human lymphocytes in comet assay with proteinase K				
				Conc. (µM)	Time (hrs)	Mean ± SD		
				0	2.5	1.08 ± 0.11		
				1	2.5	1.13 ± 0.09		
				5	2.5	1.30 ± 0.20		
				10	2.5	1.47 ± 0.13*		
				25	2.5	1.53 ± 0.08*		
				50	2.5	1.94 ± 0.30***		
DNA damage (animal studies)								
DNA strand breaks and alkali labile sites	Antimony trioxide NC: air	<i>In vivo</i> exposure (inhalation) Alkaline comet assay		Percent tail DNA			Trend tests show significant increase for both lung tissue of males and females exposed to trioxide; No increase in percent tail DNA observed in leukocytes of males or females exposed to trioxide. Normally distributed data analyzed by independent sample's t-test and linear regression; data that were not normally distributed were analyzed by the Mann-Whitney test followed	NTP 2017b
				Dose (mg/m³)	Time (mo.)	Mean ± SE		
				0	12	25.6 ± 0.78		
				3	12	33.7 ± 2.62*		
				10	12	33.5 ± 2.02**		
		30	12	37.5 ± 2.28***				
		Lung of female mice exposed via inhalation for 12 months	Percent tail DNA					
			Dose (mg/m³)	Time (mo.)	Mean ± SE			
			0	12	32.8 ± 1.11			
			3	12	35.8 ± 2.09			
10	12		36.4 ± 2.65					
30	12	45.5 ± 2.32***						

Genotoxicity endpoint	Antimony form	Assay name	Testing system	Assay endpoint ^a		Comments	References
						by the Kendall rank correlation test	
DNA strand breaks and alkali labile sites	Antimony trioxide NC: air	<i>In vivo</i> exposure (inhalation) Alkaline comet assay	Lung and blood leukocytes of male and female rats exposed via inhalation for 12 months	No statistically significant increases were observed in percent tail DNA in blood leukocytes or lung tissue in exposed rats of either sex		Normally distributed data analyzed by independent sample's t-test; data that were not normally distributed were analyzed by the Mann-Whitney test followed by the Kendall rank correlation test	NTP 2017b
DNA damage (<i>in vitro</i> studies in non-human mammalian cells)							
DNA strand breaks and alkali labile sites	Antimony trichloride	Alkaline comet assay	V79 Chinese hamster cells exposed <i>in vitro</i> (-S9)	Tail moment was significantly elevated at a minimum dose of 1 µM Sb(III); no difference could be found comparing the results obtained in presence and absence of proteinase K.		DNA damage observed below cytotoxic levels; antimony uptake measured	Gebel <i>et al.</i> 1998
DNA damage (bacterial systems)							
Growth in recombination-repair deficient bacterial strain	Antimony trioxide NC: Kanamycin (5, 10 20 µg/plate) PC: Mitomycin C (0.05, 0.1, and 0.2 µg/plate)	B. subtilis rec assay	B. subtilis M45(rec-) and H17(rec+)	H117 (Rec+) and M45 (Rec-) inhibition length		Used spore plate method	Kuroda <i>et al.</i> 1991
				Conc. (µg/plate)	Difference in Inhibition length (mm)		
				NC (5)	0		
				NC (10)	0		
				NC (20)	0.5		
				PC (0.05)	8.0		
				PC (0.1)	8.0		
				PC (0.2)	7.0		
				0.3	2.5		
				0.6	4.0		
1.1	4.5						

Genotoxicity endpoint	Antimony form	Assay name	Testing system	Assay endpoint ^a		Comments	References
Growth in recombination-repair deficient bacterial strain	Antimony trioxide PC and NC: other metals tested	<i>B. subtilis</i> rec assay	<i>B. subtilis</i> M45(rec-) and H17(rec+) (-S9)	HI17 (Rec+) and M45 (Rec-) inhibition length		Examined 127 metals; Used streak plate method; Included cold incubation step to increase contact of metal with bacteria	Kanematsu <i>et al.</i> 1980
				Solution conc. (M)	Difference in inhibition length (mm)		
				0.05	5		
Growth in recombination-repair deficient bacterial strain	Antimony trichloride NC: Kanamycin (5, 10 20 µg/plate) PC: Mitomycin C (0.05, 0.1, and 0.2 µg/plate)	<i>B. subtilis</i> rec assay	<i>B. subtilis</i> M45(rec-) and H17(rec+) (-S9)	HI17 (Rec+) and M45 (Rec-) inhibition length		Used spore plate method	Kuroda <i>et al.</i> 1991
				Conc. (µg /plate)	Difference in inhibition length (mm)		
				NC (5)	0		
				NC (10)	0		
				NC (20)	0.5		
				PC (0.05)	8.0		
				PC (0.1)	8.0		
				PC (0.2)	7.0		
				6.3	1.5		
				12.5	4.5		
23	4.5						
Growth in recombination-repair deficient bacterial strain	Antimony trichloride	<i>B. subtilis</i> rec assay	<i>B. subtilis</i> M45(rec-) and H17(rec+) (-S9)	Antimony trichloride result was negative in rec assay (tested at 0.05M)		Antimony pentachloride also negative	Nishioka 1975
Growth in recombination-repair deficient bacterial strain	Antimony trichloride PC and NC: other metals tested	<i>B. subtilis</i> rec assay	<i>B. subtilis</i> M45(rec-) and H17(rec+) (-S9)	HI17 (Rec+) and M45 (Rec-) inhibition length		Examined 127 metals; Used streak plate method; Included cold incubation step to increase contact of metal with bacteria	Kanematsu <i>et al.</i> 1980
				Solution Conc. (M)	Difference in inhibition length (mm)		
				0.01	7		

Genotoxicity endpoint	Antimony form	Assay name	Testing system	Assay endpoint ^a	Comments	References			
Induction of recombination-repair genes	Antimony trichloride	SOS chromotest for genotoxicity	<i>E. coli</i> PQ37 derived from strain GC4436 (-S9)	SOS chromotest was negative for antimony trichloride (concentration tested: 11–707 μ M)	Cytotoxicity observed at 354 μ M	Lantzsch and Gebel 1997			
Induction of recombination-repair genes	Antimony trichloride	Umu test for genotoxicity	<i>S. typhimurium</i> TA1535/pSK1002 (-S9)	Umu test was negative for antimony trichloride (concentrations tested: 1.6–820 μ M)	Data not reported	Yamamoto <i>et al.</i> 2002			
DNA Damage (biochemical assay)									
plasmid DNA nicking	Trimethylstibine	Plasmid DNA nicking assay	Plasmid pBR322 exposed <i>in vitro</i> (gaseous phase) to test reactions for 30 min.	Estimated Quantity of Open Circular form of Plasmid^d		Chemical reactions to produce trimethylstibine were conducted in situ; Plus and minus designations were estimated from images only (no quantitation of nicked and supercoiled forms). Negative results were reported for potassium antimony tartrate.	Andrewes <i>et al.</i> 2004		
	Potassium antimony tartrate PC: Trimethylarsine			Dose (μM)				Result	
				Trimethylstibine	NC			+/-	
					5			+/-	
					20			+/-	
					50			+	
					200			++	
500	+++								
5000	+++								

AP = apurinic/aprimidinic; avg = average; CI = confidence interval; conc. = concentration; ELISA = enzyme-linked immunosorbent assay; FPG = formamidopyrimidine-DNA glycosylase; hr =hour(s); mo = month(s); NC = negative control; NR=not reported; PC = positive control; SD = standard deviation; SE = standard error; VC = vehicle control.

^aLevels of significance are designated as follows: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

^bDNA damage estimated as quantity of open circular (vs supercoiled) forms from images of plasmids electrophoretically separated in ethidium bromide-stained agarose gels.

Table E.3-3. Genotoxicity of antimony compounds – chromosomal aberrations, micronucleus, and sister chromatic exchange^{a, b, c, d}

Studies are listed hierarchically according to the following criteria:

- 1 Assay, in the order of assays for chromosomal aberrations, micronucleus, and sister chromatid exchange.
- 2 Target system, in the order of studies in human cells, animal studies, *in vitro* studies, biochemical studies.
- 3 Compound, in the order of antimony trioxide (bold), antimony trichloride, other antimony(III) compounds.

Substance	Exposure and assay name	Testing system and exposure duration	Assay endpoint	Comments	References			
Chromosomal aberrations								
Antimony trioxide NC: dimethyl sulfoxide (10 µL/mL) PC: mitomycin C (0.2 µg/mL for-S9) or cyclo-phosphamide (50 µg/mL for +S9)	<i>In vitro</i> exposure Metaphase analysis	Human peripheral lymphocytes with 2 hr exposure to colcemid (-S9) Exposure time: 20 hr and 44 hr Dose: 10, 50, 100 µg/mL	Mean % aberrant cells excluding gaps			Precipitate formed at top dose level	Elliott <i>et al.</i> 1998	
			Group	HIC/LEC (µg/mL, unless specified)	Mean (%)			
			NC	–	0.5–1.5			
			PC	–	22.0–32.0**			
			Donor 1, 20 hr	100	2.0			
			Donor 2, 20 hr	100	12.5**			
		Human peripheral lymphocytes with 2 hr exposure to colcemid (+S9) Dose: Same as above	NC	–	1.0–1.5			
			PC	–	26–34.0**			
			Donor 1, 20 hr	50	4.5*			
			Donor 2, 20 hr	100	9.5**			
			Donor 2, 44 hr	100	2.0			
Antimony sodium tartrate	<i>In vitro</i> exposure Metaphase analysis	Human leucocytes Exposure time: 48 hr Concentration: 2.3 nM	12% of cells with chromatid breaks ($P < 0.05$)	Purity of test compound not reported; toxicity (marked reduction in mitotic index) reported at 10 nM	Paton and Allison 1972			

Substance	Exposure and assay name	Testing system and exposure duration	Assay endpoint			Comments	References
Antimony trioxide VC: HPMC/poly-sorbate PC: Cyclo-phosphamide	<i>In vivo</i> exposure <i>Ex vivo</i> metaphase analysis	Sprague-Dawley rat bone marrow cells (-S9) Exposure time: Once daily for 21 consecutive days by oral gavage (except PC administered on only on day 21) Dose: 250, 500, 1000 mg/kg	Frequency of cells with chromosomal aberration excluding gaps in male rats			Body-weight gain was reduced (<10%) in the top dose group of treated rats of both sexes over the 3-week dosing period.	Kirkland <i>et al.</i> 2007
			Group	HIC/LEC (mg/kg)	Mean% ± SD		
			VC	20	0 ± 0		
			PC	20	13 ± 6.63***		
			Male rat	1000	0 ± 0		
		Female rat	1000	0 ± 0			
Antimony trioxide NC: distilled water	<i>In vivo</i> exposure <i>Ex vivo</i> metaphase analysis	Male Swiss albino mice bone marrow cells (-S9) Exposure by daily oral gavage on days 7, 14 and 21. Dose: 400, 666.7, 1000 mg/kg	Frequency of aberrations excluding gap			Purity of test compound not reported; Test for trend significant for 7 and 14 days for analysis including and excluding gaps (not shown in this table). No increases in chromosomal aberrations was observed after single acute exposure at same doses and measured 6, 12, 18 and 24 hours); Highest dose was lethal.	Gurnani <i>et al.</i> 1992b
			LEC (mg/kg)	Time (days)	Mean % ± SD		
			NC	7	1.4 ± 1.140		
			400	7	2.2 ± 0.447*		
			NC	14	1.6 ± 0.547		
			400	14	3.2 ± 0.447*		
			NC	21	1.6 ± 0.547		
400	21	4.6 ± 0.547*					
Antimony trichloride NC: distilled water	<i>In vivo</i> exposure <i>Ex vivo</i> metaphase analysis	Female Swiss albino mice bone marrow cells (-S9) Dose: 70, 140, 233.3 mg/kg	Frequency of aberrations including gap			Source and purity of test compound not reported Test for trend significant for 6, 12, 18, and 24 hr analysis including and excluding gaps (not shown in this table).	Gurnani <i>et al.</i> 1992a
			LEC (mg/kg)	Time (hrs)	Mean% ± SD		
			NC	6	1.6 ± 0.547		
			70	6	2.6 ± 0.547		
			NC	12	1.0 ± 1.0		
		70	12	3.0 ± 0.0			

Substance	Exposure and assay name	Testing system and exposure duration	Assay endpoint			Comments	References
		Single exposure by oral gavage analyzed at 6, 12, 18 and 24 hrs	NC	18	1.6 ± 0.547		
			70	18	3.2 ± 0.836		
			NC	24	1.0 ± 0.0		
			70	24	4.2 ± 1.095		
Potassium antimony tartrate Control: untreated animals	<i>In vivo</i> exposure <i>Ex vivo</i> metaphase analysis	Male rats bone marrow (-S9) Exposure via single intraperitoneal injection at each dose; Also, tested repeated exposure (daily for 5 days) at each dose. Dose: 2.0, 8.4, 14.8 mg/kg	Metaphases with aberrations excluding gap			Similar findings for aberrations including gaps but statistical analysis not performed	El Nahas <i>et al.</i> 1982
			LEC (mg/kg, unless specified)	Time after treatment (hr, unless specified)	%		
			NC	n/a	0.7		
			2.0	6	2.0*		
			2.0	24	2.4*		
			8.4	48	5.2*		
			2.0 mg/kg/day x 5 days	-	7.6*		
Micronuclei							
Occupational antimony trioxide	Epidemiology study Sister chromatid exchange assay	Blood lymphocytes from textile workers exposed to low levels of antimony trioxide 23 exposed workers: 17 high exposure (0.12 ± 11 µg/m ³) and 6 lower exposure (0.052 ± 0.038 µg/m ³) 23 controls	Mean micronuclei/1000 binucleated cells did not differ between controls and two exposure groups			High exposure well below OHSA permissible exposure levels and NIOSH recommended exposure levels Exposure groups had similar ages, and smoking habits	Cavallo <i>et al.</i> 2002
Antimony trichloride NC: DMSO PC:	<i>In vitro</i> exposure Micronucleus test	Human peripheral lymphocytes (-S9) Doses: 0, 0.5, 2, 5, 25 µM	Induction of micronuclei by Sb(III)			Co-incubation with SOD or CAT had no effect on micronucleus frequency; Statistical	Schaumlöffel and Gebel 1998
			LEC (µM)	Time (hrs)	MN/1000 BN, mean ± SD		
			0	20	10 ± 1.4		

Substance	Exposure and assay name	Testing system and exposure duration	Assay endpoint			Comments	References
mitomycin C (data not shown)			5	20	30.5 ± 2.1	significant in MN observed in second experiment at 5, 10 and 25µM	
Antimony trioxide NC: air	<i>In vivo</i> exposure <i>Ex vivo</i> micronucleus test	Male mice peripheral blood erythrocytes exposed via inhalation for 12 months. Dose: 0, 3, 10, 30 mg/m ³	No significant increase in micronucleated PCEs/1,000 PCEs in male mice			Twenty thousand CD71+ reticulocytes (PCE) were scored per animal for the presence of micronuclei and 1 × 10 ⁶ erythrocytes (NCE) were counted for micronuclei. William's and Dunn's test were used for pairwise significance, and Jonckheere's test and linear regression used for trend significance. MN frequency in NCEs but not PCEs significant by trend test (<i>P</i> < 0.001) in both sexes.	NTP 2017b
			Micronucleated NCEs/1,000 NCEs				
			LEC (mg/m ³)	Time (mo.)	Mean ± SE		
		30	12	1.93 ± 0.10***			
		Female mice peripheral blood erythrocytes exposed via inhalation for 12 months Dose: 0, 3, 10, 30 mg/m ³	No significant increase in micronucleated PCEs/1,000 PCEs in female mice				
			Micronucleated NCEs/1,000 NCEs				
LEC (mg/m ³)	Time (mo.)		Mean ± SE				
30	12	1.38 ± 0.09***					
Antimony trioxide NC: air	<i>In vivo</i> exposure <i>Ex vivo</i> micronucleus test	Male rat peripheral blood erythrocytes exposed via inhalation for 12 months	No significant increase in micronucleated PCEs/1,000 PCEs or micronucleated NCEs/1000 NCEs in male rats.			Twenty thousand CD71+ reticulocytes (PCE) were scored per animal for the presence of micronuclei and 1 × 10 ⁶ erythrocytes (NCE) were counted for micronuclei. William's and Dunn's test were used for pairwise significance, and	NTP 2017b
		Female rat peripheral blood erythrocytes exposed via inhalation for 12 months	No significant increase in micronucleated PCEs/1,000 PCEs or micronucleated NCEs/1000 NCEs in female rats.				

Substance	Exposure and assay name	Testing system and exposure duration	Assay endpoint	Comments	References
				Jonckheere's test and linear regression used for trend significance. No significant changes were observed in MN frequency in rats of either sex.	
Antimony trichloride	<i>In vitro</i> exposure Micronucleus test	Human fibroblast cells (-S9) Human bronchial epithelial cells (BES-6) (-S9) Chinese hamster ovary cells (CHO-K1) (-S9) Exposure time: 4 hr, Dose: 50–400 μ M	Positive findings for all cell types at all doses	LD ₅₀ = 40 μ M in fibroblast cells LD ₅₀ = 80 μ M in BES-6 cells LD ₅₀ = 180 μ M in CHO-K1 cells	Huang <i>et al.</i> 1998
Antimony trioxide VC: DMSO PC: Cyclophosphamide (20 mg/kg)	<i>In vivo</i> exposure Micronucleus test	Mouse bone marrow (-S9) male and females Single dose study Exposure time: 24 and 48 hr Dose: 5000 mg/kg by oral gavage Repeated dose study: Exposure time: 8, 15 and 22 days Dose: 400, 667, or 1000 mg/kg by oral gavage	No increases in mean incidence of MPE/1000 PE in the single dose study (males and females) or in the repeated dose study (sex not identified).	Significantly decreased frequency of polychromatic erythrocytes observed in females at 24 hr in the single dose experiment.	Elliott <i>et al.</i> 1998
Antimony trioxide	<i>In vivo</i> exposure Micronucleus test	Sprague-Dawley male and female rat bone marrow cells (-S9)	No increase in the frequency of micronucleated PCE in male and female rats		Kirkland <i>et al.</i> 2007

Substance	Exposure and assay name	Testing system and exposure duration	Assay endpoint			Comments	References
VC: HPMC/poly-sorbate PC: Cyclophosphamide (20 mg/kg)		Exposure time: 21 days (except for PCs) by oral gavage Dose: 250, 500, 1000 mg/kg					
Antimony trioxide VC: DMSO (25 µL) PC: Mitomycin C (0.5 µM)	<i>In vitro</i> Micronucleus test with cytokinesis block	Chinese hamster V79 cells Exposure time: 24 hr Dose: 2–50 µM	Mean number of micronuclei			Study measured both antimony uptake in cells and cytotoxicity (50% neutral red uptake was found with SbCl ₃ at 83 µM)	Gebel <i>et al.</i> 1998
			Group	LEC (µM)	Mean		
			VC	–	9.5		
			PC	–	45.5		
			Antimony trioxide	25	17.5		
Sister chromatid exchange							
Occupational antimony trioxide	Epidemiology study Sister chromatid exchange assay	Peripheral blood lymphocytes from textile workers exposed to low levels of antimony trioxide 23 exposed workers: 17 high exposure (0.12 ± 11 µg/m ³) and 6 lower exposure (0.052 ± 0.038 µg/m ³) 23 controls	Mean SCE did not differ between controls and two exposure groups			High exposure well below OHSA permissible exposure levels and NIOSH recommended exposure levels Exposure groups had similar ages, and smoking habits	Cavallo <i>et al.</i> 2002
Antimony trioxide (dissolved in distilled water)	<i>In vitro</i> exposure Sister chromatid exchange assay	Human peripheral blood lymphocytes from healthy non-smokers aged 25–Human 35 years (-S9)	SCE/cell		NC was DMSO, and it is unclear whether the 0 µM result was from distilled water or DMSO. No PC was	Gebel <i>et al.</i> 1997	
			LEC (µM)	Mean ± SD			
			0	8.6 ± 3.4			
		0.5	11.5 ± 4.4*				

Substance	Exposure and assay name	Testing system and exposure duration	Assay endpoint			Comments	References	
NC: DMSO		Exposure time: 24 hrs				stated in the study. Results are from 60 metaphase cells scored on two slides.		
Antimony trichloride (dissolved in DMSO) NC: DMSO	<i>In vitro</i> exposure Sister chromatid exchange assay	Human peripheral blood lymphocytes from healthy non-smokers aged 25–35 years (-S9) Exposure time: 24 hr	SCE/cell			No PC was stated in the study. Results are from 60 metaphases scored on two slides. NC was DMSO, and it is unclear whether the 0 µM result was from distilled water or DMSO.	Gebel <i>et al.</i> 1997	
			LEC (µM)		Mean ± SD			
			0		8.8 ± 4.0			
			1		13.8 ± 5.5**			
Antimony trioxide NC: Water (100 µL) PC: Mitomycin C (0.01 µg/mL)	<i>In vitro</i> exposure Sister chromatid exchange assay	Chinese hamster V79 cells Exposure time: 28 hr Dose: 0.09–0.34 µg/mL	Frequency of sister chromatid exchanges/metaphase			Sb ₂ O ₅ was negative in the SCE assay; Similar results in experiment 1, although LEC was 0.17 µg/mL	Kuroda <i>et al.</i> 1991	
			LEC (µg/mL)	Time (hrs)	Mean ± SD			
			NC	28	6.3 ± 2.5			
			PC	28	56.0 ± 9.3**			
			0.09	28	10.6 ± 3.7**			
Antimony trichloride NC: Water (100 µL) PC: Mitomycin C (0.01 µg/mL)	<i>In vitro</i> exposure Sister chromatid exchange assay	Chinese hamster V79 cells Exposure time: 28 hr Dose: 1.3–20 µg/mL	Frequency of sister chromatid exchanges/metaphase			SbCl ₅ was negative in the SCE assay. Toxic at 20 µg/mL; similar results in experiment 2, although LEC was 5 µg/mL.	Kuroda <i>et al.</i> 1991	
			Conc. (µg/mL)	Time (hrs)	Mean ± SD			
			NC	28	4.5 ± 2.2			
			PC	28	46.8 ± 8.6**			
			2.5	28	7.5 ± 4.3*			

b.w. = body weight; FISH = fluorescence in situ hybridization; HIC = Highest ineffective concentration; hr = hour(s); LEC = Lowest effective concentration; mo = months; NC = Negative control; NR = not reported; PC = Positive Control; VC = Vehicle Control.

* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

^a Provided are the form of the test compound, study details including the testing system and exposure duration, assay endpoint results for test compounds and positive and negative controls, comments provided by reviewers, and reference.

^b Abbreviations used in this table are as follows:

^c Levels of significance are designated as follows:

^d Compounds containing pentavalent antimony are not included. For instance, trimethylantimony dichloride in Dopp *et al.* 2006 (no increase of MN formation, chromosome aberration, or sister chromatid exchange in the Chinese hamster ovary cells after exposure to at up to 1 mM. When the cells underwent electroporation to double the intake of trimethylantimony dichloride, the formation of MN was increased.) and KSbO₃ in Migliore *et al.* 1999 (non significant increase of centromere-negative MN) were not included in the table.

E.4 Studies related to cell proliferation

Table E.4-1. Mutations in the lung of mice and rats after two-year inhalation exposure to antimony trioxide (NTP 2017b).

Genotoxicity endpoint	Testing system	Assay endpoint		Comments	Reference
<i>Egfr</i> mutations	Lung tumors from exposed B6C3F1/N mice. Both nontumor lung and spontaneous tumors from control mice.	Mutation Frequency			NTP 2017b
		Concentration (mg/m³)	# with mutation/# tissues assayed		
		0 (nontumor lung)	0/10		
		0 (tumor lung)	0/9		
		3 (tumor lung)	11/28*		
		10 (tumor lung)	11/26*		
30 (tumor lung)	15/26**				
<i>Egfr</i> mutations	Lung tumors from exposed Wistar Han rats. Both nontumor lung and spontaneous tumors from control mice.	Mutation Frequency		Increase was <u>not</u> statistically significant.	NTP 2017b
		Concentration (mg/m³)	# with mutation/# tissues assayed		
		0 (nontumor lung)	0/11		
		0 (tumor lung)	0/4		
		3 (tumor lung)	3/5		
		10 (tumor lung)	6/11		
30 (tumor lung)	4/10				
<i>Kras</i> mutations	Lung tumors from exposed Wistar Han rats. Both nontumor lung and spontaneous tumors from control mice.	Mutation Frequency		Increase was <u>not</u> statistically significant.	NTP 2017b
		Concentration (mg/m³)	# with mutation/# tissues assayed		
		0 (nontumor lung)	0/11		
		0 (tumor lung)	0/4		
		3 (tumor lung)	0/5		
		10 (tumor lung)	1/11		
30 (tumor lung)	0/10				
<i>Kras</i> mutations	Lung tumors from exposed B6C3F1/N mice. Both nontumor lung and spontaneous tumors from control mice.	Mutation Frequency		Increase was <u>not</u> statistically significant.	NTP 2017b
		Concentration (mg/m³)	# with mutation/# tissues assayed		
		0 (nontumor lung)	0/10		
		0 (tumor lung)	3/9		
		3 (tumor lung)	9/28		
		10 (tumor lung)	15/26		
30 (tumor lung)	10/26				

E.5 Transcriptomic of antimony(III) potassium tartrate trihydrate in HepG2 cells

One DNA microarray study (Kawata *et al.* 2007) of *in vitro* effects of an antimony(III) compound on a human cell line was found in the National Center for Biotechnology Information Gene Expression Omnibus (NCBI GEO) database (NCBI 2017). HepG2 (human liver carcinoma) cells were exposed to bis[(+)-tartato]diantimonate(III) dipotassium trihydrate (i.e., antimony(III) potassium tartrate trihydrate, equivalent to one molecule of antimony(III) potassium tartrate plus three water molecules) at a concentration of 200 μ M for 6 hours, and the gene expression changes seen in a Human Genome Focus array (Affymetrix) were compared with changes following exposure to five other substances, including arsenic(III) oxide at 20 μ M and nickel(III) chloride hexahydrate at 6.5 nM. The gene expression profile after antimony(III) potassium tartrate trihydrate exposure was most similar to that after nickel(III) chloride hexahydrate exposure.

The microarray data were downloaded from the NCBI GEO database and analyzed in Ingenuity Pathway Analysis (Qiagen) by the NTP ORoC, using the filter of minimal 2-fold change. Of the top ten canonical pathways affected (Table E.5-1), seven were related to immune reactions (pathways 1, 2, 4, 5, 7, 8, and 9). These findings are consistent with the former use of antimony(III) potassium tartrate as an antiparasitic agent for leishmaniasis. The other three pathways were eicosanoid signaling, bladder-cancer signaling, and detoxification of oxidized guanosine triphosphate (GTP) and deoxyguanosine triphosphate (dGTP). Although antimony is not known to cause urinary-bladder cancer, the chemically similar arsenic increases the incidence of transitional-cell carcinoma of the urinary bladder in humans. An effect on the oxidized GTP and dGTP detoxification pathway is consistent with the observation that various antimony compounds increase oxidative stress (as discussed in Section 6.2).

In the upstream analysis, the top three affected regulators were vascular endothelial growth factor (VEGF), colony-stimulating factor 2 (CSF2) (a cytokine), and the triggering receptor expressed on myeloid cells 1 (TREM1), which stimulates neutrophil- and monocyte-mediated inflammatory responses (Appendix E.5, Table E.5-1). In a 2015 study, antimony(III) potassium tartrate inhibited the VEGF-induced formation of capillary-like structures in endothelial cells (Wang *et al.* 2015). In other words, antimony(III) potassium tartrate showed anti-tumor effects via anti-angiogenesis in cultured cells. Both CSF2 and TREM1 stimulate immune or inflammatory responses. These top three affected regulators are predominantly involved in skin disease and cancer. Some anti-cancer effects, such as increased differentiation of cells, were also enriched in the gene expression. To identify key factors contributing to potential carcinogenic effects, further analysis is needed. It is also possible that 6-hour exposure leads to mostly acute responses, which may differ from the long-term effects.

Table E.5-1. Top 10 upstream regulators for antimony

Upstream Regulator	Expr Fold Change	Molecule Type	Predicted Activation State	Activation z-score	Flags	p-value of overlap	Target molecules in dataset	Mechanistic Network
1 Vegf		group	Activated	9.487	bias	1.88E-09	ANGPT2,ANGPTL4,AQP4,ATF3,AURKA,AURKB,BCL2A1,BIRC5,BNC1,BTN1A1,CA2,CALB1,CALCRL,CCL7,CCNF,CD3EAP,CDC14A,CDC20,CDC25A,CDC25B,CDC25C,CDC45,CDH5,CDK1,CDKN2C,CDKN3,CELSR1,CHI3L1,CHIA,CHRN2,CHST7,CKS1B,CLCF1,CNN1,CNTFR,CPA3,CRLF1,CRYAB,CSF2,CXCL1,CXCL8,CXCR2,CXCR4,CYR61,DBF4,DPF3,DRD3,DTYMK,DUSP4,DUSP5,EDN1,EGR1,EGR3,EMCN,EMP2,ESM1,FABP4,FAIM2,FANCG,FGF16,FGF2,FLNA,FOSB,FOSL1,GATA1,GEM,GH1,GPR4,GPRC5B,HBE1,HBEGF,HDC,HMOX1,HOXB8,HPSE,HTR7,IL18,IL1A,IL3RA,IL4,ITGB3BP,JAM2,JUN,KIF15,KIF22,KIF2C,KITLG,LEF1,LPAR1,LRAT,LYVE1,MCM2,MCM5,MID1,MKI67,MMP10,MMP14,MT1G,MYCN,NDC80,NEK2,NFATC1,NGB,NR4A2,NR4A3,NRCAM,NRG1,PLK1,PLXNA2,PMAIP1,PRC1,PRKCB,PSMC3IP,PTH,RGS2,RGS20,SOCS2,SOCS3,ST8SIA4,STK10,TAAR5,TACR1,TACSTD2,TBXA2R,THBD,TNC,TNFRSF9,TNFSF15,TPX2,TRAF5,TRAIP,TRPC4,TTK,UBE2C,XCR1	
2 CSF2	8.025	cytokine	Activated	8.308	bias	1.85E-08	ADA,ADAM8,ADGRE5,ANXA1,AURKA,BCL2A1,BIRC5,C5AR1,CCL4,CCNF,CCR1,CCR5,CCR7,CD1C,CD209,CD28,CD33,CD40LG,CD69,CD8A,CDC20,CDK1,CDKN1A,CDKN2B,CDKN2C,CENPE,CHAF1A,CHAF1B,CKS1B,CLCF1,COL8A1,CSF1,CSF2,CTLA4,CXCL1,CXCL2,CXCL8,CXCR4,CYBB,EDN1,EGR1,EGR2,EGR3,EPOR,EXO1,FANCA,FCGR2B,FOLR2,FOS,FOSL1,FPR2,GATA1,GCLM,GDF15,HBEGF,HDC,HLA-DQB1,HRH4,HRK,HSPH1,IER3,IFNG,IGF1,IL1A,IL1RN,IL24,IL3RA,IL4,ITGA4,ITGAM,LEP,MCM5,MKI67,MMP1,MMP14,MRC1,NEK2,NFATC1,NFE2,NFKBIA,NR4A2,NUSAP1,OSM,PDE1B,PIM1,PLK1,POLD1,POLE,PPP1R15A,PRC1,PTGER2,RARA,RECL4,RELB,RRM2,SERPIN9,SLC1A5,SOCS2,SOCS3,SPAG5,SPI1,STMN1,THBS1,TLR2,TLR4,TNFAIP3,TNFRSF1B,TN	352 (5)

Upstream Regulator	Expr Fold Change	Molecule Type	Predicted Activation State	Activation z-score	Flags	p-value of overlap	Target molecules in dataset	Mechanistic Network
							FRSF9,TNFSF14,TNFSF15,TNFSF8,TPM4,TPX2,UBE2C,ZFP36	
3 TREM1	1.62	transmembrane receptor	Activated	4.945	bias	0.000000 203	ATF3,CASP5,CCL7,CCR7,CCRL2,CDK1,CDKN2B,CEBPB,CKS2,CSF1,CSF2,CXCL1,CXCL2,CXCL3,CXCL5,CXCL8,CXCR4,DCSTAMP,DEFB4A/DEFB4B,DUSP4,EDN1,EGR1,EGR2,EGR3,FOSL1,GADD45B,GCLM,GEM,GIPR,GLA,HAS1,HBEGF,IFNG,IL17A,IL36RN,IL4,LPL,MAD1L1,MAFF,MMP1,MMP10,MMP19,NFKBIA,NOD2,NR4A2,OSGIN1,RGS1,RRAD,SLC1A3,SNAPC1,TCEAL9,THBD,THBS1,TLR2,TLR4,TNFSF14,TNFSF15,WNT5A	
4 GATA2	2.854	transcription regulator		1.922		0.000000 237	ADGRE5,ANGPT2,ANGPTL4,ARPP21,C9,CCL21,CCR8,CD177,CD34,CD36,CD69,CD96,CDH5,CDK6,CDKN1A,CEL,CELSR3,CHGA,CHI3L1,CLDN18,CMA1,CPA1,CPA3,CST7,CYBB,CYP2F1,CYP4F11,DDX4,DLK1,E2F2,EDN1,ELANE,EMCN,EPHA3,FABP4,FCN1,GABRP,GATA1,GATA2,GP5,GP9,GPR65,GUCA2A,HBQ1,HDC,HOXA10,HSD17B1,ICAM2,IKZF1,IL3RA,IL4,IL4R,ITGAM,KLF2,KLK3,LYL1,MAFB,MEP1A,MMRN1,MPIG6B,NFE2,PAX3,PDE9A,PLK2,PRG3,RAG1,REG1A,S100A5,S100A9,S100G,SERPINB10,SLC4A1,SLC9A5,SOX18,SPI1,STR2,TAC3,TACSTD2,TAL1,THBS1,TUBA8,UBASH3A	
5 calcitriol		chemical drug		0.412		0.000000 494	ADAM19,ALPI,ANGPT2,ANKRD2,ATP5D,BIRC5,CA2,CALB1,CALCB,CASR,CCNA1,CCR8,CDC20,CDC45,CDK1,CDK5R1,CDKN1A,CEBPB,CELSR3,CHAF1A,CHAF1B,CHGA,CKM,CO4A1,CSF1,CSF2,CXCL2,CXCL3,CXCL8,CYP24A1,CYP2C9,CYP3A4,CYP46A1,CYR61,DCSTAMP,DEFB4A/DEFB4B,DUSP1,DUSP10,EDN1,EGR1,ETFB,EXO1,FABP4,FAM107A,FCE R2,FOS,GADD45A,GADD45G,GEM,HBEGF,HSPB7,IER3,IFITM1,IFNG,IGF1,IGFBP5,IL10RA,IL17A,IL18,IL1A,IL1RN,IL4,INCENP,INS,ITGA4,ITGAM,ITGB7,JUN,KIF20A,KIF22,KL,KLK13,KLK5,LEP,LIG1,LPAR1,LPL,LTBP1,MAOA,MCM2,MCM5,MMP1,MRC1,MYH8,NEK2,NFATC1,NKX2-1,NME4,NPHS1,NTHL1,NUPR1,NUSAP1,PDE9A,POLE,POU1F1,PRC1,PRKCB,PRKCD,PTGER2,PTGFR,PTH,RAB38,RAD51API,RARRES1,	140 (2)

Upstream Regulator	Expr Fold Change	Molecule Type	Predicted Activation State	Activation z-score	Flags	p-value of overlap	Target molecules in dataset	Mechanistic Network
							RBPMS,REL,RRM2,RUNX1T1,S100A9,S100G,SERPINB7,SERPINB9,SLC2A4,SLC7A7,SNPH,SOCS3,SPAG5,STMN1,SUV39H1,TACC3,TERT,THBD,THBS1,THRA,TK1,TLR2,TLR4,TNF AIP3,TPX2,TSPO,WNT11	
6 ID2	1.706	transcription regulator		-1.136		0.000000 514	AICDA,ASCL2,BATF,CCR10,CCR7,CCR8,CD40LG,CDC25B,CDK1,CDKN1A,CDKN2C,CEBPB,CSF1,CXCR4,CXCR5,DUSP1,DUSP10,DUSP4,E2F2,EGR2,EGR4,FLT3LG,FOXO3,FSHB,GADD45B,GADD45G,IFNG,IL10RA,IL4,IL4R,IL9R,IRF8,KLF6,LTA,MAP3K14,MPZ,NFAT5,NFATC1,NR4A3,PDCD1,PTPN13,PTPN14,PTPN22,RAPGEF4,REL,RPS6KA2,SELL,SEMA3F,SH2D1A,SOCS3,SOX4,SOX5,TNFRSF25,TNFSF14,TNFSF8,TRAF1,TRAF5	
7 phorbol myristate acetate		chemical drug	Activated	7.684	bias	0.000000 604	ADAM28,ADAM8,ADM,ADRB3,AGER,ALOX12,ANGPT2,ANGPTL4,ANXA1,AQP4,ATP2A3,AURKA,AURKB,BCL2A1,BDNF,BIRC5,BLM,BTG2,C5AR1,CA2,CA8,CAV1,CCL1,CCL4,CNA1,CCR7,CD209,CD28,CD36,CD40LG,CD69,CDK1,CDK5R1,CDK5R2,CDKN1A,CDKN2B,CGA,CHGA,CKM,CLCF1,CRH,CRHR1,CSF1,CSF2,CTLA4,CXCL13,CXCL2,CXCL3,CXCL8,CXCR2,CXCR4,CYBB,CYP24A1,CYP2A6 (includes others),CYR61,DEFB4A/DEFB4B,DSG1,DUSP1,DUSP2,DUSP5,E2F1,E2F3,EGR1,EGR2,EGR3,EGR4,EIF4EBP1,ELANE,EN1,EP300,EPOR,ERBB4,FGF2,FGF7,FOS,FOSB,FOSL1,FSHB,FUT9,GABRP,GAP43,GATA1,GATA2,GDF15,GEM,GML,GNRH1,GRIN2A,H1FX,HAS1,HBEGF,HDC,HMGA1,HPSE,HSD11B1,HSD17B1,HSD3B1,HTR2A,HTR7,IFNG,IGF1,IGFBP2,IGFBP5,IL12RB1,IL17A,IL18,IL1A,IL1RN,IL20RA,IL24,IL4,ITGAM,ITM2A,JUN,JUNB,JUND,KCNJ10,KIF2C,KLF2,KLF6,KLK3,KRT35,LAMB3,LOR,LPL,LTA,LYVE1,MAD1L1,MMP1,MMP11,MMP12,MMP14,MMP19,MMP7,MPZ,MRC1,MSR1,MST1R,MST2A,MUC4,MYH7,MYOZ2,NCR1,NFAT5,NFATC1,NFKBIA,NFKBIE,NKX2-1,NOCT,NR4A2,NTS,OLR1,OSM,OSR2,PAK2,PDCD1,PDE1C,PDPN,PIM1,PLIN3,PODXL2,PON1,POU1F1,PPP1R15A,PRKC	276 (3)

Upstream Regulator	Expr Fold Change	Molecule Type	Predicted Activation State	Activation z-score	Flags	p-value of overlap	Target molecules in dataset	Mechanistic Network
							B,PRKCD,PRKD1,PTGER2,PTGES,PTGFR,PTPRE,PTPRN,PTPRO,RAE1,RARA,RARB,RASGRP1,RECQL4,REL,RELB,RGS1,RGS2,RUVBL2,S100A9,SELL,SELPLG,SERPINB10,SERPINB7,SERPINB9,SLC22A1,SLC6A2,SLC6A7,SLC7A11,SNAI1,SNAP25,SOCS3,SP4,SPHK1,SRC,SRD5A2,SSTR2,STATH,TACR1,TBXAS1,TEAD4,TERT,TH,THBS1,TIE1,TK1,TLR2,TLR4,TLR6,TMOD2,TNFAIP3,TNFRSF1B,TNFSF14,TRAF1,TRPC6,ULBP2,USF2,VIP,WT1,XCR1,ZFP36	
8 HDAC1	0.743	transcription regulator		-0.945		0.000000942	ADIPOQ,AMPD3,ANGPT2,ASCL2,ATF3,BDNF,CCNA1,CCNB2,CCR8,CD27,CD34,CDC25A,CDC25C,CDK1,CDKN1A,COL1A2,COL9A1,CXCL8,E2F2,EGR1,EHMT2,FABP4,FAM107A,FOS,H2AFX,HBE1,HBG2,IFNB1,IL17A,IL24,IL4,INA,ITGB4,KLK3,LIG1,MAD1L1,MCM5,MPZ,MT1G,MUC4,MYH7,NEFH,NFATC1,NFKBIA,NKX2-5, PAX3, PLK1, PMAIP1, POLL, PPP2R2B,PRIM2,PTH,RAD54L,RAG1,RECQL4,RELB,RGS10,RRM2,RUNX2,S100A9,SATB1,SNAI1,SOX10,TAGLN,TAL1,TBX1,TBX2,TERT,TUBB3,TYMS	414 (12)
9 PTGER2	2.853	g-protein coupled receptor	Activated	5.127	bias	0.00000162	AURKA,CCNB2,CCR7,CDKN3,CENPE,CFP,CKS2,CXCL8,CXCR2,CXCR4,EGR1,FPR1,H2AFX,HAMP,HDC,HIST1H2AB,IFNG,IL17A,IL1A,KIF15,KIF20A,KIF22,KIF2C,KLRD1,MKI67,NEK2,NUSAP1,PIM1,PLK1,PRC1,PTGER3,PTGES,SPAG5,THBS1,TPX2,TROAP,TTK	
10 TNF	1.621	cytokine	Activated	8.752	bias	0.00000184	ACTA1,ADAM8,ADAMTS5,ADIPOQ,ADM,ADRB1,ADRB3,AEBP1,AGER,AICDA,AMPD3,ANGPT2,ANGPTL4,ANXA1,ARHGDI,ATF3,AURKC,BCL2A1,BDKRB1,BDKRB2,BDNF,BIK,BIRC5,BTG2,BTG3,C5AR1,CA2,CABP1,CAV1,CCK,CCL1,CCL22,CCL4,CCL7,CCR1,CCR5,CCR7,CCR8,CD1C,CD209,CD247,CD28,CD36,CD3E,CD40LG,CD5,CD69,CD82,CDC25C,CDH13,CDH5,CDK5R1,CDKN1A,CDKN2C,CDX1,CEBPB,CEBPG,CHI3L1,CHRNA4,CHRNA2,CHRNE,CHRNA3,CHST4,CHST7,CIB2,CKM,CLCF1,CLDN7,CNN1,COL15A1,COL16A1,COL1A2,COLQ,COTL1,CPA3,CRH,CRHR1,CRLF1,CRYAB,CSF1,CSF2,CSN2,CST7,CTLA4,CTSF,CX3CR1,CXCL1,CXCL13,CXCL2,C	611 (12)

Upstream Regulator	Expr Fold Change	Molecule Type	Predicted Activation State	Activation z-score	Flags	p-value of overlap	Target molecules in dataset	Mechanistic Network
							XCL3,CXCL5,CXCL8,CXCR2,CXCR4,CXCR5,CYBB,CYP26B1,CYP2C8,CYR61,CYTH3,DCSTAMP,DEFB4A/DEFB4B,DPF3,DUSP1,DUSP10,DUSP2,DUSP4,DUSP5,DVL1,E2F1,EDN1,EGFR1,EGR2,EGR3,ELF3,EMCN,EMP2,ENG,ENPP3,EREG,ESM1,FABP4,FAT2,FCAR,FCER2,FCGR2B,FGF2,FGF5,FOS,FOSB,FOSL1,FOXF1,FOXF2,FPR1,FPR2,FSCN1,G0S2,GABRA1,GADD45A,GADD45B,GADD45G,GATA2,GCLM,GDF15,GEM,GN A15,GNL1,GPR176,GPRC5B,GRIA1,HAS1,HBEGF,HDC,HIVE P1,HLA-F, HMOX1,HOXB8,HRK,HSD11B1, HSPA1A/HSPA1B,HSPG2,ICAM2,IER2,IER3,IFI27,IFITM1,IFNA1/IFNA13,IFNB1,IFNG,IGF1,IGFBP2,IGFBP5,IL10RA,IL17A,IL18,IL18R1,IL1A,IL1RN,IL24,IL3,IL36RN,IL3RA,IL4,IL4R,INS,IRF8,ITGA4,ITGAM,ITGB7,JUN,JUNB,JUND,KIF20A,KITLG,KL,KLF10,KLF2,KLF6,KLK3,LAMA4,LAMB3,LBP,LEP,LLPL,LTB4R2,LYVE1,MADCAM1,MAFF,MAP3K14,MC1R,MCF2,MECOM,MFHAS1,MGMT,MMP1,MMP10,MMP12,MMP14,MMP28,MMP7,MSR1,MST1R,MSTN,MT2A,MUC1,MUC4,MYH7,NCAN,NCF2,NEFH,NFATC1,NFKB1A,NFKB1E,NKX21,NKX6-1, NOCT,NOD2,NPHS1,NPPB,NR4A2,NR4A3,NR6A1,OAS2,OLR1,OSM,OTUD7B,P2RY6,PAK2,PAX6,PDCD1,PDE2A,PDGFRA,PDPN,PIM1,PLA2G3,PLA2G4C,PLA2G5,PLIN1,PLK2,PLP1,PMAIP1,PPP1R15A,PRKCD,PRSS23,PTGES,PTGFR,PTPRN,PYCARD,RARA,RBPMS,RCAN2,REL,RELB,RFX2,RGS1,RGS2,RGS20,RGS3,RGS5,RND1,RRAD,RRM1,RRM2,RUNX2,S100A9,SCNN1B,SCO2,SCUBE2,SELL,SELPLG,SERPINB10,SERPINB9,SLC12A1,SLC16A2,SLC1A3,SLC2A4,SLC7A8,SNAI1,SNN,SOCS2,SOCS3,SOX4,SPHK1,ST8SIA4,STMN1,SYNGR3,TAGLN,TBXAS1,TERT,TH,THBD,THBS1,THBS2,TIE1,TK1,TLR2,TLR4,TNC,TNFAIP3,TNFRSF1B,TNFRSF9,TNFSF14,TNFSF15,TNFSF8,TNFSF9,TNNC1,TRAF1,TRAF2,TRAF5,TREM2,TRIM15,TRPC3,TRPC6,TWIST1,TXNRD1,VIP,WNT10B,WNT5A,WNT7A,YY1,ZFP36	

Table E.5-2. Top ten canonical pathways affected by 6-hour exposure to 20 µM antimony(III) potassium tartrate trihydrate

Order	Ingenuity Canonical Pathways	-log(p-value)	Ratio	Molecules
1	Agranulocyte Adhesion and Diapedesis	4.29	0.358	CLDN7,CCL8,SELL,CLDN8,PODXL2,MYH3,CXCR4,IL1RN,CLDN14,C5AR1,MYH11,CXCL1,MYH7,MMP11,MADCAM1,MYL6B,CDH5,CXCL8,IL18,XCL2,CXCR2,SELPLG,IL1A,IL36RN,ITGA4,ACTA1,CXCL3,CD34,CXCL14,MMP1,MMP25,ITGA3,MMP12,ITGB7,CXCL5,CCL4,MMP28,ICAM2,CLDN17,CCL7,CLDN18,CCL1,MMP17,CCL21,CXCL2,MMP7,CCL22,MMP19,CXCL13,MMP20,MMP10,MYH8,MMP14
2	Granulocyte Adhesion and Diapedesis	3.79	0.35	CLDN7,CCL8,SELL,CLDN8,CXCR4,IL1RN,CLDN14,C5AR1,CXCL1,MMP11,TNFRSF1B,FPR2,CDH5,CXCL8,HSPB1,IL18,XCL2,CXCR2,SELPLG,IL1A,IL36RN,ITGA4,CXCL3,CXCL14,MMP1,HRH2,MMP25,ITGAM,ITGA3,MMP12,FPR1,HRH4,CXCL5,CCL4,MMP28,ICAM2,CLDN17,CCL7,CLDN18,CCL1,MMP17,CCL21,CXCL2,MMP7,CCL22,MMP19,CXCL13,MMP20,MMP10,MMP14
3	Eicosanoid Signaling	3.63	0.449	DPEP3,ALOX12,FPR2,PTGER1,LTBR2,PLA2G7,PLA2G6,DPEP1,PLA2G3,PLA2G5,PTGER2,PTGIS,PTGFR,PLA2G4C,TBXA2R,PLA2G2E,ALOX12B,PTGES,PTGIR,PTGER3,ALOX15,TBXAS1
4	Role of Cytokines in Mediating Communication between Immune Cells	3.28	0.444	IFNA10,IL3,CSF2,IFNG,IL4,IFNA7,IFNA14,CXCL8,IL26,IL18,IL1RN,IL25,IFNA1/IFNA13,IL24,IL1A,IL36RN,IL17A,IFNA16,IFNB1,IFNA4
5	Role of Hypercytokinemia/hyperchemokine in the Pathogenesis of Influenza	2.91	0.447	IFNA10,IFNG,CCR1,IFNA7,IFNA14,CXCL8,CCR5,IL18,IL1RN,IFNA1/IFNA13,IL1A,IL36RN,IL17A,CCL4,IFNA16,IFNB1,IFNA4
6	Bladder Cancer Signaling	2.33	0.351	FGF5,MMP25,FGF1,MMP12,E2F1,THBS1,FGF20,SUV39H1,CDKN1A,MMP28,FGF12,MMP11,FGF21,FGF7,FGF3,FGF2,MMP17,CXCL8,MMP7,MMP19,FGF16,MMP20,MMP10,FGF8,MMP1,FGF9,MMP14
7	Crosstalk between Dendritic Cells and Natural Killer Cells	2.24	0.346	CAMK2B,CCR7,IL3,CSF2,IFNG,TREM2,HLA-F,FSCN2,TLR4,CD209,FSCN1,FSCN3,KIR2DL2,TNFRSF1B,PRF1,IL4,LTA,NECTIN2,CD69,IL3RA,KLRD1,IL18,CD40LG,CD28,IFNA1/IFNA13,ACTA1,IFNB1
8	Role of IL-17A in Psoriasis	2.2	0.583	S100A9,CXCL1,IL17A,DEFB4A/DEFB4B,CXCL3,CXCL5,CXCL8
9	Role of Wnt/GSK-3β Signaling in the Pathogenesis of Influenza	2.1	0.362	IFNA10,IFNG,WNT5A,LEF1,FZD2,WNT5B,IFNA7,IFNA14,FZD7,DVL1,FZD9,WNT2B,WNT11,WNT8B,IFNA1/IFNA13,WNT7A,IFNA16,IFNB1,APC2,IFNA4,WNT10B
10	Oxidized GTP and dGTP Detoxification	1.99	1	RUVBL2,NUDT1,DDX6

Pathways 1, 2, 4, 5, 7, 8, and 9 (green background) are related to immune reactions. Pathway 6 (with orange background) is related to cancer. Pathway 10 (with yellow background) is related to oxidative stress.

E.6: Immune effects from compounds containing pentavalent antimony

This appendix lists immune function from compounds containing pentavalent antimony (Table E.6-1).

Table E.6-1. Effects of compounds containing pentavalent antimony on immunity

Patients, species or experimental system	Antimony compound	Immune effects	Functional or mechanistic association	Reference
Human studies				
Healthy active duty soldiers treated for leishmaniasis	Sodium stibogluconate	Transient lymphopenia (decreased CD4 ⁺ and CD8 ⁺ T cells)	Increased susceptibility to Herpes Zoster infections	Wortmann <i>et al.</i> 1998
Patients treated for cutaneous leishmaniasis	Glucantime (meglumine antimoniate)	Elevated IL-1 β , TNF- α , IL-6 and IL-8	Amplified pro-inflammatory cytokines upon exposure to antimonials	Kocyigit <i>et al.</i> 2002
Patients treated for visceral leishmaniasis	Sodium stibogluconate	Elevated IL-1 β , TNF- α , IL-6, GM-CSF, and C1q-binding circulating immune complexes (CIC)	Amplified pro-inflammatory cytokines and CIC-induced GM-CSF upon exposure to antimonials	Elshafie <i>et al.</i> 2007
Animal studies				
BALB/c mice	Antimony sodium gluconate	Activation of peritoneal macrophages associated with enhanced antigen presentation to T cells	Increased macrophage membrane fluidity and enhanced antigen presentation capacity	Ghosh <i>et al.</i> 2013
Normal C57BL/6 mice, IFN γ gene knockout mice, inducible nitric oxide synthase-knockout (iNOS KO) mice, and respiratory burst-deficient gp91 ^{phox-/-} (X-linked chronic granulomatous disease [X-CGD]) mice	Sodium stibogluconate	In IFN γ gene knockout mice, pentavalent antimony inhibited but did not kill intracellular <i>Leishmania donovani</i> ; treatment was effective in killing the parasite in normal, iNOS KO, and X-CGD mice.	Results support a role for T cell-derived IFN γ as a critical host factor required for the efficacy of antimony in promoting parasite killing	Murray and Delph-Etienne 2000
BALB/c mice	Antimony sodium gluconate	Sodium stibogluconate synergizes with IL-2 to promote IFN γ -dependent anti-Renca tumor immune response	Supports a role for pentavalent antimony in promoting IFN γ -dependent anti-tumor immune response	Fan <i>et al.</i> 2009

Patients, species or experimental system	Antimony compound	Immune effects	Functional or mechanistic association	Reference
<i>In vitro</i> studies				
Murine Baf 3 cell line and TF-1 human myeloid leukemia cells	Sodium stibogluconate	Sodium stibogluconate is a potent inhibitor of protein tyrosine phosphatases including Src homology PTPase1 (SHP-1), SHP-2, and PTP1B	Sodium stibogluconate, which contains a pentavalent antimony atom, (but not antimony(III) potassium tartrate) can alter signaling of multiple cytokines (IL-3, IFN α , and GM-CSF) that signal through receptor tyrosine kinases regulated by PTPases	Pathak and Yi 2001
Various cancer cell lines	Sodium stibogluconate	Sodium stibogluconate enhanced IFN α -induced Stat1 tyrosine phosphorylation, inactivated intracellular SHP-1 and SHP-2, and induced cellular protein tyrosine phosphorylation in cancer cell lines	Sodium stibogluconate treatment was found to synergize with IFN α to overcome cancer cell lines that were refractory to the anti-cancer effects of IFN α <i>in vitro</i> and <i>in vivo</i>	Yi <i>et al.</i> 2002
Human CD4+ and CD8+ T lymphocytes from healthy donors and melanoma patients	Sodium stibogluconate	Sodium stibogluconate synergizes with IL-2 to potentiate induction of IFN γ + T cells	Sodium stibogluconate treatment may potentiate T cell function in the presence of IL-2	Fan <i>et al.</i> 2009

Appendix F: Other Relevant Information

F.1 Studies of antimony(III) potassium tartrate carcinogenicity in experimental animals

This appendix includes neoplasms induced in experimental animals exposed to antimony potassium tartrate (Table F.1-1), details of these animal studies (Table F.1-2) and risk of bias rating of Schroeder *et al.* (1970) study (male rats in Table F.1-3, female rats in Table F.1-4) and Kanisawa and Schroeder (1969) study (Table F.1-5)

Table F.1-1. Neoplasms induced in experimental animal carcinogenicity studies by drinking water studies of antimony potassium tartrate

Studies are presented in the order of descending overall utility.

Species strain/stock*	Site	Classification	Neoplasms (Sex of animal)	Reference
Rat, Long-Evans	None	None	None (M and F)	Schroeder <i>et al.</i> 1970
Mouse, Swiss CD-1	None	None	None (M and F)	Schroeder <i>et al.</i> 1968, Kanisawa and Schroeder 1969

F = female, M = male.

Table F.1-2. Cancer studies in experimental animals exposed to antimony(III) potassium tartrate

Reference and study design	Exposure	Tumor site – Tumor type		Comments
		Dose levels	Tumor incidence (n/N) (%)	
Schroeder <i>et al.</i> 1970 Animal: Rat — Long-Evans (random bred) M, F Animal age at the beginning of exposure: NR (possibly at weaning) Study duration: ~4 years	Agent and purity: Antimony potassium tartrate NR Exposure route: Drinking water Exposure concentrations, frequency, and duration: 0 5 ppm not clearly reported (possibly ad libitum x life-span)	Whole body – Tumor NOS (M)		Survival: The survival of females at 50% death ($P < 0.025$ by chi-square analysis) and males and females for longevity (mean age of the last surviving 10%) ($P < 0.001$ by Student's t test) was significantly reduced compared to untreated controls. Body weight: Both males and females were similar to controls. Overall utility: [+] The study has low utility because of many limitations, including only reporting grossly visible tumors without organ site or tumor type.
		0	10/50 (20%)	
		5	6/50 (12%)	
		Whole body – Tumor NOS (F)		
		0	14/39 (35.9%)	
		5	18/47 (38.3%)	

Reference and study design	Exposure	Tumor site – Tumor type		Comments
		Dose levels	Tumor incidence (n/N) (%)	
Kanisawa and Schroeder 1969 Animal: Mouse — White Swiss CD-1 (Random bred) M+F (combined) Animal age at the beginning of exposure: Weanling Study duration: Life span	Agent and purity: Antimony potassium tartrate NR Exposure route: Drinking water Exposure concentrations, frequency, and duration: 0 5 µg/mL in double deionized water ad libitum x life span	Whole body – Tumor NOS		Survival: Survival was similar to controls. Body weight: Males were sporadically lower than controls at 90, 150, and 540 days, while females were more consistently lower at 150, 360, and 540 days. Other comments: The incidences were reported for both sexes combined, but it was stated that none of the neoplasms were significantly increased. Overall utility: [+] This study is of low utility due to many limitations, including only one tested concentration (below maximally tolerated dose for males, and close to or at maximally tolerated dose for females), unknown test substance purity, tumor incidences only reported in combined sexes with no histologic information, and lack of site specific information (except incidences of three sites in sexes combined). Data lack sufficient details to allow us determine whether any specific type of tumor had increased in a sex.
		0	24/71 (33.8%)	
		5	18/76 (23.7%)	
		Whole body – Malignant tumor NOS		
		0	8/71 (11.3%)	
		5	6/76 (7.9%)	
		Whole body – Benign tumor NOS		
		0	16/71 (22.5%)	
		5	12/76 (15.8%)	
		Mammary gland – Tumor NOS		
		0	1/71 (1.4%)	
		5	3/76 (3.9%)	
		Lung – Tumor NOS		
		0	15/71 (21.1%)	
5	10/76 (13.2%)			
Liver – Tumor NOS				
0	4/71 (5.6%)			
5	1/76 (1.3%)			

F = female; M = male; n/N = number of animals with neoplasms divided by the total number of animals tested in that dose group; NR = not reported; NOS = not otherwise specified

Table F.1-3. Schroeder *et al.* (1970) study of male rats and female rats exposed to antimony(III) potassium tartrate in drinking water for the life span of the animals

Utility question	Rating	Rationale
Study design		
Randomization	NR	Randomization and initial body weights were not reported.
Controls	+++	Concurrent control group, exposed to untreated drinking water, had the same number of animals as exposure group did.
Historical data		No
Animal model	+++	Both sexes of a random bred strain, which increases external validity.
Statistical power	+++	Large numbers of animals (51 males, 59 females) per concentration group were used.
Exposure		
Chemical characterization	NR	Not reported, not even purity.
Dosing regimen	++	The maximally tolerated dose level was not reached, because the treated group did not show decreased body weight compared to the control group, although the median life spans and longevity (mean age of the last surviving 10%) for both sexes were decreased by the treatment. The dose might not have been high enough to detect neoplastic effects.
Exposure duration	+++	Though exposure duration was never clearly stated, this study appears to use a life time exposure.
Dose-response	+	Only one dose level was tested and no basis for that level was reported. All other elements were administered at the same level, except for lead.
Outcome		
Pathology	+	Only grossly visible tumors were reported. The methods stated that gross tumors were fixed, but did not state that they were stained or microscopically examined. Consequently, small tumors might have been missed.
Consistency between groups	++	A pneumonia epidemic killed many rats and the death rates varied among the groups.
Study duration	+++	Life time study, because the animals were observed until their nature death (as compared to scheduled euthanization after a predetermined exposure period).
Confounding		
Confounding	++	Pneumonia killed various numbers of animals per group, before penicillin treatment controlled the disease. It is unclear that all rats, or only visibly sick rats, received penicillin. Furthermore, the disease might in effect select stronger/healthier animals (than the general population) to complete the study. Additionally, test substance purity was unknown.
Reporting and analysis		
Reporting data and statistics	++	The statistical methods and results of survival measures were reported, but statistical analysis of tumor incidences were not reported.
Combining lesions	+	Tumors were counted based on gross observation, not histological analysis occurred.

Overall utility: +. The study has low utility because of many limitations, including only reporting grossly visible tumors without organ site or tumor type.

NR = Not reported, +++ = High utility, ++ = Moderate utility, + = Low utility.

Table F.1-4. Kanisawa and Schroeder (1969) study of male and female (combined) mice exposed to antimony potassium tartrate in drinking water for the lifespan of the animals

Utility question	Rating	Rationale
Study design		
Randomization	NR	Not reported.
Controls	+++	Concurrent control group, exposed to doubly deionized water with added essential trace elements, had the same number of animals as the exposure group did.
Historical data		No
Animal model	+++	Both sexes of random bred mice were used, giving a high level of external validity.
Statistical power	+++	A large number (54) of mice per sex per group were used.
Exposure		
Chemical characterization	NR	No chemical characterization was reported, not even purity.
Dosing regimen	+	The maximally tolerated dose level was not reached, because the treated group did not show decreased body weight compared to the control group. The dose might not have been high enough to detect neoplastic effects.
Exposure duration	+++	Mice were exposed for their lifetimes.
Dose-response	+	Only one concentration was tested and no rationale for the dose selection was reported. Dose response relationships cannot be evaluated due to only one dose level.
Outcome		
Pathology	++	Only gross lesions were microscopically evaluated.
Consistency between groups	++	The treated and control groups were treated the same while mice were alive. The examination of organs/tissues varied, because only gross lesions (not all major organs) were examined microscopically.
Study duration	+++	The study duration was lifetime, up to the animals' natural death.
Confounding		
Confounding	+++	Testing substance purity and supplier were unknown. Exposure to antimony via other sources (feed, housing) was negligible because the feed was antimony free and metal exposure via housing was minimized.
Reporting and analysis		
Reporting data and statistics	++	Although tumor incidents were not statistically analyzed in the study, the data were reported and enabled us to conduct statistical analysis. Statistical methods were described as "numerical data were treated by Chi-square analysis and by Student's t test", but the reported probability in tables did not specify the result was from which method.
Combining lesions	++	Tumor incidence was reported for two sexes combined only, instead of male and female separately. Site specific (lung, liver, mammary gland, and other) information was limited to tumor incidence, with no subtype. Tumors were also grouped by origin (epithelial, non-epithelial) along with being benign or malignant. Overall information did not allow detecting of specific tumor type increase in either sex.

Overall utility: +. Due to many limitations, including only one tested concentration (below maximally tolerated dose), unknown test substance purity, tumor incidences only reported in combined sexes with no histologic information, and lack of site specific information (except incidences of three sites in sexes combined), this study is of low utility. Data lack sufficient details to allow us determine whether any specific type of tumor had increased.

NR = Not reported, +++ = High utility, ++ = Moderate utility, + = Low utility.

References

1. Andrewes P, Kitchin KT, Wallace K. 2004. Plasmid DNA damage caused by stibine and trimethylstibine. *Toxicol Appl Pharmacol* 194(1): 41-48. (Supported by the National Research Council and the US EPA. Authors affiliated with U.S. Environmental Protection Agency, NC.)
2. Asakura K, Satoh H, Chiba M, Okamoto M, Serizawa K, Nakano M, Omae K. 2009. Genotoxicity studies of heavy metals: lead, bismuth, indium, silver and antimony. *J Occup Health* 51(6): 498-512. (Supported by the IMS (Intelligent Manufacturing Systems) Promotion Center, Manufacturing Science and Technology Center. Authors affiliated with Keio University School of Medicine, Japan; Tohoku University Graduate School of Medicine, Japan; International University of Health and Welfare, Japan; Hitachi Ltd., Japan.)
3. Cavallo D, Iavicoli I, Setini A, Marinaccio A, Perniconi B, Carelli G, Iavicoli S. 2002. Genotoxic risk and oxidative DNA damage in workers exposed to antimony trioxide. *Environ Mol Mutagen* 40(3): 184-189. (Support not reported. Authors affiliated with ISPESL-National Institute for Occupational Safety and Prevention, Italy; Università Cattolica del Sacro Cuore, Italy.)
4. Chiu WA, Guyton KZ, Martin MT, Reif DM, Rusyn I. 2017. Use of high-throughput *in vitro* toxicity screening data in cancer hazard evaluations by IARC Monograph Working Groups. *ALTEX*: 14 pp. (Supported by NIH and NIEHS. Authors affiliated with Texas A&M University, TX; IARC, France; US EPA, NC; North Carolina State University, NC.)
5. Dopp E, Hartmann LM, Florea AM, von Recklinghausen U, Rabieh S, Shokouhi B, Hirner AV, Rettenmeier AW. 2006. Trimethylantimony dichloride causes genotoxic effects in Chinese hamster ovary cells after forced uptake. *Toxicol Vitro* 20(6): 1060-1065. (Supported by the German Research Foundation. Authors affiliated with University Hospital Essen, Germany; University of Duisburg-Essen, Germany.)
6. El Nahas S, Temtamy SA, de Hondt HA. 1982. Cytogenetic effect of two antimonial antibilharzial drugs: tartar emetic and bilharcid. *Environ Mutagen* 4(1): 83-91. (Support not reported. Authors affiliated with National Research Centre, Egypt; University of Waterloo, Canada.)
7. El Shanawany S, Foda N, Hashad DI, Salama N, Sobh Z. 2017. The potential DNA toxic changes among workers exposed to antimony trioxide. *Environ Sci Pollut Res Int* 24(13): 12455-12461. (Support not reported. Authors affiliated with University of Alexandria, Egypt.)
8. Elliott BM, Mackay JM, Clay P, Ashby J. 1998. An assessment of the genetic toxicology of antimony trioxide. *Mutat Res* 415(1-2): 109-117. (Support not reported. Authors affiliated with Central Toxicology Laboratory, UK.)
9. Elshafie AI, Åhlin E, Mathsson L, ElGhazali G, Rönnelid J. 2007. Circulating immune complexes (IC) and IC-induced levels of GM-CSF are increased in sudanese patients with

- acute visceral *Leishmania donovani* infection undergoing sodium stibogluconate treatment: implications for disease pathogenesis. *J Immunol* 178(8): 5383-5389. (Supported by the Swedish Society of Medicine, King Gustav V's 80-years Fund, Swedish League against Rheumatism, Ugglas Foundation, Groschinsky Foundation, Viberg Foundation, and Swedish Fund for Research Without Animal Experiments. Authors affiliated with Uppsala University, Sweden; Alribate University Hospital, Sudan; University of Khartoum, Sudan; King Fahad Medical City, Saudi Arabia.)
10. EPA. 2017. *iCSS ToxCast Dashboard*. U.S. Environmental Protection Agency. <https://actor.epa.gov/dashboard/>. Accessed on 10/10/17.
 11. EU. 2008. *European Union Risk Assessment Report. Diantimony Trioxide*. Luxembourg: European Communities. 556 pp.
 12. Fan K, Borden E, Yi T. 2009. Interferon-gamma is induced in human peripheral blood immune cells *in vitro* by sodium stibogluconate/interleukin-2 and mediates its antitumor activity *in vivo*. *J Interferon Cytokine Res* 29(8): 451-460. (Support not reported. Authors affiliated with Lerner Research Institute, OH; Cleveland Clinic Foundation, OH; Taussig Cancer Center, OH.)
 13. Gebel T, Christensen S, Dunkelberg H. 1997. Comparative and environmental genotoxicity of antimony and arsenic. *Anticancer Res* 17(4a): 2603-2607. (Support not reported. Authors affiliated with University of Goettingen, Germany.)
 14. Gebel T, Birkenkamp P, Luthin S, Dunkelberg H. 1998. Arsenic(III), but not antimony(III), induces DNA-protein crosslinks. *Anticancer Res* 18(6a): 4253-4257. (Support not reported. Authors affiliated with University of Goettingen, Germany.)
 15. Ghosh M, Roy K, Roy S. 2013. Immunomodulatory effects of antileishmanial drugs. *J Antimicrob Chemother* 68(12): 2834-2838. (Supported by the Network Project and CSIR India. Authors affiliated with CSIR-Indian Institute of Chemical Biology, India.)
 16. Gross P, Brown JH, Westrick ML, Srsic RP, Butler NL, Hatch TF. 1955. Toxicologic study of calcium halophosphate phosphors and antimony trioxide. I. Acute and chronic toxicity and some pharmacologic aspects. *AMA Arch Ind Health* 11(6): 473-478. (Supported by General Electric Company and Sylvania Electric Products, Inc. Authors affiliated with Mellon Institute, PA; Emory University, GA.)
 17. Grosskopf C, Schwerdtle T, Mullenders LH, Hartwig A. 2010. Antimony impairs nucleotide excision repair: XPA and XPE as potential molecular targets. *Chem Res Toxicol* 23(7): 1175-1183. (Supported by the DFG. Authors affiliated with Technische Universität Berlin, Germany; Westfälische Wilhelms-Universität Münster, Germany; Leiden University Medical Center, Netherlands.)
 18. Groth DH, Stettler LE, Burg JR, Busey WM, Grant GC, Wong L. 1986. Carcinogenic effects of antimony trioxide and antimony ore concentrate in rats. *J Toxicol Environ Health* 18(4): 607-626. (Supported by NIOSH and Midwest Research Institute. Authors affiliated with NIOSH, OH; Experimental Pathology Laboratories, VA; College of William and Mary, VA; Midwest Research Institute, MO.)

19. Gurnani N, Sharma A, Talukder G. 1992a. Cytotoxic effects of antimony trichloride on mice *in vivo*. *Cytobios* 70(281): 131-136. (Supported by the Council of Scientific and Industrial Research, Government of India. Authors affiliated with University of Calcutta, India; Vivekananda Institute of Medical Sciences, India.)
20. Gurnani N, Sharma A, Talukder G. 1992b. Comparison of the clastogenic effects of antimony trioxide on mice *in vivo* following acute and chronic exposure. *Biomaterials* 5(1): 47-50. (Supported by the Council of Scientific and Industrial Research. Authors affiliated with University of Calcutta, India; Vivekananda Institute of Medical Sciences, India.)
21. Hashemzaei M, Pourahmad J, Safaeinejad F, Tabrizian K, Akbari F, Bagheri G, Hosseini MJ, Shahraki J. 2015. Antimony induces oxidative stress and cell death in normal hepatocytes. *Toxicol Environ Chem* 97(2): 256-265. (Supported by the Zabol University of Medical Sciences. Authors affiliated with Zabol University of Medical Sciences, Iran; Shahid Beheshti University of Medical Sciences, Iran; Zanjan University of Medical Sciences, Iran.)
22. Huang H, Shu SC, Shih JH, Kuo CJ, Chiu ID. 1998. Antimony trichloride induces DNA damage and apoptosis in mammalian cells. *Toxicology* 129(2-3): 113-123. (Supported by the National Science Council, Republic of China. Authors affiliated with National Tsing-Hua University, Taiwan.)
23. IARC. 2017. *Some Organophosphate Insecticides and Herbicides*, IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. vol. 112, Lyon, France: International Agency for Research on Cancer. 464 pp.
24. Johns Hopkins University. 2017. *Species Specific Information: Rat*. Johns Hopkins University. <http://web.jhu.edu/animalcare/procedures/rat.html>. Accessed on 8/18/17.
25. Jones RD. 1994. Survey of antimony workers: mortality 1961-1992. *Occup Environ Med* 51(11): 772-776. (Support not reported. Authors affiliated with Corporate Health Limited, Slough.)
26. Jones SR, Atkin P, Holroyd C, Lutman E, Battle JV, Wakeford R, Walker P. 2007. Lung cancer mortality at a UK tin smelter. *Occup Med (Lond)* 57(4): 238-245. (Supported by Rio Tinto plc. Authors affiliated with Westlakes Research Institute, UK; Rio Tinto plc, UK; University of Manchester, UK.)
27. Kanematsu N, Hara M, Kada T. 1980. Rec assay and mutagenicity studies on metal compounds. *Mutat Res* 77(2): 109-116. (Supported by the Ministry of Welfare, the Ministry of Education, the Science and Technology Agency of Japan, the Takamatsunomiya Cancer Foundation and the Nissan Science Foundation. Authors affiliated with Gifu College of Dentistry, Japan; National Institute of Genetics, Japan.)
28. Kanisawa M, Schroeder HA. 1969. Life term studies on the effect of trace elements on spontaneous tumors in mice and rats. *Cancer Res* 29(4): 892-895. (Supported by the USPHS, the US Army, the American Cancer Society, and the CIBA Pharmaceutical Company. Authors affiliated with Dartmouth Medical School, NH; Brattleboro Memorial Hospital, VT.)

29. Kavlock R, Dix D. 2010. Computational toxicology as implemented by the U.S. EPA: providing high throughput decision support tools for screening and assessing chemical exposure, hazard and risk. *J Toxicol Environ Health B Crit Rev* 13(2-4): 197-217. (Support not reported. Authors affiliated with US EPA, NC.)
30. Kavlock R, Chandler K, Houck K, Hunter S, Judson R, Kleinstreuer N, Knudsen T, Martin M, Padilla S, Reif D, Richard A, Rotroff D, Sipes N, Dix D. 2012. Update on EPA's ToxCast program: providing high throughput decision support tools for chemical risk management. *Chem Res Toxicol* 25(7): 1287-1302. (Support not reported. Authors affiliated with US EPA, NC.)
31. Kawata K, Yokoo H, Shimazaki R, Okabe S. 2007. Classification of heavy-metal toxicity by human DNA microarray analysis. *Environ Sci Technol* 41(10): 3769-3774. (Supported by the 21st Century COE Program "Sustainable Metabolic System of Water and Waste for Area-Based Society," and the Ministry of Education, Science, and Culture of Japan. Authors affiliated with Hokkaido University, Japan.)
32. Kirkland D, Whitwell J, Deyo J, Serex T. 2007. Failure of antimony trioxide to induce micronuclei or chromosomal aberrations in rat bone-marrow after sub-chronic oral dosing. *Mutat Res* 627(2): 119-128. (Supported by the International Antimony Oxide Industry Association (IAOIA) and the Antimony Trioxide Stakeholders (ATOS). Authors affiliated with Covance Laboratories Ltd., UK; Eastman Chemical Company, TN; Blasland, Bouck & Lee Inc., CA.)
33. Kocyigit A, Gur S, Gurel MS, Bulut V, Ulukanligil M. 2002. Antimonial therapy induces circulating proinflammatory cytokines in patients with cutaneous leishmaniasis. *Infect Immun* 70(12): 6589-6591. (Support not reported. Authors affiliated with University of Harran Medical School, Turkey; University of Firat Medical School, Turkey.)
34. Kuroda K, Endo G, Okamoto A, Yoo YS, Horiguchi S. 1991. Genotoxicity of beryllium, gallium and antimony in short-term assays. *Mutat Res* 264(4): 163-170. (Supported by the Ministry of Education, Science and Culture of Japan. Authors affiliated with Osaka City Institute of Public Health and Environmental Sciences, Japan; Osaka City University Medical School, Japan; Seoul Health Junior College, Korea.)
35. Lantzsch H, Gebel T. 1997. Genotoxicity of selected metal compounds in the SOS chromotest. *Mutat Res* 389(2-3): 191-197. (Support not reported. Authors affiliated with University of Goettingen, Germany.)
36. Lecureur V, Lagadic-Gossmann D, Fardel O. 2002. Potassium antimonyl tartrate induces reactive oxygen species-related apoptosis in human myeloid leukemic HL60 cells. *Int J Oncol* 20(5): 1071-1076. (Support not reported. Authors affiliated with INSERM, France.)
37. Lösler S, Schlieff S, Kneifel C, Thiel E, Schrezenmeier H, Rojewski MT. 2009. Antimony-trioxide- and arsenic-trioxide-induced apoptosis in myelogenic and lymphatic cell lines, recruitment of caspases, and loss of mitochondrial membrane potential are enhanced by modulators of the cellular glutathione redox system. *Ann Hematol* 88(11): 1047-1058. (Support not reported. Authors affiliated with Charité Universitätsmedizin Berlin,

- Germany; Universität Ulm, Germany; Institut für klinische Transfusionsmedizin und Immungenetik, Germany.)
38. Mann KK, Davison K, Colombo M, Colosimo AL, Diaz Z, Padovani AM, Guo Q, Scrivens PJ, Gao W, Mader S, Miller WH, Jr. 2006. Antimony trioxide-induced apoptosis is dependent on SEK1/JNK signaling. *Toxicol Lett* 160(2): 158-170. (Supported by the CIHR, the Montreal Centre for Experimental Therapeutics in Cancer/CIHR/FRSQ, BCRP, Department of the Army, and the U.S. Army Medical Research Acquisition Activity. Authors affiliated with Montreal Centre for Experimental Therapeutics in Cancer and Lady Davis Institute for Medical Research, Canada; Université de Montréal, Canada.)
 39. Migliore L, Cocchi L, Nesti C, Sabbioni E. 1999. Micronuclei assay and FISH analysis in human lymphocytes treated with six metal salts. *Environ Mol Mutagen* 34(4): 279-284. (Support not reported. Authors affiliated with Università di Pisa, Italy; ECVAM, Italy.)
 40. Murray HW, Delph-Etienne S. 2000. Roles of endogenous gamma interferon and macrophage microbicidal mechanisms in host response to chemotherapy in experimental visceral leishmaniasis. *Infect Immun* 68(1): 288-293. (Supported by NIH. Authors affiliated with Weill Medical College of Cornell University, NY.)
 41. NCBI. 2017. *GEO Datasets*. National Center for Biotechnology Information, U.S. National Library of Medicine. <https://www.ncbi.nlm.nih.gov/gds/>. Accessed on 11/20/17.
 42. Newton PE, Bolte HF, Daly IW, Pillsbury BD, Terrill JB, Drew RT, Ben-Dyke R, Sheldon AW, Rubin LF. 1994. Subchronic and chronic inhalation toxicity of antimony trioxide in the rat. *Fundam Appl Toxicol* 22(4): 561-576. (Supported by the Antimony Oxide Industry Association. Authors affiliated with Pharmaco LSR Inc., NJ; American Petroleum Institute, Washington, DC; Antimony Oxide Industry Association, Washington, DC; University of Pennsylvania, PA.)
 43. Nielson KB, Atkin CL, Winge DR. 1985. Distinct metal-binding configurations in metallothionein. *J Biol Chem* 260(9): 5342-5350. (Supported by NIH. Authors affiliated with University of Utah, UT.)
 44. Nishioka H. 1975. Mutagenic activities of metal compounds in bacteria. *Mutat Res* 31(3): 185-189. (Support not reported. Author affiliated with Doshisha University, Japan.)
 45. NTP. 2017a. *Tox21 Toolbox*. U.S. Department of Health and Human Services. <https://ntp.niehs.nih.gov/results/tox21/tbox/>. Accessed on 10/10/17.
 46. NTP. 2017b. *Toxicology and Carcinogenesis Studies of Antimony Trioxide (CAS No. 1309-64-4) in Wistar Han [CrI:WI (Han)] Rats and B6C3F1/N Mice (Inhalation Studies)*. NTP TR 590. National Toxicology Program, National Institutes of Health. 265 pp. https://ntp.niehs.nih.gov/ntp/htdocs/lt_rpts/tr590_508.pdf.
 47. Pathak MK, Yi T. 2001. Sodium stibogluconate is a potent inhibitor of protein tyrosine phosphatases and augments cytokine responses in hemopoietic cell lines. *J Immunol* 167(6): 3391-3397. (Supported in part by Grants R01CA79891 and R01MG58893. Authors affiliated with Cleveland Clinic Foundation, OH.)

48. Paton GR, Allison AC. 1972. Chromosome damage in human cell cultures induced by metal salts. *Mutat Res* 16(3): 332-336. (Support not reported. Authors affiliated with Clinical Research Centre, UK.)
49. Schaumlöffel N, Gebel T. 1998. Heterogeneity of the DNA damage provoked by antimony and arsenic. *Mutagenesis* 13(3): 281-286. (Support not reported. Authors affiliated with University of Gottingen, Germany.)
50. Schnorr TM, Steenland K, Thun MJ, Rinsky RA. 1995. Mortality in a cohort of antimony smelter workers. *Am J Ind Med* 27(5): 759-770. (Support not reported. Authors affiliated with NIOSH, OH; American Cancer Society, GA.)
51. Schroeder HA, Mitchener M, Balassa JJ, Kanisawa M, Nason AP. 1968. Zirconium, niobium, antimony and fluorine in mice: effects on growth, survival and tissue levels. *J Nutr* 95(1): 95-101. (Supported by the Public Health Service, the National Heart Institute, the US Army and CIBA Pharmaceutical Products, Inc. Authors affiliated with Dartmouth Medical School, NH; Brattleboro Memorial Hospital, VT.)
52. Schroeder HA, Mitchener M, Nason AP. 1970. Zirconium, niobium, antimony, vanadium and lead in rats: life term studies. *J Nutr* 100(1): 59-68. (Supported by the Public Health Service, the National Heart Institute, the U.S. Army, and Cooper Laboratories, Inc. Authors affiliated with Dartmouth Medical School, NH; Brattleboro Memorial Hospital, VT.)
53. Smith MT, Guyton KZ, Gibbons CF, Fritz JM, Portier CJ, Rusyn I, DeMarini DM, Caldwell JC, Kavlock RJ, Lambert PF, Hecht SS, Bucher JR, Stewart BW, Baan RA, Coglianò VJ, Straif K. 2016. Key characteristics of carcinogens as a basis for organizing data on mechanisms of carcinogenesis. *Environ Health Perspect* 124(6): 713-721. (Supported by NIEHS. Authors affiliated with University of California Berkeley, CA; IARC, France; US EPA, Washington, D.C. and NC; ATSDR; Texas A&M University, TX; University of Wisconsin School of Medicine and Public Health, WI; University of Minnesota, MN; NIEHS, NC; University of New South Wales, Australia.)
54. Tice RR, Austin CP, Kavlock RJ, Bucher JR. 2013. Improving the human hazard characterization of chemicals: a Tox21 update. *Environ Health Perspect* 121(7): 756-765. (Support not reported. Authors affiliated with NIH, NC and MD; US EPA, NC.)
55. TNO Quality of Life. 2005. A study on the biodistribution of antimony trioxide (Sb₂O₃) in rats. *TNO Rep* 6502: 1-63. (as cited in EU 2008)
56. Wang B, Yu W, Guo J, Jiang X, Lu W, Liu M, Pang X. 2015. The antiparasitic drug, potassium antimony tartrate, inhibits tumor angiogenesis and tumor growth in nonsmall-cell lung cancer. *J Pharmacol Exp Ther* 352(1): 129-138. (Supported by the National Basic Research Program of China, the National Natural Science Foundation of China, the Science and Technology Commission of Shanghai Municipality, the Chenguang Program from Shanghai Municipal Education Commission, and the Fundamental Research Funds for the Central Universities. Authors affiliated with East China Normal University, China; Texas A&M University Health Science Center, TX.)

57. Watt WD. 1983. *Chronic Inhalation Toxicity of Antimony Trioxide: Validation of the Threshold Limit Value*. Detroit, MI: Wayne State University, PhD Thesis. (Supported by ASARCO, Inc. Author affiliated with Wayne State University, MI.)
58. Westrick ML. 1953. Physiologic responses attending administration of antimony, alone or with simultaneous injections of thyroxin. *Proc Soc Exp Biol Med* 82(1): 56-60. (Support not reported. Author affiliated with University of Pittsburgh, PA.)
59. Wingren G, Axelson O. 1993. Epidemiologic studies of occupational cancer as related to complex mixtures of trace elements in the art glass industry. *Scand J Work Environ Health* 19 Suppl 1: 95-100. (Support not reported. Authors affiliated with University Hospital - Linköping, Sweden)
60. Wortmann GW, Aronson NE, Byrd JC, Grever MR, Oster CN. 1998. Herpes zoster and lymphopenia associated with sodium stibogluconate therapy for cutaneous leishmaniasis. *Clin Infect Dis* 27(3): 509-512. (Support not reported. Authors affiliated with Walter Reed Army Medical Center, Washington, D.C; Johns Hopkins University, MD.)
61. Yamamoto A, Kohyama Y, Hanawa T. 2002. Mutagenicity evaluation of forty-one metal salts by the umu test. *J Biomed Mater Res* 59(1): 176-183. (Support not reported. Authors affiliated with National Institute for Materials Science, Japan.)
62. Yi T, Pathak MK, Lindner DJ, Ketterer ME, Farver C, Borden EC. 2002. Anticancer activity of sodium stibogluconate in synergy with IFNs. *J Immunol* 169(10): 5978-5985. (Supported by Grants R01CA7989, R01 MG58893 and CA90914. Authors affiliated with Cleveland Clinic Foundation, OH; Northern Arizona University Flagstaff, AZ.)