

Overview of the NTP Rat Model Selection: Technical Reports 583-586

Angela King-Herbert, DVM, DACLAM
National Institute of Environmental Health Sciences

NTP Technical Reports Peer Review
May 22, 2014

Objective

To provide an overview of the rat models used in the NTP studies presented today.

NTP F344/N Rat

- The F344/N rat had several characteristics that made it a good choice for the 2-year rodent bioassay.
 - Used in the rodent bioassay for about 30 years.
 - The historical database was based on a 5-year rolling window and was very robust.
 - Relatively small as compared to other rat strains and stocks.
 - Good survival rate at the end of a 2-year study.
 - Good fecundity for an inbred rat strain, with approximately 6-8 pups produced per litter.

NTP F344/N Rat

- There were several concerns with the F344/N rat as well.
 - High background incidence of:
 - Testicular tumors (interstitial cell tumor)
 - Mononuclear cell leukemia
 - The F344/N rat developed declining fertility, sporadic idiopathic seizures, and spontaneous chylothorax without evidence of trauma.
 - These issues were unique to the NTP F344/N colony.
 - Possibility of genetic drift within the colony.

Strains & Stocks Workshop

- NTP hosted a workshop in June 2005, “Animal Models for the NTP Rodent Cancer Bioassay: Stocks & Strains—Should We Switch?”
 - One of the objectives of the workshop was to determine if the F344/N rat was still an appropriate model in identifying substances that may be a carcinogenic hazard for humans.
 - Included an invited panel of scientists with expertise in rodent genetics, cancer biology, statistics and other related fields.
 - Breakout groups discussed this rodent model and its place in the NTP bioassay.

Strains & Stocks Workshop Summary

- The rat breakout group recommended discontinuing the use of the current F344/N rat strain
 - Proposed 3 options:
 - Re-establish the F344/N strain from another source, however the general concerns about the model would still be present.
 - Create an F1 hybrid such as the F344 X Brown Norway cross, FBNF1.
 - Consider using an alternative model such as an outbred rat like the Wistar Han or Sprague Dawley.

NTP Deliberations

- The NTP discontinued use of the F344/N rat in all new studies and temporarily used the F344/NTac rat from Taconic Farms, Inc.'s commercial colony.
- Scientists within the NTP deliberated the selection of an alternative rat model.



Other programmatic changes

- The NTP decided to utilize a single rat model for studies of other endpoints.
 - Minimized the need to conduct multiple preliminary and toxicokinetic studies.
 - Enhanced comparability across study endpoints.
- The NTP was moving towards a perinatal exposure design.
 - Some toxicity/carcinogenicity studies would include *in utero* exposure in rat models.
 - More developmental/reproductive toxicology studies to be performed.
 - RACB, MOG and teratology studies
 - Inbred strains were not a good choice for perinatal exposure but outbred stocks were.

General considerations when selecting animal models

Primary

- Availability
- Fecundity
- Survival
- Sensitivity to carcinogens
- Spontaneous tumor rate

Other

- Experience with the model
- Similar metabolic pathways to humans
- Similar pathology to humans
- Sensitivity to other endpoints
- Cost



NTP rat model selection

- Desired traits
 - Outbred
 - Long lifespan
 - Moderate size for an outbred rat
 - Good fertility with large litters
- 2007
 - Selected the Wistar Han model [CrI:WI(Han)] but discontinued use due to several issues
 - Low pregnancy rate
 - Litter size smaller than expected
 - Sex ratio in litters appeared skewed
- 2009
 - Changed the rat model to the Harlan Sprague Dawley [HSD:Sprague Dawley (SD)]
 - Currently use this model

Brief overview on rodent models - Alike but different

F344/N

≠

F344/NTac

N=NIH

- F344/N – NIH rat strain
- F344/NTac – Substrain of the NIH F344 rat; bred at Taconic Farms, Inc.
- In general, historical databases are not combined between substrains

Summary of rat models used in the studies presented today

	2-week study	3-month study	2-year study
Bromodichloroacetic Acid TR 583	F344/N	F344/N	F344/NTac
Indole-3-Carbinol TR 584	N/A	F344/N	Harlan Sprague Dawley
Green Tea Extract TR 585	N/A	F344/NTac	Wistar Han
Cimstar 3800 TR 586	N/A	F344/NTac	Wistar Han

Questions?

