

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 238
 Report Date: 03/22/2010

Study Summary

| Laboratory Name | Study Number | Study Start Date | Length of Study in Days | Number of Animals Total and (#/Cage) | Number Surviving and (Mean Life Span) | Maximum Mean Weekly Body Weight | Chemical |
|-----------------------------|--------------|------------------|-------------------------|--------------------------------------|---------------------------------------|---------------------------------|--|
| Male | | | | | | | |
| Southern Research Institute | 052011406 | 09/04/02 | 729 | 50(1) | 33(700) | 55.6 | Sodium dichromate dihydrate (VI) |
| Southern Research Institute | 059601906 | 10/10/01 | 729 | 50(1) | 38(696) | 54.1 | Water disinfection byproducts (Bromochloroacetic acid) |
| Southern Research Institute | 059601506 | 08/22/01 | 729 | 50(1) | 40(697) | 54.0 | Water disinfection byproducts (Dibromoacetonitrile) |
| Female | | | | | | | |
| Southern Research Institute | 052011406 | 09/04/02 | 735 | 50(5) | 35(701) | 64.3 | Sodium dichromate dihydrate (VI) |
| Southern Research Institute | 059601906 | 10/10/01 | 734 | 50(5) | 36(693) | 63.0 | Water disinfection byproducts (Bromochloroacetic acid) |
| Southern Research Institute | 059601506 | 08/22/01 | 731 | 50(5) | 35(705) | 60.9 | Water disinfection byproducts (Dibromoacetonitrile) |

NOTE: The text on page 2 ("Report Description") describes how the studies listed here are displayed on the data pages that follow.

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 239
 Report Date: 03/22/2010

| | | Male | | | Female | | | |
|--------------------------------------|-------|------------------|-------------|-------------|-------------|-----------------|-------------|-----------|
| #All Organs: | | | | | | | | |
| Benign Tumors | | | | | | | | |
| | | 42/50 (84%) | 41/50 (82%) | 32/50 (64%) | 27/50 (54%) | 32/50 (64%) | 29/50 (58%) | |
| Overall Incidence | Total | 115/150 (76.67%) | Mean 76.67% | SD 11.02% | Total | 88/150 (58.67%) | Mean 58.67% | SD 5.03% |
| | | | | | | | | |
| #All Organs: | | | | | | | | |
| Malignant Tumors | | | | | | | | |
| | | 31/50 (62%) | 36/50 (72%) | 29/50 (58%) | 22/50 (44%) | 38/50 (76%) | 31/50 (62%) | |
| Overall Incidence | Total | 96/150 (64%) | Mean 64% | SD 7.21% | Total | 91/150 (60.67%) | Mean 60.67% | SD 16.04% |
| | | | | | | | | |
| #All Organs: | | | | | | | | |
| Malignant and Benign Tumors | | | | | | | | |
| | | 49/50 (98%) | 47/50 (94%) | 41/50 (82%) | 36/50 (72%) | 43/50 (86%) | 44/50 (88%) | |
| Overall Incidence | Total | 137/150 (91.33%) | Mean 91.33% | SD 8.33% | Total | 123/150 (82%) | Mean 82% | SD 8.72% |
| | | | | | | | | |
| #All Organs: | | | | | | | | |
| Hemangioma | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 1/50 (2%) | 1/50 (2%) | 2/50 (4%) | 0/50 (0%) | |
| Overall Incidence | Total | 2/150 (1.33%) | Mean 1.33% | SD 1.15% | Total | 3/150 (2%) | Mean 2% | SD 2% |
| | | | | | | | | |
| #All Organs: | | | | | | | | |
| Hemangiosarcoma | | | | | | | | |
| | | 4/50 (8%) | 3/50 (6%) | 1/50 (2%) | 2/50 (4%) | 1/50 (2%) | 3/50 (6%) | |
| Overall Incidence | Total | 8/150 (5.33%) | Mean 5.33% | SD 3.06% | Total | 6/150 (4%) | Mean 4% | SD 2% |
| | | | | | | | | |
| #All Organs: | | | | | | | | |
| Hemangiosarcoma or Hemangioma | | | | | | | | |
| | | 4/50 (8%) | 4/50 (8%) | 2/50 (4%) | 3/50 (6%) | 3/50 (6%) | 3/50 (6%) | |
| Overall Incidence | Total | 10/150 (6.67%) | Mean 6.67% | SD 2.31% | Total | 9/150 (6%) | Mean 6% | SD 0% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 240
 Report Date: 03/22/2010

| | | Male | | | Female | | |
|--|-------|---------------|------------|-----------|-----------------------|-------------|------------|
| #All Organs: | | | | | | | |
| Histiocytic Sarcoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 2/50 (4%) | 1/50 (2%) | 4/50 (8%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total 7/150 (4.67%) | Mean 4.67% | SD 3.06% |
| | | | | | | | |
| #All Organs: | | | | | | | |
| Malignant Lymphoma: Histiocytic, Lymphocytic, Mixed, NOS, or Undifferentiated Cell Type | | | | | | | |
| | | 2/50 (4%) | 2/50 (4%) | 0/50 (0%) | 7/50 (14%) | 18/50 (36%) | 9/50 (18%) |
| Overall Incidence | Total | 4/150 (2.67%) | Mean 2.67% | SD 2.31% | Total 34/150 (22.67%) | Mean 22.67% | SD 11.72% |
| | | | | | | | |
| #All Organs: | | | | | | | |
| Mesothelioma: Benign, Malignant, NOS | | | | | | | |
| | | 2/50 (4%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 2/150 (1.33%) | Mean 1.33% | SD 2.31% | Total 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | |
| #All Organs: | | | | | | | |
| Mesothelioma: Malignant | | | | | | | |
| | | 2/50 (4%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 2/150 (1.33%) | Mean 1.33% | SD 2.31% | Total 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | |
| #All Organs: | | | | | | | |
| Osteoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total 1/150 (0.67%) | Mean 0.67% | SD 1.15% |
| | | | | | | | |
| #All Organs: | | | | | | | |
| Osteosarcoma | | | | | | | |
| | | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total 0/150 (0%) | Mean 0% | SD 0% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 241
 Report Date: 03/22/2010

| | | Male | | | Female | | |
|------------------------------------|-------|-----------------|-------------|------------|-----------|---------------|------------------------|
| #All Organs: | | | | | | | |
| Osteosarcoma or Osteoma | | | | | | | |
| | | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 1/150 (0.67%) | Mean 0.67% SD 1.15% |
| | | | | | | | |
| *Adrenal Cortex: | | | | | | | |
| Adenoma | | | | | | | |
| | | 7/49 (14%) | 9/50 (18%) | 6/50 (12%) | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 22/149 (14.77%) | Mean 14.76% | SD 3.03% | Total | 0/149 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Adrenal Cortex: | | | | | | | |
| Plasma Cell Tumor Malignant | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 0/149 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Adrenal Medulla: | | | | | | | |
| Pheochromocytoma Benign | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 1/49 (2%) | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 1/148 (0.68%) | Mean 0.68% | SD 1.18% | Total | 0/149 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Adrenal Medulla: | | | | | | | |
| Pheochromocytoma Complex | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/49 (0%) | 0/49 (0%) | 1/50 (2%) | 0/50 (0%) |
| Overall Incidence | Total | 0/148 (0%) | Mean 0% | SD 0% | Total | 1/149 (0.67%) | Mean 0.67% SD 1.15% |
| | | | | | | | |
| *Adrenal Medulla: | | | | | | | |
| Pheochromocytoma Malignant | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/49 (0%) | 0/49 (0%) | 1/50 (2%) | 0/50 (0%) |
| Overall Incidence | Total | 0/148 (0%) | Mean 0% | SD 0% | Total | 1/149 (0.67%) | Mean 0.67% SD 1.15% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 242
 Report Date: 03/22/2010

| | | Male | | | Female | | | |
|--|-------|---------------|------------|-----------|-----------|---------------|------------|----------|
| *Adrenal Medulla: | | | | | | | | |
| Pheochromocytoma: Benign, Complex, Malignant, NOS | | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 1/49 (2%) | 0/49 (0%) | 2/50 (4%) | 0/50 (0%) | |
| Overall Incidence | Total | 1/148 (0.68%) | Mean 0.68% | SD 1.18% | Total | 2/149 (1.34%) | Mean 1.33% | SD 2.31% |
| | | | | | | | | |
| *Bone Marrow: | | | | | | | | |
| Hemangioma | | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Bone Marrow: | | | | | | | | |
| Hemangiosarcoma | | | | | | | | |
| | | 1/49 (2%) | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 2/149 (1.34%) | Mean 1.35% | SD 1.17% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Bone Marrow: | | | | | | | | |
| Plasma Cell Tumor Malignant | | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Bone: | | | | | | | | |
| Hemangiosarcoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Bone: | | | | | | | | |
| Osteoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% |
| | | | | | | | | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 243
 Report Date: 03/22/2010

| | | Male | | | | Female | | | | | |
|---|-------|---------------|------|------------|------|----------|------|-------|------|---------------|------|
| #Bone: | | | | | | | | | | | |
| Osteosarcoma | | | | | | | | | | | |
| | | 1/50 | (2%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 1/150 (0.67%) | | Mean 0.67% | | SD 1.15% | | Total | | 0/150 (0%) | |
| | | | | Mean 0% | | | | | | SD 0% | |
| | | | | | | | | | | | |
| #Bone: | | | | | | | | | | | |
| Osteosarcoma or Osteoma | | | | | | | | | | | |
| | | 1/50 | (2%) | 0/50 | (0%) | 0/50 | (0%) | 1/50 | (2%) | 0/50 | (0%) |
| Overall Incidence | Total | 1/150 (0.67%) | | Mean 0.67% | | SD 1.15% | | Total | | 1/150 (0.67%) | |
| | | | | Mean 0.67% | | | | | | SD 1.15% | |
| | | | | | | | | | | | |
| *Brain: | | | | | | | | | | | |
| Meningioma Benign | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 (0%) | | Mean 0% | | SD 0% | | Total | | 0/150 (0%) | |
| | | | | Mean 0% | | | | | | SD 0% | |
| | | | | | | | | | | | |
| *Brain: | | | | | | | | | | | |
| Meningioma Malignant | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 (0%) | | Mean 0% | | SD 0% | | Total | | 0/150 (0%) | |
| | | | | Mean 0% | | | | | | SD 0% | |
| | | | | | | | | | | | |
| *Brain: | | | | | | | | | | | |
| Meningioma: Benign, Malignant, NOS | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 (0%) | | Mean 0% | | SD 0% | | Total | | 0/150 (0%) | |
| | | | | Mean 0% | | | | | | SD 0% | |
| | | | | | | | | | | | |
| *Brain: | | | | | | | | | | | |
| Schwannoma Malignant | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 1/50 | (2%) |
| Overall Incidence | Total | 0/150 (0%) | | Mean 0% | | SD 0% | | Total | | 1/150 (0.67%) | |
| | | | | Mean 0% | | | | | | Mean 0.67% | |
| | | | | | | | | | | SD 1.15% | |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 244
 Report Date: 03/22/2010

| | | Male | | | Female | | |
|---|-------|---------------|------------|-----------|-----------|------------|------------------|
| *Clitoral/Preputial Gland: Adenoma | | | | | | | |
| | | 0/50 (0%) | 0/48 (0%) | 1/49 (2%) | 0/50 (0%) | 0/43 (0%) | 0/47 (0%) |
| Overall Incidence | Total | 1/147 (0.68%) | Mean 0.68% | SD 1.18% | Total | 0/140 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Clitoral/Preputial Gland: Carcinoma | | | | | | | |
| | | 0/50 (0%) | 0/48 (0%) | 0/49 (0%) | 0/50 (0%) | 0/43 (0%) | 0/47 (0%) |
| Overall Incidence | Total | 0/147 (0%) | Mean 0% | SD 0% | Total | 0/140 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Clitoral/Preputial Gland: Carcinoma or Adenoma | | | | | | | |
| | | 0/50 (0%) | 0/48 (0%) | 1/49 (2%) | 0/50 (0%) | 0/43 (0%) | 0/47 (0%) |
| Overall Incidence | Total | 1/147 (0.68%) | Mean 0.68% | SD 1.18% | Total | 0/140 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Clitoral/Preputial Gland: Hemangioma | | | | | | | |
| | | 0/50 (0%) | 0/48 (0%) | 0/49 (0%) | 0/50 (0%) | 0/43 (0%) | 0/47 (0%) |
| Overall Incidence | Total | 0/147 (0%) | Mean 0% | SD 0% | Total | 0/140 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Clitoral/Preputial Gland: Hemangiosarcoma | | | | | | | |
| | | 0/50 (0%) | 0/48 (0%) | 0/49 (0%) | 0/50 (0%) | 0/43 (0%) | 0/47 (0%) |
| Overall Incidence | Total | 0/147 (0%) | Mean 0% | SD 0% | Total | 0/140 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Clitoral/Preputial Gland: Squamous Cell Carcinoma | | | | | | | |
| | | 0/50 (0%) | 0/48 (0%) | 0/49 (0%) | 0/50 (0%) | 0/43 (0%) | 0/47 (0%) |
| Overall Incidence | Total | 0/147 (0%) | Mean 0% | SD 0% | Total | 0/140 (0%) | Mean 0% SD 0% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 245
 Report Date: 03/22/2010

| | | Male | | | Female | | | |
|--------------------------------|-------|-----------------|-------------|------------|------------|-----------------|-------------|----------|
| #Esophagus: | | | | | | | | |
| Squamous Cell Carcinoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Eye: | | | | | | | | |
| Sarcoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Harderian Gland: | | | | | | | | |
| Adenoma | | | | | | | | |
| | | 6/50 (12%) | 5/50 (10%) | 5/50 (10%) | 4/50 (8%) | 1/50 (2%) | 6/50 (12%) | |
| Overall Incidence | Total | 16/150 (10.67%) | Mean 10.67% | SD 1.15% | Total | 11/150 (7.33%) | Mean 7.33% | SD 5.03% |
| | | | | | | | | |
| #Harderian Gland: | | | | | | | | |
| Carcinoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 1/50 (2%) | 2/50 (4%) | 2/50 (4%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 5/150 (3.33%) | Mean 3.33% | SD 1.15% |
| | | | | | | | | |
| #Harderian Gland: | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | |
| | | 6/50 (12%) | 5/50 (10%) | 5/50 (10%) | 5/50 (10%) | 3/50 (6%) | 8/50 (16%) | |
| Overall Incidence | Total | 16/150 (10.67%) | Mean 10.67% | SD 1.15% | Total | 16/150 (10.67%) | Mean 10.67% | SD 5.03% |
| | | | | | | | | |
| #Harderian Gland: | | | | | | | | |
| Sarcoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 246
 Report Date: 03/22/2010

| | | Male | | | Female | | |
|--|-------|---------------|------------|-----------|-----------|------------|------------------|
| *Heart: | | | | | | | |
| Hemangioma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Heart: | | | | | | | |
| Hemangiosarcoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Intestine Large: Cecum: | | | | | | | |
| Carcinoma | | | | | | | |
| | | 1/50 (2%) | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 2/150 (1.33%) | Mean 1.33% | SD 1.15% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Intestine Large: Cecum: | | | | | | | |
| Carcinoma or Adenoma | | | | | | | |
| | | 1/50 (2%) | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 2/150 (1.33%) | Mean 1.33% | SD 1.15% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Intestine Large: Cecum: | | | | | | | |
| Leiomyoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Intestine Large: Colon/Rectum: | | | | | | | |
| Leiomyosarcoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 247
 Report Date: 03/22/2010

| | | Male | | | Female | | | |
|------------------------------------|-------|---------------|------------|-----------|-----------|---------------|------------|----------|
| #Intestine Small: Duodenum: | | | | | | | | |
| Adenoma | | | | | | | | |
| | | 1/50 (2%) | 3/50 (6%) | 1/50 (2%) | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | |
| Overall Incidence | Total | 5/150 (3.33%) | Mean 3.33% | SD 2.31% | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% |
| | | | | | | | | |
| #Intestine Small: Duodenum: | | | | | | | | |
| Carcinoma | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Intestine Small: Duodenum: | | | | | | | | |
| Leiomyosarcoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Intestine Small: Duodenum: | | | | | | | | |
| Polyp Adenomatous | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Intestine Small: Ileum: | | | | | | | | |
| Carcinoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Intestine Small: Ileum: | | | | | | | | |
| Hemangiosarcoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 248
 Report Date: 03/22/2010

| | | Male | | | | Female | | | |
|--|-------|---------------|------------|-----------|-------|---------------|------------|-----------|--|
| #Intestine Small: Ileum: | | | | | | | | | |
| Polyp Adenomatous | | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Intestine Small: Jejunum: | | | | | | | | | |
| Adenoma | | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Intestine Small: Jejunum: | | | | | | | | | |
| Carcinoma | | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 1/50 (2%) | | 1/50 (2%) | 1/50 (2%) | 0/50 (0%) | |
| Overall Incidence | Total | 2/150 (1.33%) | Mean 1.33% | SD 1.15% | Total | 2/150 (1.33%) | Mean 1.33% | SD 1.15% | |
| | | | | | | | | | |
| #Intestine Small: Jejunum: | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Intestine Small: Jejunum: | | | | | | | | | |
| Leiomyosarcoma | | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Intestine Small: Site Unspecified: | | | | | | | | | |
| Adenoma | | | | | | | | | |
| | | 1/50 (2%) | 3/50 (6%) | 1/50 (2%) | | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | |
| Overall Incidence | Total | 5/150 (3.33%) | Mean 3.33% | SD 2.31% | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 249
 Report Date: 03/22/2010

| | | Male | | | Female | | |
|---|-------|---------------|------------|-----------|-----------|---------------|------------------------|
| #Intestine Small: Site Unspecified: Carcinoma | | | | | | | |
| | | 0/50 (0%) | 2/50 (4%) | 1/50 (2%) | 1/50 (2%) | 1/50 (2%) | 0/50 (0%) |
| Overall Incidence | Total | 3/150 (2%) | Mean 2% | SD 2% | Total | 2/150 (1.33%) | Mean 1.33% SD 1.15% |
| | | | | | | | |
| #Intestine Small: Site Unspecified: Carcinoma or Adenoma | | | | | | | |
| | | 1/50 (2%) | 5/50 (10%) | 1/50 (2%) | 1/50 (2%) | 2/50 (4%) | 0/50 (0%) |
| Overall Incidence | Total | 7/150 (4.67%) | Mean 4.67% | SD 4.62% | Total | 3/150 (2%) | Mean 2% SD 2% |
| | | | | | | | |
| #Intestine Small: Site Unspecified: Hemangiosarcoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Intestine Small: Site Unspecified: Leiomyosarcoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Intestine Small: Site Unspecified: Polyp Adenomatous | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Islets, Pancreatic: Adenoma | | | | | | | |
| | | 3/49 (6%) | 1/50 (2%) | 1/50 (2%) | 1/49 (2%) | 1/49 (2%) | 1/49 (2%) |
| Overall Incidence | Total | 5/149 (3.36%) | Mean 3.37% | SD 2.38% | Total | 3/147 (2.04%) | Mean 2.04% SD 1.1% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 250
 Report Date: 03/22/2010

| | | Male | | | | Female | | | |
|-------------------------------|-------|---------------|------------|-----------|-------|---------------|------------|-----------|--|
| *Islets, Pancreatic: | | | | | | | | | |
| Carcinoma | | | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/49 (0%) | 0/49 (0%) | 0/49 (0%) | |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 0/147 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| *Islets, Pancreatic: | | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | | |
| | | 3/49 (6%) | 1/50 (2%) | 1/50 (2%) | | 1/49 (2%) | 1/49 (2%) | 1/49 (2%) | |
| Overall Incidence | Total | 5/149 (3.36%) | Mean 3.37% | SD 2.38% | Total | 3/147 (2.04%) | Mean 2.04% | SD 1.5% | |
| | | | | | | | | | |
| *Kidney: Renal Tubule: | | | | | | | | | |
| Adenoma | | | | | | | | | |
| | | 0/50 (0%) | 4/50 (8%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 4/150 (2.67%) | Mean 2.67% | SD 4.62% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| *Kidney: Renal Tubule: | | | | | | | | | |
| Carcinoma | | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| *Kidney: Renal Tubule: | | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | | |
| | | 0/50 (0%) | 4/50 (8%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 4/150 (2.67%) | Mean 2.67% | SD 4.62% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| *Kidney: Renal Tubule: | | | | | | | | | |
| Hemangioma | | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 251
 Report Date: 03/22/2010

| | | Male | | | | Female | | | |
|---|-------|----------------------------|-------------------------|-----------------------|-------|----------------------------|-------------------------|-----------------------|--|
| *Kidney: Renal Tubule: Hemangiosarcoma | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | |
| *Kidney: Renal Tubule: Plasma Cell Tumor Malignant | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | |
| *Kidney: Renal Tubule: Sarcoma | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | |
| *Liver: Fibrous Histiocytoma | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/49 (0%) 0/149 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | |
| *Liver: Hemangioma | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 1/150 (0.67%) | 0/50 (0%) Mean 0.67% | 1/50 (2%) SD 1.15% | Total | 1/49 (2%) 1/149 (0.67%) | 0/50 (0%) Mean 0.68% | 0/50 (0%) SD 1.18% | |
| *Liver: Hemangiosarcoma | | | | | | | | | |
| Overall Incidence | Total | 1/50 (2%) 3/150 (2%) | 2/50 (4%) Mean 2% | 0/50 (0%) SD 2% | Total | 1/49 (2%) 3/149 (2.01%) | 0/50 (0%) Mean 2.01% | 2/50 (4%) SD 2% | |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 252
 Report Date: 03/22/2010

| | | Male | | | Female | | |
|--|-------|------------------|-------------|-------------|-------------|-----------------|--------------------------|
| *Liver: | | | | | | | |
| Hepatoblastoma | | | | | | | |
| | | 17/50 (34%) | 4/50 (8%) | 1/50 (2%) | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 22/150 (14.67%) | Mean 14.67% | SD 17.01% | Total | 0/149 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Liver: | | | | | | | |
| Hepatocellular Adenoma | | | | | | | |
| | | 36/50 (72%) | 27/50 (54%) | 29/50 (58%) | 14/49 (29%) | 27/50 (54%) | 19/50 (38%) |
| Overall Incidence | Total | 92/150 (61.33%) | Mean 61.33% | SD 9.45% | Total | 60/149 (40.27%) | Mean 40.19% SD 12.86% |
| | | | | | | | |
| *Liver: | | | | | | | |
| Hepatocellular Carcinoma | | | | | | | |
| | | 14/50 (28%) | 19/50 (38%) | 24/50 (48%) | 8/49 (16%) | 14/50 (28%) | 10/50 (20%) |
| Overall Incidence | Total | 57/150 (38%) | Mean 38% | SD 10% | Total | 32/149 (21.48%) | Mean 21.44% SD 5.97% |
| | | | | | | | |
| *Liver: | | | | | | | |
| Hepatocellular Carcinoma or Hepatoblastoma | | | | | | | |
| | | 25/50 (50%) | 21/50 (42%) | 24/50 (48%) | 8/49 (16%) | 14/50 (28%) | 10/50 (20%) |
| Overall Incidence | Total | 70/150 (46.67%) | Mean 46.67% | SD 4.16% | Total | 32/149 (21.48%) | Mean 21.44% SD 5.97% |
| | | | | | | | |
| *Liver: | | | | | | | |
| Hepatocellular Carcinoma or Hepatocellular Adenoma | | | | | | | |
| | | 42/50 (84%) | 34/50 (68%) | 37/50 (74%) | 17/49 (35%) | 31/50 (62%) | 27/50 (54%) |
| Overall Incidence | Total | 113/150 (75.33%) | Mean 75.33% | SD 8.08% | Total | 75/149 (50.34%) | Mean 50.23% SD 14.04% |
| | | | | | | | |
| *Liver: | | | | | | | |
| Hepatocellular Carcinoma, Hepatocellular Adenoma, or Hepatoblastoma | | | | | | | |
| | | 46/50 (92%) | 35/50 (70%) | 37/50 (74%) | 17/49 (35%) | 31/50 (62%) | 27/50 (54%) |
| Overall Incidence | Total | 118/150 (78.67%) | Mean 78.67% | SD 11.72% | Total | 75/149 (50.34%) | Mean 50.23% SD 14.04% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 253
 Report Date: 03/22/2010

| | | Male | | | | Female | | | | | |
|------------------------------------|-------|-------|---------|------------|----------|--------|-------|------|---------|-------|------|
| *Liver: | | | | | | | | | | | |
| Hepatocholangiocarcinoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/49 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/149 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| *Liver: | | | | | | | | | | | |
| Hepatocholangioma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/49 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/149 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| *Liver: | | | | | | | | | | | |
| Ito Cell Tumor Benign | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/49 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/149 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| *Liver: | | | | | | | | | | | |
| Ito Cell Tumor Malignant | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 2/50 | (4%) | 0/49 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 2/150 | (1.33%) | Mean 1.33% | SD 2.31% | Total | 0/149 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| *Liver: | | | | | | | | | | | |
| Plasma Cell Tumor Malignant | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/49 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/149 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| *Liver: | | | | | | | | | | | |
| Sarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/49 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/149 | (0%) | Mean 0% | SD 0% | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 254
 Report Date: 03/22/2010

| | | Male | | | Female | | | |
|---|-------|-----------------|-------------|-------------|-----------|-----------------|-------------|----------|
| *Lung: | | | | | | | | |
| Adenoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Lung: | | | | | | | | |
| Alveolar/Bronchiolar Adenoma | | | | | | | | |
| | | 15/50 (30%) | 4/50 (8%) | 6/50 (12%) | 1/50 (2%) | 2/50 (4%) | 3/50 (6%) | |
| Overall Incidence | Total | 25/150 (16.67%) | Mean 16.67% | SD 11.72% | Total | 6/150 (4%) | Mean 4% | SD 2% |
| | | | | | | | | |
| *Lung: | | | | | | | | |
| Alveolar/Bronchiolar Carcinoma | | | | | | | | |
| | | 3/50 (6%) | 12/50 (24%) | 4/50 (8%) | 2/50 (4%) | 6/50 (12%) | 6/50 (12%) | |
| Overall Incidence | Total | 19/150 (12.67%) | Mean 12.67% | SD 9.87% | Total | 14/150 (9.33%) | Mean 9.33% | SD 4.62% |
| | | | | | | | | |
| *Lung: | | | | | | | | |
| Alveolar/Bronchiolar Carcinoma or Alveolar/Bronchiolar Adenoma | | | | | | | | |
| | | 16/50 (32%) | 15/50 (30%) | 10/50 (20%) | 2/50 (4%) | 8/50 (16%) | 9/50 (18%) | |
| Overall Incidence | Total | 41/150 (27.33%) | Mean 27.33% | SD 6.43% | Total | 19/150 (12.67%) | Mean 12.67% | SD 7.57% |
| | | | | | | | | |
| *Lung: | | | | | | | | |
| Fibrosarcoma | | | | | | | | |
| | | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Lung: | | | | | | | | |
| Fibrous Histiocytoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 255
 Report Date: 03/22/2010

| | | Male | | | | Female | | | | | |
|------------------------------------|-------|-------|------|---------|-------|--------|-------|------|---------|-------|------|
| *Lung: | | | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| *Lung: | | | | | | | | | | | |
| Plasma Cell Tumor Malignant | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| *Lung: | | | | | | | | | | | |
| Sarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Lymph Node: | | | | | | | | | | | |
| Fibrous Histiocytoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Lymph Node: | | | | | | | | | | | |
| Hemangioma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Lymph Node: | | | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 256
 Report Date: 03/22/2010

| | | Male | | | | Female | | | | | |
|------------------------------------|-------|-------|------|---------|-------|--------|-------|------|---------|-------|------|
| #Lymph Node: | | | | | | | | | | | |
| Plasma Cell Tumor Malignant | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Lymph Node: | | | | | | | | | | | |
| Sarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Lymph Node, Bronchial: | | | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Lymph Node, Mandibular: | | | | | | | | | | | |
| Fibrous Histiocytoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Lymph Node, Mandibular: | | | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Lymph Node, Mandibular: | | | | | | | | | | | |
| Mast Cell Tumor Malignant | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 257
 Report Date: 03/22/2010

| | | Male | | | Female | | |
|------------------------------------|-------|------------|-----------|-----------|-----------|------------|------------------|
| #Lymph Node, Mandibular: | | | | | | | |
| Plasma Cell Tumor Malignant | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Lymph Node, Mesenteric: | | | | | | | |
| Fibrous Histiocytoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Lymph Node, Mesenteric: | | | | | | | |
| Hemangioma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Lymph Node, Mesenteric: | | | | | | | |
| Hemangiosarcoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Lymph Node, Mesenteric: | | | | | | | |
| Plasma Cell Tumor Malignant | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Mammary Gland: | | | | | | | |
| Adenoacanthoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 258
 Report Date: 03/22/2010

| | | Male | | | | Female | | | |
|---|-------|-------------------------|----------------------|--------------------|-------|-------------------------|----------------------|--------------------|--|
| #Mammary Gland: Adenoma | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | |
| #Mammary Gland: Carcinoma | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 3/150 (2%) | 2/50 (4%) Mean 2% | 1/50 (2%) SD 2% | |
| #Mammary Gland: Carcinoma or Adenoma | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 3/150 (2%) | 2/50 (4%) Mean 2% | 1/50 (2%) SD 2% | |
| #Mammary Gland: Fibroadenoma | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | |
| #Mammary Gland: Fibroma, Fibroadenoma or Adenoma | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | |
| #Mammary Gland: Fibroma, Fibroadenoma, Carcinoma, or Adenoma | | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 3/150 (2%) | 2/50 (4%) Mean 2% | 1/50 (2%) SD 2% | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 259
 Report Date: 03/22/2010

| | | Male | | | | Female | | | | | | | |
|---|-------|-------|------|---------|-------|--------|-------|---------|------------|----------|------|------|------|
| #Mesentery: | | | | | | | | | | | | | |
| Fibrosarcoma | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 3/50 | (6%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 3/150 | (2%) | Mean 2% | SD 3.46% | | | |
| | | | | | | | | | | | | | |
| #Mesentery: | | | | | | | | | | | | | |
| Hemangioma | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | | | |
| | | | | | | | | | | | | | |
| #Mesentery: | | | | | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | | | |
| | | | | | | | | | | | | | |
| #Mesentery: | | | | | | | | | | | | | |
| Sarcoma | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 1/50 | (2%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 1/150 | (0.67%) | Mean 0.67% | SD 1.15% | | | |
| | | | | | | | | | | | | | |
| *Nose: | | | | | | | | | | | | | |
| Mast Cell Tumor Malignant | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | | | |
| | | | | | | | | | | | | | |
| #Oral Cavity (Oral Mucosa, Tongue, Pharynx, Tooth, Gingiva): | | | | | | | | | | | | | |
| Papilloma Squamous or Papilloma | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | | | |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 260
 Report Date: 03/22/2010

| | Male | | | | Female | | | |
|---|-------|------------|-----------|-----------|--------|---------------|------------|-----------|
| #Oral Cavity (Oral Mucosa, Tongue, Pharynx, Tooth, Gingiva): | | | | | | | | |
| Squamous Cell Carcinoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Oral Cavity (Oral Mucosa, Tongue, Pharynx, Tooth, Gingiva): | | | | | | | | |
| Squamous Cell Carcinoma, Papilloma Squamous, or Papilloma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Ovary: | | | | | | | | |
| Choriocarcinoma | | | | | | | | |
| | | | | | | 0/50 (0%) | 0/47 (0%) | 1/50 (2%) |
| Overall Incidence | Total | | | | Total | 1/147 (0.68%) | Mean 0.67% | SD 1.15% |
| | | | | | | | | |
| *Ovary: | | | | | | | | |
| Cystadenoma | | | | | | | | |
| | | | | | | 1/50 (2%) | 2/47 (4%) | 1/50 (2%) |
| Overall Incidence | Total | | | | Total | 4/147 (2.72%) | Mean 2.75% | SD 1.3% |
| | | | | | | | | |
| *Ovary: | | | | | | | | |
| Fibrous Histiocytoma | | | | | | | | |
| | | | | | | 0/50 (0%) | 0/47 (0%) | 0/50 (0%) |
| Overall Incidence | Total | | | | Total | 0/147 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Ovary: | | | | | | | | |
| Granulosa Cell Tumor Benign | | | | | | | | |
| | | | | | | 0/50 (0%) | 0/47 (0%) | 1/50 (2%) |
| Overall Incidence | Total | | | | Total | 1/147 (0.68%) | Mean 0.67% | SD 1.15% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 261
 Report Date: 03/22/2010

| | Male | | Female | |
|---|-------|---------------|------------|-----------|
| *Ovary: | | | | |
| Granulosa Cell Tumor Malignant | | | | |
| | | 0/50 (0%) | 0/47 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/147 (0%) | Mean 0% | SD 0% |
| | | | | |
| *Ovary: | | | | |
| Granulosa Cell Tumor: Benign, Malignant, NOS | | | | |
| | | 0/50 (0%) | 0/47 (0%) | 1/50 (2%) |
| Overall Incidence | Total | 1/147 (0.68%) | Mean 0.67% | SD 1.15% |
| | | | | |
| *Ovary: | | | | |
| Hemangioma | | | | |
| | | 0/50 (0%) | 0/47 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/147 (0%) | Mean 0% | SD 0% |
| | | | | |
| *Ovary: | | | | |
| Hemangiosarcoma | | | | |
| | | 0/50 (0%) | 0/47 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/147 (0%) | Mean 0% | SD 0% |
| | | | | |
| *Ovary: | | | | |
| Leiomyosarcoma | | | | |
| | | 0/50 (0%) | 0/47 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/147 (0%) | Mean 0% | SD 0% |
| | | | | |
| *Ovary: | | | | |
| Luteoma | | | | |
| | | 0/50 (0%) | 0/47 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/147 (0%) | Mean 0% | SD 0% |
| | | | | |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 262
 Report Date: 03/22/2010

| | | Male | | | | Female | | | |
|------------------------------------|-------|---------------|--|------------|--|-----------|--|-----------|--|
| *Ovary: | | | | | | | | | |
| Plasma Cell Tumor Malignant | | | | | | | | | |
| | | 0/50 (0%) | | 0/47 (0%) | | 0/50 (0%) | | | |
| Overall Incidence | Total | 0/147 (0%) | | Mean 0% | | SD 0% | | | |
| | | | | | | | | | |
| *Ovary: | | | | | | | | | |
| Teratoma Benign | | | | | | | | | |
| | | 0/50 (0%) | | 0/47 (0%) | | 0/50 (0%) | | | |
| Overall Incidence | Total | 0/147 (0%) | | Mean 0% | | SD 0% | | | |
| | | | | | | | | | |
| *Ovary: | | | | | | | | | |
| Teratoma Malignant | | | | | | | | | |
| | | 0/50 (0%) | | 0/47 (0%) | | 0/50 (0%) | | | |
| Overall Incidence | Total | 0/147 (0%) | | Mean 0% | | SD 0% | | | |
| | | | | | | | | | |
| *Ovary: | | | | | | | | | |
| Tubulostromal Adenoma | | | | | | | | | |
| | | 0/50 (0%) | | 0/47 (0%) | | 1/50 (2%) | | | |
| Overall Incidence | Total | 1/147 (0.68%) | | Mean 0.67% | | SD 1.15% | | | |
| | | | | | | | | | |
| *Pancreas: | | | | | | | | | |
| Adenoma | | | | | | | | | |
| | | 0/49 (0%) | | 0/50 (0%) | | 0/50 (0%) | | 0/48 (0%) | |
| Overall Incidence | Total | 0/149 (0%) | | Mean 0% | | SD 0% | | 0/49 (0%) | |
| | | | | | | | | SD 0% | |
| | | | | | | | | | |
| *Pancreas: | | | | | | | | | |
| Carcinoma | | | | | | | | | |
| | | 0/49 (0%) | | 0/50 (0%) | | 0/50 (0%) | | 0/48 (0%) | |
| Overall Incidence | Total | 0/149 (0%) | | Mean 0% | | SD 0% | | 1/49 (2%) | |
| | | | | | | | | SD 1.18% | |
| | | | | | | | | | |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 263
 Report Date: 03/22/2010

| | | Male | | | | Female | | | |
|---|-------|---------------|------------|-----------|-------|---------------|------------|-----------|--|
| *Pancreas: | | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/48 (0%) | 1/49 (2%) | 0/49 (0%) | |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 1/146 (0.68%) | Mean 0.68% | SD 1.18% | |
| ----- | | | | | | | | | |
| *Pituitary Gland: Pars Distalis or Unspecified Site: | | | | | | | | | |
| Adenoma | | | | | | | | | |
| | | 0/49 (0%) | 1/50 (2%) | 0/49 (0%) | | 3/48 (6%) | 4/48 (8%) | 2/48 (4%) | |
| Overall Incidence | Total | 1/148 (0.68%) | Mean 0.67% | SD 1.15% | Total | 9/144 (6.25%) | Mean 6.25% | SD 2.08% | |
| ----- | | | | | | | | | |
| *Pituitary Gland: Pars Distalis or Unspecified Site: | | | | | | | | | |
| Carcinoma | | | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/49 (0%) | | 0/48 (0%) | 0/48 (0%) | 0/48 (0%) | |
| Overall Incidence | Total | 0/148 (0%) | Mean 0% | SD 0% | Total | 0/144 (0%) | Mean 0% | SD 0% | |
| ----- | | | | | | | | | |
| *Pituitary Gland: Pars Distalis or Unspecified Site: | | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | | |
| | | 0/49 (0%) | 1/50 (2%) | 0/49 (0%) | | 3/48 (6%) | 4/48 (8%) | 2/48 (4%) | |
| Overall Incidence | Total | 1/148 (0.68%) | Mean 0.67% | SD 1.15% | Total | 9/144 (6.25%) | Mean 6.25% | SD 2.08% | |
| ----- | | | | | | | | | |
| *Pituitary Gland: Pars Intermedia: | | | | | | | | | |
| Adenoma | | | | | | | | | |
| | | 0/49 (0%) | 1/50 (2%) | 0/49 (0%) | | 0/48 (0%) | 0/48 (0%) | 0/48 (0%) | |
| Overall Incidence | Total | 1/148 (0.68%) | Mean 0.67% | SD 1.15% | Total | 0/144 (0%) | Mean 0% | SD 0% | |
| ----- | | | | | | | | | |
| *Pituitary Gland: Pars Intermedia: | | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | | |
| | | 0/49 (0%) | 1/50 (2%) | 0/49 (0%) | | 0/48 (0%) | 0/48 (0%) | 0/48 (0%) | |
| Overall Incidence | Total | 1/148 (0.68%) | Mean 0.67% | SD 1.15% | Total | 0/144 (0%) | Mean 0% | SD 0% | |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 264
 Report Date: 03/22/2010

| | | Male | | | | Female | | | |
|----------------------------------|-------|-------|---------|------------|------|----------|------|---------|------------|
| *Prostate: | | | | | | | | | |
| Adenoma | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | | |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | | SD 0% | | | |
| | | | | | | | | | |
| *Prostate: | | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | | |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | | SD 0% | | | |
| | | | | | | | | | |
| *Prostate: | | | | | | | | | |
| Hemangioma | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | | |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | | SD 0% | | | |
| | | | | | | | | | |
| *Salivary Glands: | | | | | | | | | |
| Carcinoma | | | | | | | | | |
| | | 0/49 | (0%) | 0/50 | (0%) | 1/50 | (2%) | 0/50 | (0%) |
| Overall Incidence | Total | 1/149 | (0.67%) | Mean 0.67% | | SD 1.15% | | Total | 0/150 (0%) |
| | | | | | | | | Mean 0% | SD 0% |
| | | | | | | | | | |
| *Salivary Glands: | | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | | |
| | | 0/49 | (0%) | 0/50 | (0%) | 1/50 | (2%) | 0/50 | (0%) |
| Overall Incidence | Total | 1/149 | (0.67%) | Mean 0.67% | | SD 1.15% | | Total | 0/150 (0%) |
| | | | | | | | | Mean 0% | SD 0% |
| | | | | | | | | | |
| *Salivary Glands: | | | | | | | | | |
| Mast Cell Tumor Malignant | | | | | | | | | |
| | | 0/49 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/149 | (0%) | Mean 0% | | SD 0% | | Total | 0/150 (0%) |
| | | | | | | | | Mean 0% | SD 0% |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 265
 Report Date: 03/22/2010

| | | Male | | | | Female | | | |
|--|-------|-------|------|---------|------|--------|-------|---------|-------|
| #Seminal Vesicle: | | | | | | | | | |
| Adenoma | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | | |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | | SD 0% | | | |
| | | | | | | | | | |
| #Skeletal Muscle: | | | | | | | | | |
| Fibrosarcoma | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | | SD 0% | Total | 0/150 | (0%) |
| | | | | | | | | Mean 0% | SD 0% |
| | | | | | | | | | |
| #Skeletal Muscle: | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | | SD 0% | Total | 0/150 | (0%) |
| | | | | | | | | Mean 0% | SD 0% |
| | | | | | | | | | |
| #Skeletal Muscle: | | | | | | | | | |
| Myxosarcoma | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | | SD 0% | Total | 0/150 | (0%) |
| | | | | | | | | Mean 0% | SD 0% |
| | | | | | | | | | |
| #Skeletal Muscle: | | | | | | | | | |
| Rhabdomyosarcoma | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | | SD 0% | Total | 0/150 | (0%) |
| | | | | | | | | Mean 0% | SD 0% |
| | | | | | | | | | |
| #Skeletal Muscle: | | | | | | | | | |
| Rhabdomyosarcoma or Rhabdomyoma | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | | SD 0% | Total | 0/150 | (0%) |
| | | | | | | | | Mean 0% | SD 0% |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 266
 Report Date: 03/22/2010

| | | Male | | | | Female | | | | | |
|--|-------|-------|---------|------------|----------|--------|-------|------|---------|-------|------|
| #Skeletal Muscle: | | | | | | | | | | | |
| Sarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | |
| Basal Cell Adenoma or Basosquamous Tumor Benign | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | |
| Basal Cell Adenoma, Basosquamous Tumor Benign, or Trichoepithelioma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | |
| Basal Cell Carcinoma | | | | | | | | | | | |
| | | 1/50 | (2%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 1/150 | (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | |
| Basal Cell Carcinoma or Basosquamous Tumor (malignant or NOS) | | | | | | | | | | | |
| | | 1/50 | (2%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 1/150 | (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | |
| Basal Cell Carcinoma, Basal Cell Adenoma, Basosquamous Tumor (benign, malignant or NOS), or Trichoepithelioma | | | | | | | | | | | |
| | | 1/50 | (2%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 1/150 | (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 267
 Report Date: 03/22/2010

| | Male | | | | | Female | | | | |
|---|-------|---------------|------------|-----------|-------|---------------|------------|-----------|-----------|--|
| #Skin: | | | | | | | | | | |
| Basal or Sq. Cell Carcinoma, Carcinoma, Basosq. Tumor (M or B), Basal Cell Adenoma, Adenoma, Papilloma, Sq Papilloma, Keratoacanthoma, Trichoepithelioma | | | | | | | | | | |
| | | 1/50 (2%) | 1/50 (2%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 2/150 (1.33%) | Mean 1.33% | SD 1.15% | Total | 0/150 (0%) | Mean 0% | SD 0% | | |
| | | | | | | | | | | |
| #Skin: | | | | | | | | | | |
| Basosquamous Tumor Benign | | | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% | | |
| | | | | | | | | | | |
| #Skin: | | | | | | | | | | |
| Fibroma | | | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 (0%) | Mean 0% | SD 0% | | |
| | | | | | | | | | | |
| #Skin: | | | | | | | | | | |
| Fibroma, Fibrosarcoma, Sarcoma, Myxoma, Myxosarcoma, or Fibrous Histiocytoma | | | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | | 2/50 (4%) | 2/50 (4%) | 4/50 (8%) | | |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 8/150 (5.33%) | Mean 5.33% | SD 2.31% | | |
| | | | | | | | | | | |
| #Skin: | | | | | | | | | | |
| Fibrosarcoma | | | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 1/50 (2%) | 0/50 (0%) | 4/50 (8%) | | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 5/150 (3.33%) | Mean 3.33% | SD 4.16% | | |
| | | | | | | | | | | |
| #Skin: | | | | | | | | | | |
| Fibrosarcoma, Sarcoma, Myxosarcoma, or Fibrous Histiocytoma | | | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 2/50 (4%) | 2/50 (4%) | 4/50 (8%) | | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 8/150 (5.33%) | Mean 5.33% | SD 2.31% | | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 268
 Report Date: 03/22/2010

| | | Male | | | | Female | | | | | | | |
|-----------------------------------|-------|-------|------|---------|-------|--------|-------|---------|------------|----------|------|------|------|
| #Skin: | | | | | | | | | | | | | |
| Fibrous Histiocytoma | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 1/50 | (2%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 1/150 | (0.67%) | Mean 0.67% | SD 1.15% | | | |
| | | | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | | | |
| Granular Cell Tumor Benign | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | | | |
| | | | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | | | |
| Hemangioma | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 2/50 | (4%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 2/150 | (1.33%) | Mean 1.33% | SD 2.31% | | | |
| | | | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 1/50 | (2%) | | |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 1/150 | (0.67%) | Mean 0.67% | SD 1.15% | | | |
| | | | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | | | |
| Liposarcoma | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | | | |
| | | | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | | | |
| Mast Cell Tumor Benign | | | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | | | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 269
 Report Date: 03/22/2010

| | | Male | | | | Female | | | | | |
|--|-------|-------|------|---------|-------|--------|-------|---------|------------|----------|------|
| #Skin: | | | | | | | | | | | |
| Myxosarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | |
| Neurofibrosarcoma or Schwannoma (malignant or NOS) | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | |
| Neurofibrosarcoma, Neurofibroma, or Schwannoma (benign, malignant or NOS) | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | |
| Osteosarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | |
| Sarcoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 1/50 | (2%) | 1/50 | (2%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 2/150 | (1.33%) | Mean 1.33% | SD 1.15% | |
| | | | | | | | | | | | |
| #Skin: | | | | | | | | | | | |
| Schwannoma Malignant | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 270
 Report Date: 03/22/2010

| | | Male | | | | Female | | | |
|---|-------|---------------|------------|-----------|-------|------------|-----------|-----------|--|
| #Skin: | | | | | | | | | |
| Squamous Cell Carcinoma | | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Skin: | | | | | | | | | |
| Squamous Cell Carcinoma, Basal Cell Carcinoma, Basosquamous Tumor (malignant or NOS), or Carcinoma | | | | | | | | | |
| | | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Skin: | | | | | | | | | |
| Squamous Cell Papilloma | | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Skin: | | | | | | | | | |
| Squamous Cell Papilloma or Papilloma | | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Skin: | | | | | | | | | |
| Squamous Cell Papilloma, Papilloma, Squamous Cell Carcinoma or Keratoacanthoma | | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Skin: | | | | | | | | | |
| Squamous Cell Papilloma, Papilloma, or Keratoacanthoma | | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 (0%) | Mean 0% | SD 0% | |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 271
 Report Date: 03/22/2010

| | | Male | | | | Female | | | | | |
|--------------------------------|-------|-------|------|---------|-------|--------|-------|---------|------------|----------|------|
| #Skin: | | | | | | | | | | | |
| Trichoepithelioma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: Sebaceous Gland: | | | | | | | | | | | |
| Adenoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 1/50 | (2%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 1/150 | (0.67%) | Mean 0.67% | SD 1.15% | |
| | | | | | | | | | | | |
| #Skin: Sebaceous Gland: | | | | | | | | | | | |
| Carcinoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| #Skin: Sebaceous Gland: | | | | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 1/50 | (2%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 1/150 | (0.67%) | Mean 0.67% | SD 1.15% | |
| | | | | | | | | | | | |
| #Spinal Cord: | | | | | | | | | | | |
| Meningioma Malignant | | | | | | | | | | | |
| | | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/50 | (0%) |
| Overall Incidence | Total | 0/150 | (0%) | Mean 0% | SD 0% | Total | 0/150 | (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | | | |
| *Spleen: | | | | | | | | | | | |
| Fibrous Histiocytoma | | | | | | | | | | | |
| | | 0/49 | (0%) | 0/50 | (0%) | 0/50 | (0%) | 0/48 | (0%) | 0/48 | (0%) |
| Overall Incidence | Total | 0/149 | (0%) | Mean 0% | SD 0% | Total | 0/145 | (0%) | Mean 0% | SD 0% | |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 272
 Report Date: 03/22/2010

| | | Male | | | Female | | | |
|------------------------------------|-------|---------------|------------|-----------|-----------|---------------|------------|----------|
| *Spleen: | | | | | | | | |
| Hemangioma | | | | | | | | |
| | | 0/49 (0%) | 1/50 (2%) | 0/50 (0%) | 0/48 (0%) | 0/49 (0%) | 0/48 (0%) | |
| Overall Incidence | Total | 1/149 (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/145 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Spleen: | | | | | | | | |
| Hemangiosarcoma | | | | | | | | |
| | | 2/49 (4%) | 1/50 (2%) | 1/50 (2%) | 1/48 (2%) | 1/49 (2%) | 0/48 (0%) | |
| Overall Incidence | Total | 4/149 (2.68%) | Mean 2.69% | SD 1.2% | Total | 2/145 (1.38%) | Mean 1.37% | SD 1.19% |
| | | | | | | | | |
| *Spleen: | | | | | | | | |
| Mast Cell Tumor Malignant | | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | 0/48 (0%) | 0/49 (0%) | 0/48 (0%) | |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 0/145 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Spleen: | | | | | | | | |
| Plasma Cell Tumor Malignant | | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | 0/48 (0%) | 0/49 (0%) | 0/48 (0%) | |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 0/145 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Stomach, Forestomach: | | | | | | | | |
| Mast Cell Tumor Malignant | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Stomach, Forestomach: | | | | | | | | |
| Squamous Cell Carcinoma | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% | Total | 0/150 (0%) | Mean 0% | SD 0% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 273
 Report Date: 03/22/2010

| | | Male | | | Female | | | |
|--|-------|---------------|------------|-----------|-----------|---------------|------------|----------|
| #Stomach, Forestomach: | | | | | | | | |
| Squamous Cell Carcinoma or Papilloma Squamous | | | | | | | | |
| | | 0/50 (0%) | 2/50 (4%) | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | 1/50 (2%) | |
| Overall Incidence | Total | 2/150 (1.33%) | Mean 1.33% | SD 2.31% | Total | 2/150 (1.33%) | Mean 1.33% | SD 1.15% |
| | | | | | | | | |
| #Stomach, Forestomach: | | | | | | | | |
| Squamous Cell Papilloma | | | | | | | | |
| | | 0/50 (0%) | 2/50 (4%) | 0/50 (0%) | 1/50 (2%) | 0/50 (0%) | 1/50 (2%) | |
| Overall Incidence | Total | 2/150 (1.33%) | Mean 1.33% | SD 2.31% | Total | 2/150 (1.33%) | Mean 1.33% | SD 1.15% |
| | | | | | | | | |
| #Stomach, Glandular: | | | | | | | | |
| Adenoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| #Stomach, Glandular: | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Testes: | | | | | | | | |
| Adenoma | | | | | | | | |
| | | 0/50 (0%) | 1/50 (2%) | 1/50 (2%) | | | | |
| Overall Incidence | Total | 2/150 (1.33%) | Mean 1.33% | SD 1.15% | | | | |
| | | | | | | | | |
| *Thymus: | | | | | | | | |
| Thymoma Benign | | | | | | | | |
| | | 0/49 (0%) | 0/46 (0%) | 0/44 (0%) | 0/46 (0%) | 0/49 (0%) | 0/46 (0%) | |
| Overall Incidence | Total | 0/139 (0%) | Mean 0% | SD 0% | Total | 0/141 (0%) | Mean 0% | SD 0% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 274
 Report Date: 03/22/2010

| | | Male | | | Female | | | |
|---|-------|---------------|------------|-----------|-----------|---------------|------------|----------|
| *Thymus: | | | | | | | | |
| Thymoma Malignant | | | | | | | | |
| | | 0/49 (0%) | 0/46 (0%) | 0/44 (0%) | 0/46 (0%) | 0/49 (0%) | 0/46 (0%) | |
| Overall Incidence | Total | 0/139 (0%) | Mean 0% | SD 0% | Total | 0/141 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Thymus: | | | | | | | | |
| Thymoma: Benign, Malignant, NOS | | | | | | | | |
| | | 0/49 (0%) | 0/46 (0%) | 0/44 (0%) | 0/46 (0%) | 0/49 (0%) | 0/46 (0%) | |
| Overall Incidence | Total | 0/139 (0%) | Mean 0% | SD 0% | Total | 0/141 (0%) | Mean 0% | SD 0% |
| | | | | | | | | |
| *Thyroid Gland: C-Cell: | | | | | | | | |
| Carcinoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 1/50 (2%) | 0/49 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 1/149 (0.67%) | Mean 0.67% | SD 1.15% |
| | | | | | | | | |
| *Thyroid Gland: C-Cell: | | | | | | | | |
| Carcinoma or Adenoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 1/50 (2%) | 0/49 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 1/149 (0.67%) | Mean 0.67% | SD 1.15% |
| | | | | | | | | |
| *Thyroid Gland: Follicular Cell: | | | | | | | | |
| Adenoma | | | | | | | | |
| | | 1/50 (2%) | 3/50 (6%) | 3/50 (6%) | 1/50 (2%) | 3/50 (6%) | 1/49 (2%) | |
| Overall Incidence | Total | 7/150 (4.67%) | Mean 4.67% | SD 2.31% | Total | 5/149 (3.36%) | Mean 3.35% | SD 2.3% |
| | | | | | | | | |
| *Thyroid Gland: Follicular Cell: | | | | | | | | |
| Carcinoma | | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/49 (0%) | |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/149 (0%) | Mean 0% | SD 0% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 275
 Report Date: 03/22/2010

| | | Male | | | Female | | | |
|--|-------|----------------------------|-------------------------|-----------------------|--------|----------------------------|-------------------------|----------------------|
| *Thyroid Gland: Follicular Cell: Carcinoma or Adenoma | | | | | | | | |
| Overall Incidence | Total | 1/50 (2%) 7/150 (4.67%) | 3/50 (6%) Mean 4.67% | 3/50 (6%) SD 2.31% | Total | 1/50 (2%) 5/149 (3.36%) | 3/50 (6%) Mean 3.35% | 1/49 (2%) SD 2.3% |
| #Tissue NOS: Carcinoma | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% |
| #Tissue NOS: Hemangiosarcoma | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% |
| #Tissue NOS: Plasma Cell Tumor Malignant | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% |
| #Tissue NOS: Sarcoma | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% |
| #Tongue: Fibroma | | | | | | | | |
| Overall Incidence | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% | Total | 0/50 (0%) 0/150 (0%) | 0/50 (0%) Mean 0% | 0/50 (0%) SD 0% |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 276
 Report Date: 03/22/2010

| | | Male | | | Female | | |
|--|-------|------------|-----------|-----------|-----------|------------|------------------|
| #Tongue: | | | | | | | |
| Squamous Cell Carcinoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Tongue: | | | | | | | |
| Squamous Cell Carcinoma or Papilloma Squamous | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Tongue: | | | | | | | |
| Squamous Cell Papilloma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| #Tooth: | | | | | | | |
| Odontoma | | | | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Urinary Bladder: | | | | | | | |
| Hemangioma | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |
| | | | | | | | |
| *Urinary Bladder: | | | | | | | |
| Hemangiosarcoma | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% SD 0% |

*: Denominator is number of animals with tissues examined microscopically
 #: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
Route: ORAL
Vehicle: WATER

Page: 277
 Report Date: 03/22/2010

| | | Male | | | | Female | | | |
|------------------------------------|-------|------------|-----------|-----------|-------|------------|-----------|-----------|--|
| *Urinary Bladder: | | | | | | | | | |
| Plasma Cell Tumor Malignant | | | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| *Urinary Bladder: | | | | | | | | | |
| Sarcoma | | | | | | | | | |
| | | 0/49 (0%) | 0/50 (0%) | 0/50 (0%) | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | Total | 0/149 (0%) | Mean 0% | SD 0% | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Uterus: | | | | | | | | | |
| Adenoma | | | | | | | | | |
| | | | | | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | | | | | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Uterus: | | | | | | | | | |
| Carcinoma | | | | | | | | | |
| | | | | | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | | | | | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Uterus: | | | | | | | | | |
| Hemangioma | | | | | | | | | |
| | | | | | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | | | | | Total | 0/150 (0%) | Mean 0% | SD 0% | |
| | | | | | | | | | |
| #Uterus: | | | | | | | | | |
| Hemangiosarcoma | | | | | | | | | |
| | | | | | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) | |
| Overall Incidence | | | | | Total | 0/150 (0%) | Mean 0% | SD 0% | |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 278
 Report Date: 03/22/2010

| | Male | Female |
|---|-----------------|-------------------------------------|
| #Uterus: | | |
| Leiomyoma | | |
| Overall Incidence | Total 0/50 (0%) | 0/50 (0%) Mean 0% SD 0% |
| | | |
| #Uterus: | | |
| Leiomyosarcoma | | |
| Overall Incidence | Total 0/50 (0%) | 0/50 (0%) Mean 0% SD 0% |
| | | |
| #Uterus: | | |
| Polyp Stromal | | |
| Overall Incidence | Total 1/50 (2%) | 2/50 (4%) Mean 2.67% SD 1.15% |
| | | |
| #Uterus: | | |
| Sarcoma Stromal | | |
| Overall Incidence | Total 0/50 (0%) | 0/50 (0%) Mean 0% SD 0% |
| | | |
| #Uterus: | | |
| Sarcoma Stromal or Polyp Stromal | | |
| Overall Incidence | Total 1/50 (2%) | 2/50 (4%) Mean 2.67% SD 1.15% |
| | | |
| #Uterus: | | |
| Schwannoma Malignant | | |
| Overall Incidence | Total 0/50 (0%) | 0/50 (0%) Mean 0% SD 0% |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied

Version: Mar2010
 Contract/Lab: All Laboratories
 Species: MICE
 Strain: B6C3F1
 Length of Study: CHRONIC

Toxicology Data Management System
 Tumor Incidence for Selected Control Animal Groups
 Route: ORAL
 Vehicle: WATER

Page: 279
 Report Date: 03/22/2010

| | Male | | Female | |
|--------------------------------|-------|---------------|------------|-----------|
| #Vagina: | | | | |
| Liposarcoma | | | | |
| | | 0/50 (0%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 0/150 (0%) | Mean 0% | SD 0% |
| | | | | |
| #Vagina: | | | | |
| Squamous Cell Papilloma | | | | |
| | | 1/50 (2%) | 0/50 (0%) | 0/50 (0%) |
| Overall Incidence | Total | 1/150 (0.67%) | Mean 0.67% | SD 1.15% |
| | | | | |

*: Denominator is number of animals with tissues examined microscopically

#: Denominator is number of animals necropsied