2,4-Hexadienal

CHEMICAL IDENTIFICATION:

Structure: CH3-CH=CH-CHO

Molecular Formula: C₆H₈O

Molecular Weight: 96.13

CAS Registry Number: 142-83-6

Other CAS Nos.: 53398-76-8 (trans, cis-2,4-Hx)

80466-34-8 (mixed isomers)

Chemical Abstr. Name: 2,4-Hexadienal,(E,E)-(9CI)

Synonyms/Trade Names: Sorbic aldehyde

Sorbaldehyde

2-propyleneacrolein

1,3-pentadiene-1-carboxaldehyde

trans,trans-2,4-hexadienal all-trans-2,4-hexadienal

hexa-2,4-dienal 2,4-hexadien-1-al

2,4-Hx

CHEMICAL AND PHYSICAL PROPERTIES

2,4-Hexadienal (2,4-Hx) is a colorless (yellowish) liquid with pungent sweet-green citrusy odor (Ford *et al.*, 1988). It is insoluble in water, soluble in alcohol (Bedoukian, 1993), and reacts with strong oxidizing and reducing agents. It has a boiling point of 64°C (147°F, Lancaster Synthesis, Inc., 1991), a specific gravity of 0.871, a vapor density of >1, and a vapor

pressure of 1.6 mm Hg at 20° C. Its refractive index is 1.535 and flash point is 130°F (67°C, Aldrich, 1991; 1992). 2,4-Hx is available from Aldrich Chemical Co. in 95% purity. No information on impurities is available.

NOMINATION RATIONALE

2,4-Hx is a natural constituent of meat, vegetable, and fish oils. It is one of the lipid peroxidation products of polyunsaturated oils which undergoes auto-oxidation especially during storage (Snyder et al., 1985) and have been implicated in the development of off or tainted flavor. 2,4-Hx is also used as a food additive or flavoring agent.

Lipid hydroperoxides have been shown to give rise to low intracellular levels of α,β -unsaturated aldehydes, including 2,4-Hx and 2,4-decadienal (2,4-De). Some of the α,β -unsaturated aldehydes have been shown to be reactive with DNA (Frankel et al., 1987). Ingested lipid oxidation products and oxidized fats have been reported to cause increased excretion of mutagens, cellular injury to liver and kidneys, increased cell proliferation in the gastro-intestinal tract, and other non-specific tissue injury and irritation effects resulting from induced oxidative stress.

There is an overall lack of data generated from testing of dienals for either carcinogenicity or mutagenicity and a lack of evidence specifically relating exposures to cancer in humans. The role of consumed oxidized oils in gastro-intestinal carcinogenesis including the effects of oral intake of different doses of various biologically active compounds present in heated oils, effects of oxidative stress induced by chronic consumption of repeatedly heated oils, as well as interactions with other modulating dietary factors, including both macro- and micro-nutrients have not been investigated (Hageman et al., 1991). Testing of 2,4-Hx and 2,4-De for toxicity and carcinogenicity would help mitigate the data gap.

The CSWG on 9/18/92 recommended with high priority evaluation of the

metabolism, distribution, mutagenesis, and carcinogenesis of 2,4-Hx with 2,4-De as a matched pair. These commonly found auto-oxidation products of unsaturated fats appear to be inactive in most standard mutagenicity prescreens, both show signs of mutagenic activity in tests using TA104, or by increasing the bacterial density, or by adding glutathione to the mix (Eder et al., 1992).

PRODUCTION AND PRODUCERS

둉 .

2,4-Hx is prepared by condensation of acetaldehyde (Keller et al., 1983). 2,4-Hx is available for sale from Aldrich Chemical Co., American Tokyo Kasei, Chem Service, Inc., Crescent Chemical Co., Pfaltz & Bauer, Inc., and Lancaster Synthesis, Inc. Current production levels are not available but the FDA's Priority-based Assessment of Food Additives (PAFA) database contains the following data (FDA, 1992):

Year # of Reporting Companies Market Disappearance Rate.lb/yr

1970	5	43
1975	-	6
1976	5	7
1982	7	26
1987	6	3.3

USE PATTERN

- 2,4-Hx is used as a flavoring agent in the manufacture of aroma chemical, 3,5,7-nonatrien-2-one. It is a chemical intermediate used in various organic synthetic reactions. 2,4-Hx has been used as the starting material for the manufacture of sorbic acid (Keller et al., 1983) widely used as a food preservative.
- 2,4-Hx is also used as chemical intermediate in the manufacture of polymethine dyes (Sturmer and Diehl, 1982), pharmaceutical intermediate

for manufacture of mitomycins and antihypercholesteremics, corrosion inhibitor for steel in oil field operations (Growcock et al., 1989), monomer reacted with a silane comonomer in polyalkenyloxysilane polymer manufacture, fumigant against larvae of the Caribbean fruit fly, reducing agent in agricultural research studies for photoreduction of nitrophenyl ether herbicides to nitro radical anions

Environmental Occurrence

2,4-Hx occurs naturally as an autoxidation product of unsaturated fatty acids, e.g. linoleic acid. It has been identified in numerous oxidized glyceridic oils, including canola (low erucic acid rapeseed) oil, soybean oil, cottonseed oil, sunflower oil, sesame oil and palm oil. In addition it has been detected in the essential oils of lovage, thyme leaf and dill and in solid alfalfa extract. It has been associated with rancidity and off-flavor in fish, including farm raised catfish, Gulf of Mexico menhaden and Upper Wisconsin River walleye and northern pike (Heil and Lindsay, 1988). occurs naturally in a variety of plant products, including cotton, tomatoes with poor flavor, mango, kiwi and Chinese quince, and is a component of tobacco leaf and tobacco smoke volatiles (Takeoka et al., 1986; Weeks et al., 1989; Wright and Harris, 1985; Zeringue and McCormick, 1989). It has been detected as a volatile of piled (rather than picked) Toyama Kurocha tea processed in Japan. 2,4-Hx was not found in fresh steamed or fermented tea leaves, but was reported at a concentration of 0.4 mg/100 mg in the solar dried product and 0.2 mg/100 mg in the product stored for 1 year (Kawakami and Shibamoto, 1991).

2, 4-Hx was identified as one of several ring fragmentation products present in polluted urban air (Dumdei et al., 1988). 2,4-Hx was also identified as a low-level carbonyl impurity in commercial ethanol, even in distilled premium grades (Sherman and Kavasmaneck, 1980).

The presence of 2,4-Hx in mussels, oysters at 35 mg/gm (35 ppb) and clams at 7.5 mg/gm (7.5 ppb) from Lake Pontchartrain has been attributed

to water pollution by VOCs (Ferrario et al., 1985). 2,4-Hx has also been cited in a recent Russian review of aldehydic environmental pollutants.

Human Exposure

2,4-Hx has been detected in tobacco and tobacco smoke (Florin et al., 1980; Pettersson et al., 1980) and is present in seafood (Ferrario et al., 1985), oxidized edible fats and oils, heated oils for food frying, and rancid or off-flavored fish oils (Przybylski and Hougen, 1989; Selke and Rohwedder, 1983; Suzuki and Bailey, 1985; White and Hammond, 1983). Other consumed products in which it has been detected include meat fat, cow's milk fat, potato chips, bread crust, dried and stored piled tea, herbs and spices, and tropical fruits.

According to the Flavor & Extract Manufacturers Association total yearly consumption of 2,4-Hx is 2.48 kg of which 0.9 kg (36%) is used as a flavor ingredient and 1.58 kg is consumed as a component of food (FEMA No. 3429). According to Ford, et al. (1988) the reported maximum concentration of this chemical as an ingredient in consumer products is 0.1%.

Regulatory Status

- 2,4-Hx is listed in EPA's TSCA inventory. No standards or guidelines have been set for occupational exposures to or environmental levels of 2,4-Hx. The American Conference of Governmental Industrial Hygienists (ACGIH) has not adopted a TLV/TWA for this compound.
- 2,4-Hx was given GRAS (generally recognized as safe) status after a review of flavoring ingredients and food additives by the Flavoring Extract Manufacturers' Association (FEMA) and was listed in 1981 by the Council of Europe as a flavoring substance that may be added to food (Ford *et al.*, 1988).

Toxicity

 α,β -unsaturated aldehydes are direct-acting alkylating agents capable of covalent binding without prior metabolism to cellular nucleophilic groups (Eder et al., 1993). The potential for being toxic and/or capable of modifying cellular processes is inherently present. Very little is known about the in vivo effects of 2,4-Hx, their target tissues, and mechanisms of toxicity.

Human Data: No information was found in the literature associating 2,4-Hx with a cancer risk in humans. Health hazard advisory information in the Aldrich MSDS includes severe irritant and toxic effects following inhalation or dermal absorption, with tissue destruction of mucous membranes of the upper respiratory tract, eyes and skin. Ford et al. (1988) cited the results of dermal 48-hr closed patch tests carried out at a concentration of 1% 2,4-Hx in petroleum on the backs of 59 volunteers in which no irritation but 1 case of sensitization was reported.

High rates of colon cancer in the northeastern U.S. have been associated with lifestyle, especially diet. Furthermore, it has been postulated that the high colon cancer incidence may be linked to a high fat diet or one low in fruits and vegetables and low in vitamin A (Urbany, 1992). Free radicals, singlet oxygen and other reactive species formed in the peroxidation of lipids are considered biologically harmful and are implicated in cellular damage and cancer (Frankel et al., 1987). The difficulties of assessing the cancer risk from multiple low level exposures to a wide variety of food toxicants is daunting. Marnett et al. (1985) have postulated that, "since carbonyl compounds are widely distributed in foods, are generated during cellular metabolism, and are present in body fluids, they make a significant contribution to the risk of human cancer."

Animal Data: The feeding of lipid oxidation products and oxidized fats has been reported to cause adverse biological effects in laboratory animals, including growth retardation, teratogenicity, tissue damage and increased

weights of the liver and kidneys (Alexander et al., 1987; Izaki et al., 1984; Kanazawa et al., 1985; 1986), cellular damage to the testes and epididymides, increased peroxidation of membrane and tissue lipids and induction of cytochrome P450 activities in the liver and colon (Crawford and Wheeler, 1983). No information was found in the literature associating 2,4-Hx with a cancer risk in animals.

The following information is recorded in RTECS database (NLM, 1992):

Oral rat LD₅₀:

Dermal rabbit LD₅₀:

Dermal guinea pig LD₁₀:

Dermal guinea pig LD₅₀

Z,500 mg/kg

5,000 mg/kg

Inhalation rat LC₁₀:

2,000 ppm/4H

Dermal rabbit: Severe irritant
Ocular rabbit: Severe irritant

Dermal guinea pig: Severe irritant and sensitizer

The irritating effect of 2,4-Hx may cause cellular injury and cell proliferation in the esophageal tissues and other parts of the alimentary tract following oral administration.

Mutagenicity

Florin et al. (1980) reported that 2,4-Hx was non-mutagenic when tested at a concentration of 3 μ M/plate in Ames/Salmonella tester strains TA98, TA100, TA1535, and TA1537. In the NCI/DCE Short-Term Test Program, 2,4-Hx was negative in the Ames/Salmonella mutagenicity assay using tester strains TA98, TA100, TA1535, TA1537, AND TA1538 when tested at doses up to 10,000 μ g/plate with and without metabolic activation. In the mouse lymphoma L5178Y TK+/- assay, 2,4-Hx was positive when tested at doses ranging from 3.1 to 0.6 μ g/ml without metabolic activation but was negative when tested at doses ranging from 74 to 23 μ g/ml with activation (NCI, 1992a).

Marnett et al. (1985) reported that 2,4-Hx tested positive for mutagenicity a liquid pre-incubation procedure measured in Ames/Salmonella assay using strain TA104, with a result of 960 revertants/μM. No mutagenicity was observed with strain TA102. authors found that 2,4-Hx was cytotoxic but the toxicity could be reduced by addition of glutathione. Both TA102 and TA104 are base substitution strains developed to detect peroxides and other oxidants. TA104 carries a nonsense mutation (-TAA-) at the site of revrsion that is present in single copy on the chromosome and TA102 contains the same histidine mutation on a multicopy plasmid. 2,4-Hx was also tested positive in TA100 without S9 in a modified preincubation Ames test (Eder et al., 1992; 1993). Positive results were obtained by increasing the bacterial number (3-fold) and by using glutathione as a chase. (Note: the molecular mechanisms leading to backmutation in strain TA104, which carries the his G428 mutation cannot be directly compared with those of the his G46 strains).

3

Eder et al. (1993) demonstrated that toxicity of 2,4-Hx interferes with mutagenicity testing in Salmonella strain TA100. But with higher cell numbers toxicity can be partly compensated and mutagenicity of 2,4-Hx without S9 was demonstrated in preincubation assays. In SOS chromotest using E. coli strain PQ37, 2,4-De was positive when ethanol was used as solvant, but was negative when dimethylsulfoxide was used (Eder et al., 1993). 2,4-Hx induced DNA strand breaks at high dose (500 μmole) as measured in the alkaline elution technique when cytoxicity was also observed. 2,4-Hx also interacted with guanine and formed DNA adducts (1,2-cyclic deoxzyguanosine adduct and 7,8-cyclic guanosine adduct) (Eder et al., 1993)

Strand breaks can be due to direct DNA interaction but also to cell death-associated release of endonucleolytic enzymes. Brambila et al. (1986) found that lipid peroxidation products can remain associated with the lipid core of the enoplasmatic reticulum and that they can diffuse into the cell nucleus without encountering cytotosolic detoxifying enzymes. Thus, relatively high concentrations can interact with DNA in vivo.

 α,β -unsaturated aldehydes have been shown to cause unscheduled DNA synthesis in primary cultures of rat hepatocytes (Griffin et al., 1986), DNA fragmentation and sister chromatid exchange in Chinese hamster ovary cells (Brambila et al., 1986), and inhibition of the DNA repair enzyme O⁶-methylguanine-DNA methyltransferase (Krokan et al., 1985).

DISPOSITION

Hydroperoxides and low-molecular weight secondary oxidation products of linoleic acid have been reported to be readily absorbed. After administration of a single dose of 14C-labelled methyl linoleate hydroperoxides by gastric intubation to rats, up to 80% of the dose were recovered mainly in tissues of the stomach and liver (Bergan and Draper, 1971, Kanazawa et al., 1985; Oarada et al., 1986). Studies of dienals have not been conducted.

Aldehydes are principally metabolized in the liver. Alcohol dehydrogenase is capable of catalyzing the metabolic reduction of aldehydes to primary alcohols in the reversal of the metabolic reaction involving the oxidation of primary alcohols to aldehydes. In mammals, however, this is not necessarily found to occur, since aldehydes are preferentially oxidized to the corresponding acids. The acids formed as major metabolites of aldehydes may then be excreted or form conjugates which are excreted. Aldehyde dehydrogenase, with NADH as cofactor, has been shown to dehydrogenate short chain aliphatic aldehydes as well as aromatic aldehydes (McMahon, 1982; Esterbauer et al., 1985).

Biological Effects

While free radical and lipoperoxides produce direct damage at the cell structures where they are produced, aldehydes are more diffusible and long-lived and may induce damage at distant sites.

1. Interact with DNA

 α,β -Unsaturated aldehydes ingested or produced endogenously as a result of lipid peroxidation during normal metabolic process or induced by exogenous chemicals (CCl4, CHCl3, DDT, PCB, etc) are strong electrophilic reagents and react readily with nucleophilic groups. Thus, 2,4-Hx is expected to interact with DNA.

Eder et al. (1993) demonstrated that members of the β -alkyl-substituted acrolein congeners (pentenal, hexenal, 3,3-dimethylacrolein) form 1,N²-cyclic adducts and 7,8-cyclic adducts with deoxyguanosine in vitro similar to those observed with crotoaldehyde. Their data showed that 2,4-Hx would formed similar adducts with deoxyguanosine although they could not isolate the adducts in sufficient quantity. They postulated that since these adducts are premutagenic DNA lesions and that crotonaldehyde is carcinogenic, the β -alkyl-substituted acrolein congeners are considered carcinogenic.

A fluorescence is generated when DNA interacts with a lipid degradation product. Using this technique Frankel et al. (1987) demonstrated that 2,4-De is readily interact with calf thymus DNA in vitro in the presence of ferric chloride and ascorbic acid.

2. Interact with glutathione

Administration of acrolein via the oral route cause a dose-dependent depletion of glutathione in the liver. GSH conjugation of acrolein mediated by glutathione transferases is considered a detoxicifcation mechanism. But evidence has been shown that conjugates may transport the chemical to be activated at a new site (Witz, 1989). Thus, the biological fate of thiol conjugates of α,β -unsaturated aldehydes remains to be explored.

3. Inhibition of cell proliferation (Carcinostatic/therapeutic /Inhibition of tumor cell growth)

Aldehydes have been shown to inhibit cell proliferation. (tumor cells are

more sensitive to aldehydes than normal cells, due to aldehyde dehydrogenase activity). The mechanism may involve interaction with tubulin, inhibition of polyamine metabolism, adenylate cyclase, lysosomes (Dianzani, 1982).

The sulfhydryl reactivity of α,β -unsaturated aldehydes may play a role in carcinostatic action, e.g. 1:2crotonaldehyde-cysteine adduct and the 1:1 trans-4-hydroxypentenal-cysteine adduct. The action is due to the reactivity of the double bond in interaction with sulfhydryls. The aldehydes may react with enzymes containing the essential sulfhydryl groups.

2,4-Hx was investigated for use as a food preservative but was found inactive in retarding the growth of food molds (fungi) (Troller and Olsen, 1967). On the other hand, Gueldner et al. (1985) reported that 2,4-Hx, occurring naturally in corn ears, appeared to act as a natural mycostatic insecticide, inhibiting growth of Aspergillus flavus.

4. Inhibition of enzymes (membrane bound enzymes?)

DNA repair enzyme O⁶-methylguanine-DNA methyl transferase by forming cysteine-trans-4-hydroxynonenal adduct.

The effect of α,β -unsaturated aldehydes on rat liver microsomal glucose 6-phsphatase has been studied. Depending on the chain-length, the Michaelis constant, K_m and the maximal rate of reaction, V, were affected. However, both 2,4-Hx and 2, 4-De did not alter the kinetic constants and K_m of the enzyme. The lack of effects may be attributed to the rather rigid planar structure around the 3 conjugated double bonds giving rise to a severe sterical hindrance at the β - and δ -carbon atoms (Jorgensen et al., 1992).

Other enzymes known to be inhibited: cytochrome P-450
aminopyrine demethylase
adenylate cyclase

5. Cytotoxicity

Cytotoxic effect of α,β-unsaturated aldehydes may be due to inhibition of superoxide anion radical production in macrophage and neutrophils or decrease in membrane lipid fluidity (see Witz, 1989). For examples, in a study of the effects of tobacco smoke components, Thelestam et al. (1980) found that 2,4-Hx at a concentration of 25 mM caused an increase of 20% in the membrane permeability of human lung fibroblasts incubated for 30 minutes. 2,4-Hx at a concentration of 0.01 mM in ethanol inhibited murine Ascites sarcoma BP8 cells 44% while 0.1 and 1.0 mM concentrations were 100% cytotoxic (Pilloti et al., 1975). 2,4-Hx at a concentration of 0.1 mM inhibited noradrenaline induced oxidative metabolism in isolated hamster brown-fat cells and at 1 mM caused 100% inhibition (Pettersson et al., 1980). A concentration of 5 mM 2,4-Hx applied to chicken embryo tracheal organ cultures for 6 minutes caused complete cessation of ciliary activity (Petterrson, et al., 1980).

6. α,β -Unsaturated aldehydes, including 2,4-Hx, react with thiobarbituric acid (TBA) forming a reddish pigment which is the basis of human clinical lipid peroxidation analysis. Kosugi *et al.* (1988) observed synergism between 2,4-alkadienals and other aldehydes and hydroperoxides as evidenced by the intensity of the red pigment formation.

Structure/Activity Relationships

The Interagency Testing Committee (ITC) has classified chemicals containing the closely related substructure, α,β -unsaturated aldehyde, as a group of chemicals likely to be associated with adverse health and ecological effects. Their concern for potential health effects resulting from exposures to this class of chemicals includes oncogenicity, mutagenicity and membrane irritation.

The known cancer risk of α, β -unsaturated aldehydes in the diet is summarized below:

Chemicals		Mutagen- icity	Carcino- genicity	Ref.
Acrolein	CH ₂ =CH-CHO	-,+w	Deferred	NTP
2-Chloroacrolein			+	Robinson et al.
2-Bromoacrolein			+	Robinson et al.
Malondialdehyde		-	MR ⁺ FR ⁺ MM ⁻ FM ⁻	NTP
нос-сн=сн-сно				
Crotonaldehyde		+	+	Chung et al., 1986
сн3-сн=сн-сно				(in drinking water)
Furfural		?	MR ⁺ FR ⁻ MM ⁺ FM ⁺	NTP
	СНО			·
Citral		-	on going	NTP
нзс-с=сн	I- (CH2) 2-C=CH-CHO			
I Сн3	 СНЗ			
vanillin		-		NTP
glycidaldehyde		+		NTP
			Animal carcin	nogen, IARC
benzaldehyde		-	MR ⁻ FR ⁻ MM ⁺ FM ⁺	NTP
2,4,5-Trimethoxybenzaldehyde		_	Deferred	NTP
cinnamaldehyde .		?	on going	NTP
C ₆ H ₅ -CH=0	Сн-сно			
Trimethoxycinnamaldehyde		+		
anisaldehyde				
Fecapentaenes		+	Assoc. human	Vertegaal et al., 1992
			colorectal CA	4
2,4-hexadienal		+		Eder et al., 1992
CH ₃ -CH=CH	-СН=СН-СНО			
2,4-Decadienal		+		Marnett et al., 1985
C ₅ H ₁₁ -CH=CH-CH=CH-CHO				Eder et al., 1993

MR: male rats; FR: female rats; MM: male mice, FM: female mice

The central structural feature of an α,β -unsaturated aldehyde is its bifunctional group, the carbon-carbon double bond conjugated with the aldehydic carbonyl group. α,β -unsaturated aldehydes are direct-acting alkylating agens capable of covalent binding without prior metabolism to cellular nucleophilic groups, the potential for being toxic and/or capable of modifying cellular processes is inherently present in every member of this class (Witz, 1989).

In studying mutagenesis of β -alky substituted acrolein congeners (crotonaldehyde, trans-2-pentenal, trans-2-hexenal, 2,4-hexadienal, and trans-2-heptenal) Eder et al. (1992) observed that toxicity increases as a function of both chain length and lipophilicity:

- 1. Increasing chain length increases toxicity and decreases mutagenicity This effect may be explained by increasing toxicity as mutagenic responses cannot be observed in some cases because of competing cytotoxicity (Eder et al., 1992).
- 2. Increasing β -chain length increases lipophilicity and toxicity. The dependence of toxicity on increasing β -chain length can be ascribed to increasing lipophilicity which allows a better penetration through the cell membrane.
- 3. The conjugated double bond in the β-alkyl chain (2,4-Hx) increases mutagenesis; this is also in accordance with the hypothesis that lipophilicity determines toxicity. (In in vitro studies only, what about in vivo?)
- 4. Decrease in mutagenicity with increasing molecular weight. This is due to decrease in reactivity of α,β -unsaturated carbonyl compounds towards nucleosides and nucleotides with increase molecular weight (Eder et al., 1993).

RECOMMENDATIONS

Since 2,4-Hx is mutagenic and cytotoxic and it has been shown that 2,4-Hx induced DNA strand breaks and forming adducts with guanine, its potential to induce cancer is high. However, it remains to be shown whether it can induce these events in vivo and ultimately cancer. The chemical is produced endogenously and humans are exposed to it mainly by mouth through food. The organism normally has a protective mechanism in detoxifying the chemical. In order to induce toxicity and carcinogenicity, the chemical has to be administered in large doses to overwhelm the natural detoxifying mechanism. The following studies are recommended:

- 1. Disposition studies be conducted. There is no information on absorption, toxicokinetics, metabolism, distribution, and excretion of 2,4-De in the literature.
- 2. Gavage studies in 0.5% methyl cellulose be conducted: 14-day and 90-day dose range finding studies and 2-year carcinogenesis studies.
- 3. Stability study be conducted: Stability of 2,4-De in 0.5% methyl cellulose.
- 4. The role of glutathione and thiols and vitamins C and E be investigated. It has been shown in vitro that the presence of glutathione reduces toxicity of a,b-unsaturated carbonyl compounds. Administration of 2,4-De may depletic liver glutathione and blood vitamins C and E.
- 5. Cell proliferation studies be conducted. Increased proliferative activity in the esophagus have been shown to occur after feeding with heated polyunsaturated oil (Hageman et al., 1991). Bull et al. (1984; 1988) and Craven et al. (1987) also showed that hydroperoxides of linoleic acids as well as other oxidative degradation products and metabolites of polyunsaturated fatty acids enhanced cell proliferation in the colonic tissue. Therefore, esophagus, stomach, colon, small and large intestine, mammary gland, kidney, and liver shall be processed for possible PCNA staining.

REFERENCES

Aikawa, K. and Chikuni, K. (1988) Antimutagenic effect of volatile decomposition products from thermally oxidized linoleate. Mutat. Res., 208(3-4):163-166

Aldrich (1991) Aldrich's 1991-1992 Flavors and Fragrances Catalog, Aldrich Chemical Co., Inc. Milwaukee, WI, p. 25

- 15 -

Aldrich (1992) Material Safety Data Sheet (MSDS). Aldrich Chemical Co., Inc., Milwaukee, WI

Alexander JC, Valli VE and Chanin BE (1987) Biological observations from feeding heated corn oil and heated peanut oil to rats. J. Txociol. Environ. Health, 21, 295-309.

Beppu, M., Murakami, K., and Kikugawa, K. (1986) Fluorescent and cross-linked proteins of human erythrocyte ghosts formed by reaction with hydroperoxylinoleic acid, malonaldehyde, and monofunctional aldehydes. Chem. Pharm. Bull., 34(2):781-788

Bergan JG and Draper HH (1971) Absorption and metabolism of 1-14-C-methyl linoleate hydroperoxide. Lipids, 5, 976-982.

Brambilla G, Sciaba L, Faggin P, Maura A, Marinari UM, Ferro M and Esterbauer H (1986) Cytotoxicity, DNA fragmentation and sister-chromatid exchange in Chinese hamster ovary cells exposed to the lipid peroxidation product 4-hydroxynonenal and homologous aldehydes. Muta. Res., 171, 169-176.

Brophy, P.M. and Barrett, J. (1990) Detoxification of secondary products of lipid peroxidation in the cytosol of a mouse fibroblast cell line. Biochem. Cell Biol., 68(11):1288-1291

Bull AW, Nigro ND, Golembieski WA, Crissman JD and Marnett LJ (1984) In vivo stimulation of DNA synthesis and induction of ornithine decarboxylase in rat colon by fatty acid hydroperoxides, autoxidation products of unsaturated fatty acids. Cancer Res., 44, 4924-4928..

Bull AW, Nigro ND and Marnett LJ (1988) Structural requirements for stimulation of colonic cell proliferation by oxidized faty acids. Cancer Res., 48, 1771-1776.

Canonero, R., Martelli, A., Marinari, U.M. and Brambilla, G. (1990) Mutation induction in Chinese hamster lung V79 cells by 5 alk-2-enals produced by lipid peroxidation. Mutat. Res., 244(2):153-156

Crescent Chemical Co. (1991) Heraeus/Crescent Catalog, Hauppauge, NY

Crawford L and Wheeler EL (1983) Microsomal P450 induction by some secondary products from thermal oxidation of dietary lipids: epidermal hyperplasia, mutagenicity and cytochrome P-450. cancer Letters, 21, 211-217.

Dianzani MU (1982) Biochemical effects of saturated and unsaturated aldehydes. In:Free Radicals, Lipid Peroxidation and Cancer. DCH McBrien and TF Slater. New York: Academic Press, 1982. pp129-158.

Dumdei, B.E., Kenny, D.V., Shepson, P.B., Kleindienst, T.E., Nero, C.M., Cupitt, L.T. and Claxton, L.D. (1988) MS/MS analysis of the products of toluene photooxidation and measurement of their mutagenic activity. *Environ. Sci. Technol.*, 22(12):1493-1498

Eder E, Deininger C, Neudecker T and Deininger D (1992) Mutagenicity of Balkyl substituted acrolein congeners in the salmonella typhimurium strain TA100 and genotoxicity testing in the SOS chromotest. Environ Mol Mutag., 19, 338-45.

Eder E, Scheckenbach S, Deininger C and Hoffman C (1993) The possible role of α, β -unsaturated carbonyl compounds in m utagenesis and carcinogenesis. Toxicol. Letters, 67, 87-103.

FDA (1992) Center for Food Safety and Applied Nutrition, Priority-based assessment of food additives (PAFA) database, Washington, D.C. [personnel communication between Dorothy Cannon of TRI and Dan Benz of FDA, March, 1992]

Feron, V.J., Til, H.P., de Vrijer, F., Woutersen, R.A., Cassee, F.R., and van Bladeren, P.J. (1991) Aldehydes: occurrence, carcinogenic potential, mechanism of action and risk assessment. Mutat. Res., 259(3-4):363-385

Ferrario, J.B., Lawler, G.C., DeLeon, I.R., and Laseter, J.L. (1985) Volatile organic pollutants in biota and sediments of Lake Pontchartrain, Bull. Environ. Contam. Toxicol., 34(2):246-255

- Florin, I., Rutberg, L., Curvall, M., and Enzell, C.R. (1980) Screening of tobacco smoke constituents for mutagenicity using the Ames' test. Toxicol., 18:219-232
- Ford, R.A., Letizia, C. and Api, A.M. (1988) Monographs on fragrance raw materials: trans, trans-2,4-hexadienal. Food Chem. Toxicol., 26(4):337-338
- Frankel, E.N., Neff, W.E., Brooks, D.D., and Fujimoto, K. (1987) Fluorescence formation from the interaction of DNA with lipid oxidation degradation products. Biochem. Biophys. Acta., 919(3):239-244
- Griffin DS, Segall HJ (1986) Lipid peroxidation and cellular damage caused by the pyrrolizidine alkaloid senecionine. In: Free Radicals. Cell Damage and Disease, Rice-Evans, ed. London: The Richelieu Press, pp. 201,220.
- Growcock, F.B., Frenier, W.W. and Andreozzi, P.A. (1989) Inhibition of steel corrosion in HCl by derivatives of cinnamaldehyde. Part II. Structure-activity correlations. Corrosion, 45(12):1007-1015
- Gueldner, R.C., Wilson, D.M., and Heidt, A.R. (1985) Volatile compounds inhibiting Aspergillus flavus. J. Agri. Food Chem., 33(3):411-413
- Gupta, I., Suzuki, K., Bruce, W.R., Krepinsky, J.J., and Yates, P. (1984) A model study of fecapentaenes: mutagens of bacterial origin with alkylating properties. Science, 225(4661):521-523
- Hageman, G., Verhagen, H., Schutte, B. and Kleinjans, J. (1991) Biological effects of short-term feeding to rats of repeatedly used deep-frying fats in relation to fat mutagen content. Fd. Chem. Toxic. 29(10):689-698
- Heil, T.P. and Lindsay, R.C. (1988) Volatile compounds in flavor-tainted fish from the Upper Wisconsin River. J. Environ. Sci. Health, Part B, B23(5):489-512 [STN abstract, CA110(11):93708q]
- Izaki Y, Yoshikawa S and Uchiyama M (1984) Effect of igestion of thermally oxidized frying oil on peroxidative criteria in rats. Lipids, 5, 324-331.
- Jorgensen BM, Agerbo P, Jensen B, Borresen T and Holmer G (1992) Inhibition of microsmal glucose 6-phosphatase by unsaturated aliphatic aldehydes and ketones. Chem-Biol. Interactions, 81, 209-218.

Kanazawa K, Kanazawa E and Natake M (1985) Uptake of secondary autoxidation products of linoleic acid by the rat. Lipids, 20, 412-419.

Kanazawa K, Ashida H, Minamoto S and Natake M (1986) The effect of orally administered secondary autoxidation products of linoleic acid on the activity of detoxifying enzymes in the rat liver. Biochim. Biophys. Acta, 879, 36-43.

Kaneko, T., Honda, S., Shun-Ichiro, N., and Matsuo, M. (1987) Lethal effects of a linoleic acid hydroperoxide and its autooxidation products unsaturated aliphatic aldehydes on human diploid fibroblasts. Chem.-Bio. Interact., 63(2):127-138

Kaneko, T., Kaji, K., and Matsuo, M. (1988) Cytotoxicities of a linoleic acid hydroperoxide and its related aliphatic aldehydes toward cultured human umbilical vein endothelial cells. Chem.-Biol. Interact., 67(3-4):295-304

Katsuki, Y., Matsumoto, S., and Tsuyuki, H. (1987) Air oxidation of unsaturated triglycerides and prevention of oxidation. II. Volatile decomposition products in the air oxidation of triolein and trilinolein. Shokuhin Eiseigaku Zasshi, 28(6):466-472 [STN abstract, CA109:5391g]

Kawakami, M. and Shibamoto, T. (1991) The volatile constituents of piled tea: Toyama kurocha. Agric Biol. Chem., 55(7):1839-1847

Keller, C.L., Balaban, S.M., Hickey, C.S. and DiFate, V.G. (1983) Sorbic acid. In: Mark, H.F., Othmer, D.F., Oveberger, C.G., Seaborg, G.T. & Grayson, M., Eds., Kirk-Othmer Encyclopedia of Chemical Technology, 3rd Ed., Vol. 21, John Wiley and Sons, Inc., New York, NY, p. 406

Kosugi, H. and Kikugawa, K. (1989) Enhancement of red pigment formation in thiobarbituric acid reaction with a combination of alkenals, alkadienals, and organic hydroperoxides. Yukagaku, 38(3):224-230 [STB abstract, CA111(6):41341r]

Kosugi, H., Kato, T, and Kikugawa, K. (1988) Formation of a red pigment by a two-step 2-thiobarbituric acid reaction of alka-2,4-dienals. Potential products of lipid oxidation. Lipids, 23(11):1024-1031 [STN abstract, CA110(7):53977v]

Kosugi, H., Kojima, T. and Kikugawa, K. (1991) Characteristics of the thiobarbituric acid reactivity of oxidized fats and oils. J. Am. Oil Chem. Soc., 68(1):51-55 [STN abstract, CA114(13):120453u]

Krokan H, Grafstrom RC, Sundqvist K, Esterbauer H, Harris L (1985) Cytotoxicity, thiol depletion and inhibition of O6-mekthylguanine DNA methyltransferase by various aldehydes in cultural human bronchial fibroblasts. Carcinogenesis, 6, 1755-1759.

Lancaster Synthesis, Inc. (1991) Lancaster Catalogue 1991/92, MTM Research Chemicals, Windham, NH

Marnett, L.J., Hurd, H.K., Hollstein, M.L., Levin, D.E., Esterbauer, H., and Ames, B.N. (1985) Naturally occurring carbonyl compounds are mutagens in Salmonella tester strain TA104. Mutat. Res., 148(1-2):25-34

McMahon, R.E. (1982) Alcohols, aldehydes, and ketones. In: Jakoby, W.B., Bend, J.R. and Caldwell, J. Eds., *Metabolic basis of detoxication: metabolism of functional groups*, Academic Press, New York

NCI (1992a) NCI/DCE Short-term testing program, in vitro results, 3/81-8/92. National Cancer Institute, Bethesda, MD

NCI (1992b) NCI tracking file, National Cancer Institute, Bethesda, MD

NLM (1992) National Library of Medicine, Toxicology Data Network (TOXNET) information Services, databases searched January to July 1992

NTP (1992) NTP Results Report: Results and Status Information on All NTP Chemicals. January, 1992, Report

O'Brien, P.J., Kaul, H., McGirr, L., Drolet, D., and Silva, J.M. (1989) Molecular mechanisms for the involvement of the aldehydic metabolites of lipid peroxides in cytotoxicity and carcinogenesis. In: Kabara, J.J., Ed. *Pharmacological Effects of Lipids 3*, AOCS, Champaign, IL, pp. 266-280

Oarada M, Miyazawa T and Kaneda T (1986) Distribution of 14C after oral administration of (U-14C) labeled methyl linoleate hydjroperoxides and their secondary oxidation products in rats. Lipids, 21, 150-154.

Pettersson, B., Curvall, M. and Enzell, C.R. (1980) Effects of tobacco smoke compounds on the noradrenaline induced oxidative metabolism in isolated brown fat cells. Toxicol., 18(1):1-15

Pilotti, A., Ancker, K., Arrhenius, E. & Enzell, C. (1975) Effects of tobacco smoke constituents on cell multiplication in vitro. Toxicol., (5):49-52

Povey, A.C., Schiffman, M., Taffe, B.G., and Harris C.C. (1991) Laboratory and epidemiology studies of fecapentaenes. Mutat. Res., 259:387-397

Przybylski, R. and Hougen, F.W.. (1989) Simple method for estimation of volatile carbonyl compounds in edible oils and fried potato chips. J. Am. Oil Chem. Soc., 66(10):1465-1468

Selke, E. and Rohwedder, W.K. (1983) Volatile components from trilinolenin heated in air. J. Am. Oil Chem. Soc., 60(11):1853-1858

Sherman, P.D., Jr. and Kavasmaneck, P.R. (1980) Ethanol. In: Mark, H.F., Othmer, D.F., Overberger, C.G., Seaborg, G.T., and Grayson, M., eds., Kirk-Othmer Encyclopedia of Chemical Technology, 3rd Ed. Vol. 9, John Wiley and Sons, Inc., New York, NY, p. 355

Snyder, J.M., Frankel, E.N. and Selke, E. (1985) Capillary gas chromatographic analyses of headspace volatiles from vegetable oils. J. Am. Oil Chem. Soc., 62(12):1675-1679

STN (1992) Chemical Abstracts Service databases (file CA), STN International, Columbus, Ohio

Sturmer, D.M. and Diehl, D.R. (1982) Polymethine dyes. In: Mark, H.F., Othmer, D.F., Overberger, C.G., Seaborg, G.T. and Grayson, M., Eds., Kirk-Othmer Encyclopedia of Chemical Technology, 3rd Ed. Vol. 18, John Wiley and Sons, Inc., New York, NY, p. 866

Suzuki, J. and Bailey, M.E. (1985) Direct sampling capillary GLC analysis of flavor volatiles from ovine fat. J. Agric. Food Chem., 33(3):343-347 [STN abstract, CA102(23):202676k]

Takeoka, G.R., Giintert, M., Flath, R.A., Wurz, R.E. and Jennings, W. (1986) Volatile constituents of kiwi fruit (*Acinidia chinesis* Plauch). J. Agric. Food Chem., 34(3)576-578

Thelestam, M., Curvall, M. and Enzell, C.R. (1980) Effect of tobacco smoke on the plasma membrane of cultured human lung fibroblasts. Toxicology, 15(3):203-17

Troller, J.A. and Olsen, R.A. (1967) Derivatives of sorbic acid as food preservatives. J. Food Sci., 32(2):228-231

Urbany, D. (1992) State cancer rate linked to lifestyles. Gazette Newspapers, Bethesda, MD, March 4, 1992, p.A-10

Vertegaal, L.B.J., Voogd, C.E., Mohn, G.R. and van der Gen, A. (1992) Further studies on the mutagenic activity of fecapentaene-12 analogues and conclusions about their molecular mode of action. Mutat. Res., 281:93-98

Weeks, W.W., Chaplin, J.F. and Campbell, C.R. (1989) Capillary chromatography: evaluation of volatiles from flue-cured tobacco varieties. J. Agric. Food Chem., 37(4):1038-1045

White, P.J. and Hammond, E.G. (1983) Quantification of carbonyl compounds in oxidized fats as trichlorophenylhydrazones. J. Am. Oil Chem. Sec., 60(10):1769-1773

Williamson, G. (1989) Purification of glutathione-S-transferase from lamb muscle and its effect on lipid peroxidation. J. Sci. Food Agric., 48(3):347-360

Williamson, G. and Ball, S.K.M. (1988) Purification of glutathione-S-transferase from lean pork muscle and its reactivity with some lipid oxidation products. J. Sci. Food Agric., 44(4):363-374

Witz, G., Lawrie, N.J., Zaccaria, A., Ferran, Jr., H.E., and Goldstein, B.D. (1986) The reaction of 2-thiobarbituric acid with biologically active alpha, beta-unsaturated aldehydes. J. Free Rad. Biol. Med., 2:33-39

Witz G (1989) Biological interactions of a,b-unsaturated aldehydes. Free Radical Biology Med., 7, 333-349.

Wright, D.H. and Harris, N.D. (1985). Effect of nitrogen and potassium fertilization on tomato flavor. J. Agric. Food Chem., 33(3):355-358

Zeringue, H.J. and McCormick, S.P. (1989) Relationships between cotton leaf-derived volatiles and growth of A. flavus. J. Am. Oil Chem. Soc., 66(4):581-585