

## Erratum

TABLE J2

Tissue Concentrations and Excreta Content of Chromium in Female Mice in the 2-Year Drinking Water Study of Sodium Dichromate Dihydrate

Bolded numbers are corrected values.

	0 mg/mL	14.3 mg/mL	57.3 mg/mL	172 mg/mL	516 mg/mL
n	3	3	3	3	3
Feces (µg) <sup>c,d</sup>					
Day 6	2.599 ± 0.899	<b>9.382 ± 1.405</b>	<b>13.032 ± 2.779</b>	<b>43.741 ± 10.325</b>	<b>162.632 ± 31.032</b>
Day 13	2.348 ± 1.173	12.103 ± 3.355	<b>85.804 ± 21.392</b>	<b>46.934 ± 4.884</b>	<b>109.687 ± 29.885</b>
Day 182	1.275 ± 0.569	<b>5.217 ± 0.471</b>	<b>17.606 ± 3.100</b>	<b>24.624 ± 5.736</b>	<b>122.928 ± 21.563</b>
Day 371	7.485 ± 2.863	<b>9.505 ± 1.569</b>	<b>18.719 ± 2.172</b>	<b>41.017 ± 3.214</b>	<b>72.197 ± 19.151</b>

<sup>c</sup> Cumulative chromium content for 48 hours ending on the day indicated

<sup>d</sup> Measured values less than the experimental limit of quantitation (ELOQ) (2.00 µg Cr/g feces; 0.100 µg Cr/g urine) were included in the analysis as ½ ELOQ x sample mass.