NTP Executive Committee Working Group for the Report on Carcinogens (RG2) Commentary on Human Exposure to Cobalt Sulfate

At the May 1, 2002 RG2 review of the cobalt sulfate nomination for listing in the RoC the RG2 recommended that the human exposure information presented in Section 2 of the cobalt sulfate background document be revised so that the human exposure information could be updated using the information in the most recent ATSDR Toxicological Profile for Cobalt (September 2001). It was also recommended that this section be reorganized according to information specific for cobalt sulfate followed by information on cobalt compounds. There is more information for exposure to specific cobalt compounds. The RG2 wanted the information to be presented so that it clearly defined what type of information (e.g. production, occupational exposure) was available specifically for cobalt sulfate and generally for cobalt compounds.

This commentary is not meant to be a new draft of Section 2 of this background document. The information on human exposure to cobalt sulfate specifically and cobalt compounds in general has been updated to include any new information contained in the September 2001 ATSDR Toxicological Profile for Cobalt and reorganized so that information specific for cobalt sulfate is clearly discernable from information on cobalt compounds as a class.

Human Exposure

Information on human exposure has been divided into information specific for cobalt sulfate and information on cobalt compounds.

1 Cobalt sulfate

1.1 Use

Cobalt sulfate is used in the electroplating and electrochemical industries, where it is added to nickel plating baths in order to improve the smoothness, brightness, hardness, and ductility of the deposits. It also is used as a drier for lithographic inks, varnishes, paints, and linoleum and in storage batteries. Cobalt sulfate is employed as a coloring agent in ceramics, enamels, and glazes to prevent discoloring and as a co-pigment for decorating porcelain. In addition, cobalt sulfate has been used in animal feeds as a mineral supplement (Budavari *et al.* 1996, Kirk and Othmer 1999).

Cobalt sulfate has been mixed, in small quantities, with fertilizers for use on pastures where the forage is cobalt deficient, to provide enough cobalt for ruminants to produce vitamin B_{12} . In the United States in 1996, the total amount of fertilizer consumed containing cobalt sulfate was two tons. All of the fertilizer use was in Washington State, where the highest concentrations of cobalt in fertilizers were 44.8 to 222 mg/kg (dry weight) (EPA 1999, Washington State 1999).

In the 1960s, some breweries added cobalt sulfate to their beers to improve the stability of the foam by counteracting the antifoaming activity of detergent residues left on poorly rinsed glasses. Although only a small amount (1 ppm) was used in the beer, this practice was stopped after an epidemic of "beer drinker's cardiomyopathy" was linked to the cobalt (NTP 1998).

Cobalt sulfate has also been used in veterinary medication to prevent and treat cobalt deficiency in ruminants, which causes reduction in feed intake and body weight, accompanied by emaciation, anemia, and debility. Cobalt sulfate had been used in the past to improve hematocrit, hemoglobin, and erythrocyte levels in human patients with refractory anemia, including sickle-cell disease, thalassemia, chronic infection or renal disease, anemia associated with neoplastic disease, and various other refractory anemias of unknown cause. In 1985, cobalt was used clinically only in the treatment of normochromic, normocytic anemia associated with severe renal failure (HSDB 2000, Hillman and Finch 1985). There is no listing for cobalt or cobalt sulfate in the current *Goodman & Gilman's Pharmacological Basis of Therapeutics* (Goodman and Gilman 2001).

1.2 Production

Cobalt sulfate is formed by the interaction of cobalt oxide, hydroxide, or carbonate with sulfuric acid. Production of cobalt sulfate in the United States in 1983 was estimated at 450,000 lb (NTP 1998). Current production levels are not available. There are currently 11 U.S. suppliers of cobalt sulfate (ChemFinder 2001).

Import of cobalt sulfate in 1986 was reported to be 79,700 lb (HSDB 2000). U.S. imports for consumption of cobalt sulfates were 1,360 metric tons in 1999 and 1,040 metric tons in 1998, valued at \$9,840,000 and \$10,400,000, respectively. No specific information on cobalt sulfate exports was identified (USGS 2001).

Chem Sources identified 15 suppliers of cobalt(II) sulfate, four suppliers of cobalt(II) sulfate monohydrate, and 16 suppliers of cobalt(II) sulfate heptahydrate in the United States (Chem Sources 2001). The Hazardous Substances Data Bank listed seven manufacturers of cobaltous sulfate (HSDB 2000).

1.3 Analysis

No information was found that specifically identified analysis for cobalt sulfate.

1.4 Environmental occurrence

No information was found that specifically identified environmental occurrence to cobalt sulfate.

1.5 Environmental fate

Seawater formation of cobalt sulfate is not estimated because of the high concentration of chloride ion. No other information was found that specifically identified environmental fate to cobalt sulfate.

1.6 Environmental exposure

No information was found that specifically identified environmental exposure to cobalt sulfate.

1.7 Occupational exposure

No information was found that specifically identified occupational exposure to cobalt sulfate.

1.8 Biological indices of exposure

Cobalt sulfate, like other water-soluble metallic salts, dissolves directly into blood serum (362 g/L at 20°C) (Jensen and Tüchsen 1990). Cobalt can be detected in urine, blood, and tissues; however, there currently is no way to correlate cobalt sulfate exposure with cobalt levels observed in these matrices.

1.9 Regulations

No specific U.S. Environmental Protection Agency (EPA) regulations for cobalt sulfate were identified.

2 Cobalt compounds as a class

2.1 Use

The largest use of metallic cobalt is in super alloys that are used in gas turbine aircraft engines. Superalloys are alloys developed for applications where elevated temperature and high mechanical stress are encountered. Cobalt compounds are used as pigments in glass, ceramics, and paints; as catalysts in the petroleum industry; as paint driers; an as trace element additives in agriculture and medicine.

2.2 Production

The United States did not mine or refine cobalt in 2000, although negligible amounts of cobalt were produced as a byproduct of mining operations. The U.S. supply of cobalt in 2000 included imports, stock releases, and secondary materials. Stock releases originated from the U.S. government reserve (National Defense Stockpile) for military, industrial, and civilian use during a national emergency. Sales of the National Defense Stockpile of cobalt began in March of 1993. Seven companies were known to be active in the production of cobalt compounds. It was estimated that 45% of U.S. cobalt usage was in superalloys, 9% in cemented carbides, 9% in magnetic alloys, and the remaining 37% in various other metallic and chemical uses (USGS 2001). Table D-1 summarizes recent patterns of cobalt production, import, export, and consumption in the United States.

	Metric tons of cobalt	
Salient statistics	1999	2000
United States:		
Production:		
Mine	NR	NR
Secondary	2,720	2,800
Consumption:		
Reported	8,420	8,400
Apparent	10,700	10,900
Imports for consumption	8,150	8,000
Exports	1,550	2,300
World production:		
Mine	29,900	32,300
Refinery	31,200	NR

Table 1-1. Cobalt production, consumption, import, and export

Sources: Shedd 1999, USGS 2001.

NR = not reported.

Reported 1999 U.S. cobalt consumption was 2,530 metric tons for chemical and ceramic uses and 64 metric tons for miscellaneous and unspecified uses. Reported 1999 U.S. consumption of cobalt chemical compounds (organic and inorganic) was 1,910 metric tons. Imports of cobalt sulfates and other cobalt salts (acetates, carbonates, and chlorides) from 10 countries totaled \$12,400,000. Most imports were from Finland. The United States exported \$49,700,000 of cobalt and cobalt compounds in 1999. Table D-2 summarizes U.S. cobalt consumption patterns in early 2001.

Consumption information	Date	Metric Tons	Compounds and uses
Reported consumption of cobalt materials	Jan–May	669	oxide and other chemical compounds
Reported consumption of cobalt by end use	Jan–May	883	chemical uses including catalysts, driers in paints, feed or nutritive additive, glass decolorizer, ground coat frit, pigments, other uses
Reported consumption of cobalt by end use	Jan–May	127	miscellaneous and unspecified uses
Imports by consumption	Jan–April	498	salts and compounds including acetates, carbonates, chlorides, and sulfates
Exports	Jan–April	74	salts and compounds

Table 1-2. Patterns of cobalt consumption in the United States in 2001

Source: USGS 2001.

2.3 Analysis

Determination of cobalt, especially in biological samples containing low levels of cobalt, is accurate only if samples are not contaminated. Contamination from disposable syringes and technical-grade anticoagulants was responsible for erroneous reports in earlier literature of grossly high levels of cobalt in biological specimens. The common classical methods used for determining cobalt concentration in biological samples are polarographic and colorimetric methods. However, these older methods are unsuitable for determining low levels of cobalt in many biological samples, and samples must be chemically pretreated before quantification. The most common single-element instrumental techniques used are electrothermal atomic absorption spectrometry (AAS) and voltammetric techniques (ATSDR 2001). Analytical methods for determining cobalt in biological matrices are summarized in Table 2-3. The samples analyzed in the studies presented in this table were primarily from cobalt-exposed and non-exposed workers (Heinrich and Angerer 1984, Ichikawa et al. 1985, Alexandersson 1988). However, one study used samples from laboratory volunteers (Bouman *et al.* 1986), and another used hospital patients with knee or hip prostheses (Sunderman et al. 1989). IARC (1991) reported that serum cobalt concentrations in humans were in the range of 0.1 to 0.3 μ g/L. As shown in Table 2-3, the detection limit for cobalt in serum by direct injection into electrothermal AAS with Zeeman background correction is 0.02 µg/L.

Matrix	Analytical method	Detection limit
Urine	electrothermal Atomic Absorption Spectroscopy (AAS) with Zeeman background correction — direct injection	0.3 μg/L 0.1 μg/L
	electrothermal AAS with Zeeman background correction — chemical preparation	2.4 μg/L
	electrothermal AAS with deuterium background correction — chemical preparation	0.1 μg/L
	differential pulse cathodic stripping voltametry (DPCSV) — chemical preparation	0.2 μg/L
Whole blood	electrothermal AAS with deuterium background correction	2 μg/L
	DPCSV — chemical preparation	0.8 µg/L
	colorimetry — chemical preparation	0.15 μg/L
Serum	electrothermal AAS with Zeeman background correction — direct injection	0.02 μg/L
Blood	inductively coupled plasma-atomic emission spectrometry (ICP-AES) — chemical preparation	10 µg/g
Tissue	ICP–AES — chemical preparation	0.2 µg/g

 Table 1-3. Analytical methods for determining cobalt in biological materials

Source: ATSDR 2001.

Because of its rapidity, accuracy, and low detection limit, electrothermal AAS with Zeeman background correction is the method most commonly used to quantify cobalt levels in environmental samples. To meet detection limits of some of the analytical methods, preconcentration may be necessary for some environmental samples (e.g., seawater). As with biological samples, contamination of environmental samples during collection, storage, and treatment are concerns (ATSDR 2001). Analytical methods for determining cobalt in environmental samples are detailed in Table 4.

Matrix	Analytical method	Detection limit	Recovery ^a
Air (workroom)	□-spectrometry with lithium-drifted germanium detector	0.17 µg/m ³	_
Air (occupational)	flame AAS with background correction	$0.4 \ \mu g/m^3$	98% with 12- to 96-µg spiked filter
	ICP–AES	0.5 µg/m ³	95%–100% with 2.5- to 1,000-μg spiked filter
Water (low ionic strength)	electrothermal AAS with Zeeman or deuterium background correction	< 0.5 µg/L	93%–115% at 8.5–30 µg/L
Lake water	ICP-AES	$< 0.004 \ \mu g/L$	_
Rainwater	photon-induced X-ray emission	0.08 µg/L	-
Seawater	electrothermal AAS with Zeeman background correction	0.0002 μg/L	90%
Water and waste water	flame AAS	0.05 mg/L	97%–98% at 0.2– 5.0 mg/L
	electrothermal AAS with background correction	1 µg/L	_
Groundwater or leachate	flame AAS with background correction	0.05 mg/L	97%–98% at 0.2– 5.0 mg/L
Groundwater or leachate	electrothermal AAS with background correction	1 µg/L	_
Food	electrothermal AAS with background correction	1.88 µg/L	100%-107% at 0.2-0.6 mg/kg in leaves and liver

 Table 1-4. Analytical methods for determining cobalt in environmental samples.

Source: ATSDR 2001.

 a^{-} = no data available.

2.4 Environmental occurrence

2.4.1 Air

Sources of cobalt in the atmosphere are both natural and anthropogenic. Natural sources include wind-blown continental dust, seawater spray, volcanoes, forest fires, and continental and marine biogenic emissions. The worldwide emissions from natural sources have been estimated to range from 13 to 15 million pounds per year. Cobalt in the atmosphere probably exists in particulate form (ATSDR 2001).

In the United States, the average ambient atmospheric concentration of cobalt was reported to be approximately 0.4 ng/m³ (ATSDR 2001). Although the HSDB (2000) described the atmospheric concentration of cobalt in remote areas as being very low, the only value given was less than 1 ng/m³ for the Antarctic. The same source reported that the air concentration of cobalt can reach or exceed 81 ng/m³ in heavily industrialized cities. Near a beryllium-copper alloy facility, cobalt levels as high as 610 ng/m³ were observed (HSDB 2000).

Atmospheric cobalt concentrations are much higher near cobalt manufacturing and production facilities. In the ambient air of a facility that manufactured cobalt salts, cobalt concentrations measured by personal sampling ranged from 0.1 to 3.0 mg/m³, with a mean of 0.2 mg/m³, and mean concentrations measured by stationary sampling were 0.049 and 1.046 mg/m³. The cobalt concentration in the ambient air during painting of pottery with soluble cobalt salts ranged from 0.07 to 8.61 mg/m³ (HSDB 2000). In Illinois, levels of cobalt in urban air ranged from 0.0025 to 0.010 mg/m³.

2.4.2 Water

Concentrations of cobalt in uncontaminated freshwater have been reported to range from 0.1 to 10 μ g/L (IARC 1991). The geometric mean and median cobalt levels from 6,805 ambient surface water stations were 2.9 and 2.0 μ g/L, respectively. The average concentration in seawater ranged from 0.078 μ g/L in the Caribbean Sea to 0.39 μ g/L in the Indian Ocean (ATSDR 2001).

The National Community Water Supply Study found that cobalt concentrations in drinking water in the United States ranged from nondetectable to 19 μ g/L, with 62% of the water samples containing a concentration less than 1 μ g/L. The average cobalt concentration in drinking water was 2.2 μ g/L (ATSDR 2001).

2.4.3 Soil

The average concentration of cobalt is 20-25 mg/kg in the earth's crust, 18 mg/kg in igneous rocks, and 7.2 mg/kg in U.S. soils. Soils with cobalt concentrations less than 3 mg/kg are considered cobalt deficient, because plants that grow in these soils will not contain enough cobalt to meet the dietary needs of cattle and sheep (< 0.08 - 0.1 mg/kg). Soils near ore deposits, phosphate rocks, ore traffic sites, or industrial pollution sites have been reported to contain cobalt at concentrations of up to 800 mg/kg (ATSDR 2001).

2.5 Environmental fate

2.5.1 Air

It is generally accepted that anthropogenic cobalt originating from combustion sources would primarily be in the form of the oxide, whereas cobalt arsenide and sulfide could be released during ore extraction processes. Very few data, however, were available on the potential transformation of these forms to other chemical species, such as the sulfate. It is speculated that chemical speciation of cobalt oxide in the air could lead to the formation of more-soluble forms such as cobalt sulfate, which would result in greater quantities being washed out of the atmosphere in rain (ATSDR 2001)

2.5.2 Water

Many factors will affect the speciation and transport of cobalt in natural waters and sediments. Dissolved cobalt appears to be precipitated in the adsorbed state with oxides of iron and manganese and with crystalline sediments such as aluminosilicate and goethite. In addition, cobalt precipitates out as carbonate and hydroxide in water (ATSDR 2001). In freshwater, it is estimated that speciation may yield 76% free Co^{+2} , 9.8% carbonate, 9.6% bicarbonate, 4% humic complexes, and 0.4% cobalt sulfate. Species of cobalt in seawater are $CoCl^+$, free Co^{+2} , carbonate, or carbonate and sulfate. Organic waste concentration and pH play an important role in cobalt speciation (ATSDR 2001). Bioconcentration of cobalt in marine fish is expected to occur, with bottom-feeders accumulating high levels of cobalt (HSDB 2000, ATSDR 2001).

2.5.3 Soil

The speciation of cobalt is regulated primarily by pH, the concentration of chelating or complexing agents in the soil, and the redox potential of the soil. At low pH, cobalt is oxidized to trivalent cobalt and usually is associated with iron. In the process of weathering, cobalt is readily taken into solution. It also is adsorbed to a great extent by hydrolysate or oxidate sediments (HSDB 2000).

2.6 Environmental exposure

Exposure to cobalt occurs through inhalation of ambient air and ingestion of food and drinking water. The average intake of cobalt in foods by adults in Canada was 11 μ g/day, with the intake varying between 4 and 15 μ g/day for various age groups. The median concentration of cobalt in U.S. drinking water is less than 2.0 μ g/L and the average concentration in ambient air is approximately 0.4 ng/m³ (ATSDR 2001). The major source of cobalt is food, in the form of green leafy vegetables, which may contain as much as 0.5 mg/kg dry weight (HSDB 2000). Cobalt is an essential trace element in humans, because a cobalt atom is present in each molecule of vitamin B₁₂ (cobalamin). Cobalt's presence in vitamin B₁₂ is its only known essential function in humans (Anderson 2000). An adult human body contains approximately 1.1 mg of cobalt (NTP 1998).

The National Health and Nutrition Examination Survey in 1999 measured cobalt levels in the urine of 1,007 participants aged 6 years or older, to provide physicians with a reference range of cobalt in the urine of the U.S. population for use in determining whether individuals have been exposed to cobalt (CDC 2001). The geometric mean was 0.36 μ g/L of urine (95% CI = 0.32 to 0.40). The geometric mean of the creatinine-adjusted levels was 0.33 μ g/g of creatinine (95% CI = 0.29 to 0.36).

2.7 Occupational exposure

It has been estimated by Jensen and Tüchsen (1990) that more than a million workers in the United States potentially are exposed to cobalt or cobalt compounds, though for many, the degree of potential exposure is limited (HSDB 2000, NTP 1998). Occupational exposure to cobalt occurs principally in refining processes, in production of alloys, and in the tungsten carbide hard-metal industry (Kazantzis 1981). In addition, many workers are exposed to a limited degree when using cobalt-containing paint dryers. Occupational exposure is primarily dermal or through inhalation of cobalt metal dusts or fumes (HSDB 2000, NTP 1998).

Cobalt metal has been reported in the air of metal manufacturing, welding, and grinding factories at concentrations ranging from 1 to $300 \ \mu g/m^3$ and in the dust of an electric welding factory at 4.2 $\mu g/g$ (ATSDR 2001). Occupational exposure to cobalt also has been assessed from the concentrations of cobalt in workers' tissues and body fluids. Alexandersson (1988) reported a high degree of conformity between the concentration of cobalt in blood and urine and the average levels of cobalt in the air during a workweek. In workers exposed to high levels (0.09 mg/m³), cobalt concentrations in the blood were 20 times those of the control group, while in the low-exposure workers (0.01 mg/m³), the concentrations were only slightly higher than in the controls. Other studies have shown that lungs from occupationally exposed workers, such as coal miners and metal-industry workers, contained from 2.5 to 6 times as much cobalt as lungs from control groups (ATSDR 2001).

2.8 Biological indices of exposure

Cobalt can be detected in urine, blood, and tissues; however, there currently is no way to correlate cobalt sulfate exposure with cobalt levels observed in these matrices. Based on reports of accidental exposure to radioactive cobalt (60 Co) and intravenous or oral administration of 60 Co to volunteer human subjects (Smith *et al.* 1972), approximately 90% of inhaled, injected, or ingested cobalt is eliminated within a few days; however, the remaining 10% has a half-life in the body of two years after parenteral administration or 5 to 15 years after inhalation. No biological use of cobalt is known other than its presence in vitamin B₁₂ (HSDB 2000).

2.9 Regulations

See cobalt sulfate background document, Section 2.9

References

- 1. Alexandersson, R. 1988. Blood and urinary concentrations as estimators of cobalt exposure. *Arch Environ Health* 43:299-303.
- 2. Anderson, J.J.B. 2000. Minerals. In Food, Nutrition, & Diet Therapy. Mahan, L.K. and S. Escott-Stump, eds. W. B. Saunders Company, Philadelphia. pp. 110-152.
- 3. ATSDR. 2001. Toxicological Profile for Cobalt. Draft. Agency for Toxic Substances and Disease Registry, U.S. Public Health Service. Available at http://www.atsdr.cdc.gov/toxprofiles/tp33.html.
- 4. Bouman, A.A., A.J. Platenkamp, and F.D. Posma. 1986. Determination of cobalt in urine by flameless atomic absorption spectroscopy. Comparison of direct analysis using Zeeman background correction and indirect analysis using extraction in organic solution. *Ann Clin Biochem* 23:346-350.
- 5. Budavari, S., M.J. O'Neil, A. Smith, P.E. Heckelman, and J.F. Kinneary. 1996. The Merck Index. Merck & Co., Inc., Whitehouse Station, NJ. pp. 414.
- 6. CDC. 2001. National Report on Human Exposure to Environmental Chemicals: Cobalt. Centers for Disease Control and Prevention, National Center for Environmental Health, Division of Laboratory Sciences. Available at (http://www.cdc.gov/nceh/dls/report/results/cobalt.htm.).
- 7. ChemFinder. 2001. Cobalt sulfate. CambridgeSoft. Available at (http://chemfinder.cambridgesoft.com and search 10124-43-3).
- 8. Chem Sources. 2001. Chemical Sources International. Available at (http://www.chemsources.com and search 10124-43-3 or 10026-24-1).
- 9. EPA. 1999. Background Report on Fertilizer Use, Contaminants and Regulations. Office of Pollution Prevention and Toxics, U.S. Environmental Protection Agency, Washington, DC. Available at (<u>www.epa.gov/opptintr/fertilizer.pdf</u>).
- Goodman, L.S. and A.G. Gilman. 2001. Goodman & Gilman's The Pharmacological Basis of Therapeutics, Tenth Edition. Hardman, J.G., L.E. Limbird and A.G. Gilman eds. McGraw-Hill, New York.
- 11. Heinrich, R. and J. Angerer. 1984. Determination of cobalt in biological materials by voltammetry and electrothermal atomic absorption spectrometry. *Int J Environ Anal Chem* 16:305-314.
- 12. Hillman, R.S. and C.A. Finch. 1985. Drugs effective in iron-deficiency and other hypochromic anemias. In Goodman & Gilman's The Pharmacological Basis of Therapeutics, Seventh Edition. Gilman, A.G., L.S. Goodman, T.W. Rall and F. Murad, eds. Macmillan Publishing Company, New York. pp. 1308-1322.

- HSDB. 2000. Cobaltous Sulfate. Revised February 8, 2000. Last reviewed August 23, 1989. Hazardous Substances Data Bank, National Library of Medicine, Bethesda, MD. Available at (http://toxnet.nlm.nih.gov/cgi-bin/sis/htm.gen?HSDB and search 10124-43-3 and select Cobaltous Sulfate in search results).
- IARC. 1991. Chlorinated Drinking-Water: Chlorination By-products; Some Other Halogenated Compounds: Cobalt and Cobalt Compounds, Vol. 52. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. World Health Organization, International Agency for Research on Cancer, Lyon, France. 363-472 pp.
- 15. Ichikawa, Y., Y. Kusaka, and S. Goto. 1985. Biological monitoring of cobalt exposure, based on cobalt concentrations in blood and urine. *Int Arch Occup Environ Health* 55:269-276.
- 16. Jensen, A.A. and F. Tüchsen. 1990. Cobalt exposure and cancer risk. *Crit Rev Toxicol* 20:427-437.
- 17. Kazantzis, G. 1981. Role of cobalt, iron, lead, manganese, mercury, platinum, selenium, and titanium in carcinogenesis. *Environ Health Perspect* 40:143-161.
- Kirk, R.E. and D.F. Othmer. 1999. Kirk-Othmer Encyclopedia of Chemical Technology. Herman, F., J.J. McKetta, D.F. Othmer and A. Standen eds. Wiley, New York, pp. 482-484.
- NTP. 1998. Toxicology and Carcinogenesis Studies of Cobalt Sulfate Heptahydrate in F344/N Rats and B6C3F₁ Mice (Inhalation Studies). National Toxicology Program TR No 471. 471 pp.
- 20. Shedd, K.B. 1999. Cobalt. Revised August 30, 2000. U.S. Geological Survey Minerals Yearbook. Available at (http://minerals.usgs.gov/minerals/pubs/commodity/myb/ and select Cobalt).
- 21. Smith, T., C.J. Edmonds, and C.F. Barnaby. 1972. Absorption and retention of cobalt in man by whole-body counting. *Health Phys* 22:359-367.
- Sunderman, F.W., Jr., S.M. Hopfer, T. Swift, W.N. Rezuke, L. Ziebka, P. Highman, B. Edwards, M. Folcik, and H.R. Gossling. 1989. Cobalt, chromium, and nickel concentrations in body fluids of patients with porous-coated knee or hip prostheses. *J Orthop Res* 7:307-315.
- 23. Washington State. 1999. Screening Survey for Metals and Dioxins in Fertilizer Products and Soils in Washington State. Washington State Department of Ecology. Available at (<u>www.ecy.wa.gov/pubs/99309.pdf</u>).
- 24. USGS. 2001. Mineral Industry Surveys. U.S. Department of the Interior. Available at (http://minerals.usgs.gov/minerals/pubs/commodity/cobalt/21000501.pdf).