


Hexachlorobenzene

CAS No. 118-74-1

Reasonably anticipated to be a human carcinogen

First listed in the *Third Annual Report on Carcinogens* (1983)



Carcinogenicity

Hexachlorobenzene is *reasonably anticipated to be a human carcinogen* based on sufficient evidence of carcinogenicity from studies in experimental animals.

Cancer Studies in Experimental Animals

Oral exposure to hexachlorobenzene caused tumors in several rodent species and at two different tissue sites. Dietary administration of hexachlorobenzene caused liver tumors (hepatocellular tumors) in female rats and mice and in hamsters of both sexes. In hamsters of both sexes, it also caused blood-vessel tumors in the liver (hemangioendothelioma) and benign thyroid-gland tumors (follicular-cell adenoma) (IARC 1979, Smith and Cabral 1980).

Since hexachlorobenzene was listed in the *Third Annual Report on Carcinogens*, additional studies in rats have been identified. Dietary exposure caused benign and malignant liver tumors (bile-duct adenoma and hepatocellular carcinoma) and benign blood-vessel tumors in the liver (hemangioma) in females and benign kidney tumors (adenoma) in both sexes. Perinatal exposure to hexachlorobenzene followed by dietary exposure for up to 130 weeks caused benign liver tumors (hepatocellular adenoma) in females, benign parathyroid-gland tumors (adenoma) in males, and benign adrenal-gland tumors (pheochromocytoma) in both sexes (IARC 1987, 2001).

Cancer Studies in Humans

At the time hexachlorobenzene was listed in the *Third Annual Report on Carcinogens*, no epidemiological studies had evaluated the relationship between human cancer and exposure specifically to hexachlorobenzene. Since then, several case-control studies, mostly of breast cancer, have been published. The International Agency for Research on Cancer concluded that there was inadequate evidence in humans for the carcinogenicity of hexachlorobenzene (IARC 2001). No association between exposure to hexachlorobenzene and breast cancer risk was found in five small case-control studies or three larger studies that assessed hexachlorobenzene exposure by measuring it in biological samples obtained close to the time of breast-cancer diagnosis. In a fourth large study, which assessed exposure from banked serum samples collected prior to diagnosis, breast-cancer risk was higher among women with higher serum concentrations of hexachlorobenzene than among women with the lowest serum concentrations, based on sampling close to the time of diagnosis; however, no dose-response relationship was observed. No significant associations between serum hexachlorobenzene concentration and risk of cancer at other tissue sites were found; however, only one study was available for each tissue site.

Since the IARC (2001) review, a number of additional studies have been conducted, mainly of breast cancer and non-Hodgkin lymphoma (NHL). Two studies reported significantly higher serum hexachlorobenzene levels in women with breast cancer than in control

subjects (Charlier *et al.* 2003, 2004), but four other studies found no significant association between serum hexachlorobenzene level and breast cancer (Lopez-Carrillo *et al.* 2002, Pavuk *et al.* 2003, Iwasaki *et al.* 2008, Itoh *et al.* 2009). One study of NHL found a significant dose-related risk associated with serum hexachlorobenzene (Spinelli *et al.* 2007), and two studies found a significantly increased risk among patients with high Epstein-Barr virus antibody titers (also associated with NHL) (Hardell *et al.* 2001, 2009). However, no association with NHL was observed in a study using banked serum samples collected up to 20 years prior to diagnosis and analyzed for hexachlorobenzene (Cantor *et al.* 2003) or in a multicenter study of lymphoma patients using blood levels of hexachlorobenzene measured close to the time of diagnosis (Cocco *et al.* 2008).

Properties

Hexachlorobenzene is a chlorinated aromatic hydrocarbon that exists as a white needle-like crystalline solid at room temperature (HSDB 2010). It is practically insoluble in water, sparingly soluble in cold alcohol and carbon tetrachloride, and soluble in benzene, chloroform, ether, and carbon disulfide. It is stable under normal temperatures and pressures (Akron 2010). It is combustible but it does not ignite readily. When hexachlorobenzene decomposes, it emits highly toxic fumes of hydrochloric acid, other chlorinated compounds, carbon monoxide, and carbon dioxide. Physical and chemical properties of hexachlorobenzene are listed in the following table.

Property	Information
Molecular weight	284.8
Density	2.044 g/cm ³ at 23°C
Melting point	231.8°C
Boiling point	325°C
Log K_{ow}	5.73
Water solubility	4.7×10^{-6} g/L at 25°C
Vapor density relative to air	9.83

Source: HSDB 2010.

Use

No commercial uses of hexachlorobenzene as an end product in the United States were identified (ATSDR 2002). Previously, it was used as a seed-treatment fungicide for onions, sorghum, wheat, and other grains (IARC 1979). All registered pesticide uses in the United States were voluntarily cancelled in 1984 (ATSDR 2002). Hexachlorobenzene was also used as a chemical intermediate in dye manufacturing, in the synthesis of other organic chemicals, and in the production of pyrotechnic compositions for the military. It was used as a raw material for synthetic rubber, as a plasticizer for polyvinyl chloride, as a porosity controller in the manufacture of electrodes, and as a wood preservative (IARC 1979, ATSDR 2002).

Production

Commercial production of hexachlorobenzene in the United States was first reported in 1933 (IARC 1979). In 1975, 3,200 lb of hexachlorobenzene was produced, but it has not been produced commercially in the United States since the late 1970s. In 1972, an estimated 2.5 million to 4.9 million pounds of hexachlorobenzene was produced in the United States as a by-product of production of other chlorinated solvents and pesticides such as tetrachloroethylene, trichloroethylene, carbon tetrachloride, vinyl chloride, atrazine, propazine, simazine, pentachlorophenol, chlorothalonil, and pentachloronitrobenzene. In addition, hexachlorobenzene may be formed during combustion of municipal waste or in waste streams from chlor-alkali and wood-preserving plants (IARC 1979, ATSDR 2002).

In 2002, nine U.S. chemical companies produced hexachlorobenzene for on-site use and processing, as a by-product, or as an impurity (ATSDR 2002). In 2009, hexachlorobenzene was available from 19 suppliers worldwide, including 13 U.S. suppliers (ChemSources 2010). U.S. imports of hexachlorobenzene totaled about 5,400 lb in 1977 and 38,000 lb in 1982 (ATSDR 2002, HSDB 2010). Imports of hexachlorobenzene and dichlorodiphenyltrichloroethane (DDT) (reported together) have generally been low since 1989. However, 2.3 million pounds was imported in 1993 and 4.9 million pounds in 2001, even though neither hexachlorobenzene nor DDT is used in the United States (USITC 2010). Imports were zero in 2007 and 11 lb in both 2006 and 2008. U.S. exports in this category have remained at about 1 million pounds or less since 1989, reaching a low of 7,000 lb in 2008. Reports filed under the U.S. Environmental Protection Agency's Toxic Substances Control Act Inventory Update Rule indicated that U.S. production plus imports of hexachlorobenzene totaled 10,000 to 500,000 lb in 1998 and 2002 (EPA 2004).

Exposure

Hexachlorobenzene is highly persistent in the environment and highly resistant to degradation; therefore, the general population may be exposed at low concentrations (ATSDR 2002). When hexachlorobenzene is released to the environment, it may be taken up by plants and animals and can bioaccumulate through the food chain. Hexachlorobenzene has been detected in terrestrial, freshwater, and marine food chains in the Great Lakes and Arctic regions. Populations with the greatest potential for exposure include those who ingest fish caught from contaminated water bodies or who reside near former manufacturing or waste-disposal sites.

According to EPA's Toxics Release Inventory, environmental releases of hexachlorobenzene ranged from over 1 million pounds in 1989 and 1991 to a low of about 12,600 lb in 1997. In 2008, 49 facilities released a total of 50,636 lb of hexachlorobenzene, mostly to on-site and off-site landfills. The majority of releases came from 5 facilities, and 12 facilities reported releases of more than 100 lb (TRI 2010). When hexachlorobenzene is released to air, it tends to remain mainly in the vapor phase and can therefore be transported great distances (for example, from temperate to polar regions). When released to water, hexachlorobenzene is strongly adsorbed to particles and sediment and is not degraded or hydrolyzed (ATSDR 2002). In the Great Lakes region, hexachlorobenzene was found in drinking and surface water and, at higher levels, in soil and sediment. In 1972, it was detected in agricultural soils where it had been used as a pesticide, at lower levels in urban soils, and at higher levels in soils near uncontrolled hazardous-waste sites. It was found at high concentrations in sediments near industrial sites at Galveston Bay, Texas (ATSDR 2002).

In dietary surveys conducted by the U.S. Food and Drug Administration, the frequency at which hexachlorobenzene was detected in foods declined from 9% in the early 1980s to less than 2% in 1994 (ATSDR 2002). Consequently, the U.S. average daily intake of hexachlorobenzene through foods declined by a factor of 5 over this period. In the FDA Total Diet Study, hexachlorobenzene was detected in 229 of 1,748 samples (13%) of 42 different foods; the highest concentration was found in butter (FDA 2006).

Hexachlorobenzene has been detected in the blood of numerous groups of people, especially indigenous populations of Arctic regions, in the blood and breast milk of pregnant and lactating women, and in the placenta and cord blood. Organochlorine compounds were found in maternal blood in circumpolar populations in Greenland, Canada, Alaska, Norway, Sweden, Iceland, Finland, and Russia (Van Oostdam *et al.* 2004). In Arctic Canada, hexachlorobenzene was detected in all samples of maternal blood, and at higher concentrations in blood

from Inuit women than from Caucasian women in the region. Cord-blood plasma concentrations showed a similar trend (Butler Walker *et al.* 2003). Breast-milk concentrations of hexachlorobenzene were elevated in populations of women who ate contaminated local fish in New York State and Finland (Greizerstein *et al.* 1999, Kostyniak *et al.* 1999, Fitzgerald *et al.* 2001, Damgaard *et al.* 2006). Hexachlorobenzene was found in all blood samples from pregnant women in an agricultural community in California (Fenster *et al.* 2006). The diet of the Inuit population in Greenland was studied to determine the source of the high and increasing concentration of hexachlorobenzene. The blood levels of hexachlorobenzene in Greenland Arctic populations appeared to correlate with consumption of meals containing seal and whale (Deutch *et al.* 2004, 2006). Hexachlorobenzene was detected in all adipose tissue samples collected at autopsy from Greenlanders (Dewailly *et al.* 1999). Hexachlorobenzene was detected in 98% of the blood samples collected from Akwesasne Mohawk youth living along the St. Lawrence River in New York State and Quebec; levels were somewhat higher in youths who had been breastfed as infants (Schell *et al.* 2003). In a study of consumers of sport fish in New York State, the mean blood hexachlorobenzene concentration was not significantly greater than that of nonconsumers of sport fish (Bloom *et al.* 2005).

The National Occupational Exposure Survey (conducted from 1981 to 1983) estimated that 1,038 workers at 10 facilities, including 26 women, potentially were exposed to hexachlorobenzene (NIOSH 1990). The largest numbers of exposed workers were chemical technicians (467 workers) and their supervisors (187 workers). Occupations with the highest potential for exposure included fungicide application, organic-chemical synthesis, synthetic-rubber production, seed disinfection, pesticide production, and wood preservation.

Regulations

Department of Transportation (DOT)

Hexachlorobenzene is considered a hazardous material, and special requirements have been set for marking, labeling, and transporting this material.

Environmental Protection Agency (EPA)

Clean Air Act

National Emission Standards for Hazardous Air Pollutants: Listed as a hazardous air pollutant.

New Source Performance Standards: Manufacture of hexachlorobenzene is subject to certain provisions for the control of volatile organic compound emissions.

Urban Air Toxics Strategy: Identified as one of 33 hazardous air pollutants that present the greatest threat to public health in urban areas.

Clean Water Act

Effluent Guidelines: Chlorinated benzenes are listed as a toxic pollutant.

Water Quality Criteria: Based on fish or shellfish and water consumption = 0.000079 µg/L; based on fish or shellfish consumption only = 0.000079 µg/L.

Comprehensive Environmental Response, Compensation, and Liability Act

Reportable quantity (RQ) = 10 lb.

Emergency Planning and Community Right-To-Know Act

Toxics Release Inventory: Listed substance subject to reporting requirements.

Resource Conservation and Recovery Act

Characteristic Hazardous Waste: Toxic characteristic leaching procedure threshold = 0.13 mg/L.

Listed Hazardous Waste: Waste codes for which the listing is based wholly or partly on the presence of hexachlorobenzene = U127, F024, F025, K016, K018, K030, K042, K085, K149, K150, K151.

Listed as a hazardous constituent of waste.

Safe Drinking Water Act

Maximum contaminant level (MCL) = 0.001 mg/L.

Food and Drug Administration (FDA)

Maximum permissible level in bottled water = 0.001 mg/L.

Guidelines

American Conference of Governmental Industrial Hygienists (ACGIH)

Threshold limit value – time-weighted average (TLV-TWA) = 0.002 mg/m³.
Potential for dermal absorption.

References

- Akron. 2010. *The Chemical Database*. The Department of Chemistry at the University of Akron. <http://ull.chemistry.uakron.edu/erd> and search on CAS number. Last accessed: 1/8/10.
- ATSDR. 2002. *Toxicological Profile for Hexachlorobenzene*. Agency for Toxic Substances and Disease Registry. <http://www.atsdr.cdc.gov/ToxProfiles/tp90.pdf>.
- Bloom MS, Vena JE, Swanson MK, Moysich KB, Olson JR. 2005. Profiles of ortho-polychlorinated biphenyl congeners, dichlorodiphenyldichloroethylene, hexachlorobenzene, and Mirex among male Lake Ontario sportfish consumers: the New York State angler cohort study. *Environ Res* 97(2): 178-194.
- Butler Walker J, Seddon L, McMullen E, Houseman J, Tofflemire K, Corriveau A, et al. 2003. Organochlorine levels in maternal and umbilical cord blood plasma in Arctic Canada. *Sci Total Environ* 302(1-3): 27-52.
- Cantor KP, Strickland PT, Brock JW, Bush D, Helzlsouer K, Needham LL, Zahm SH, Comstock GW, Rothman N. 2003. Risk of non-Hodgkin's lymphoma and prediagnostic serum organochlorines: β -hexachlorocyclohexane, chlordane/heptachlor-related compounds, dieldrin, and hexachlorobenzene. *Environ Health Perspect* 111(2): 179-183.
- Charlier C, Albert A, Herman P, Hamoir E, Gaspard U, Meurisse M, Plomteux G. 2003. Breast cancer and serum organochlorine residues. *Occup Environ Med* 60(5): 348-351.
- Charlier C, Foidart JM, Pitance F, Herman P, Gaspard U, Meurisse M, Plomteux G. 2004. Environmental dichlorodiphenyltrichloroethane or hexachlorobenzene exposure and breast cancer: Is there a risk? *Clin Chem Lab Med* 42(2): 222-227.
- ChemSources. 2010. *Chem Sources - Chemical Search*. Chemical Sources International, Inc. <http://www.chemsources.com/chemonline.html> and search on hexachlorobenzene. Last accessed: 1/8/10.
- Cocco P, Brennan P, Ibba A, de Sanjose Llongueras S, Maynadie M, Nieters A, et al. 2008. Plasma polychlorobiphenyl and organochlorine pesticide level and risk of major lymphoma subtypes. *Occup Environ Med* 65(2): 132-140.
- Damgaard IN, Skakkebaek NE, Toppari J, Virtanen HE, Shen H, Schramm KW, et al. 2006. Persistent pesticides in human breast milk and cryptorchidism. *Environ Health Perspect* 114(7): 1133-1138.
- Deutch B, Pedersen HS, Hansen JC. 2004. Dietary composition in Greenland 2000, plasma fatty acids and persistent organic pollutants. *Sci Total Environ* 331(1-3): 177-188.
- Deutch B, Dyerberg J, Pedersen HS, Asmund G, Moller P, Hansen JC. 2006. Dietary composition and contaminants in north Greenland, in the 1970s and 2004. *Sci Total Environ* 370(2-3): 372-381.
- Dewailly E, Mulvad G, Pedersen HS, Ayotte P, Demers A, Weber JP, Hansen JC. 1999. Concentration of organochlorines in human brain, liver, and adipose tissue autopsy samples from Greenland. *Environ Health Perspect* 107(10): 823-828.
- EPA. 2004. *Non-confidential IUR Production Volume Information*. U.S. Environmental Protection Agency. <http://www.epa.gov/oppt/iur/tools/data/2002-vol.html> and search on CAS number.
- FDA. 2006. *Total Diet Study Market Baskets 1991-3 through 2003-4*. U.S. Food and Drug Administration. <http://www.fda.gov/downloads/Food/FoodSafety/FoodContaminantsAdulteration/TotalDietStudy/UCM184304.pdf>. <http://www.cfsan.fda.gov/~acrobot/tds1byps.pdf>
- Fenster L, Eskenazi B, Anderson M, Bradman A, Harley K, Hernandez H, Hubbard A, Barr DB. 2006. Association of *in utero* organochlorine pesticide exposure and fetal growth and length of gestation in an agricultural population. *Environ Health Perspect* 114(4): 597-602.
- Fitzgerald EF, Hwang SA, Deres DA, Bush B, Cook K, Worswick P. 2001. The association between local fish consumption and DDE, mirex, and HCB concentrations in the breast milk of Mohawk women at Akwesasne. *J Expo Anal Environ Epidemiol* 11(5): 381-388.
- Greizerstein HB, Stinson C, Mendola P, Buck GM, Kostyniak PJ, Vena JE. 1999. Comparison of PCB congeners and pesticide levels between serum and milk from lactating women. *Environ Res* 80(3): 280-286.
- Hardell E, Eriksson M, Lindstrom G, Van Bavel B, Linde A, Carlberg M, Liljegren G. 2001. Case-control study on concentrations of organohalogen compounds and titers of antibodies to Epstein-Barr virus antigens in the etiology of non-Hodgkin lymphoma. *Leuk Lymphoma* 42(4): 619-629.
- Hardell K, Carlberg M, Hardell L, Bjornfoth H, Ericson Jogsten I, Eriksson M, Van Bavel B, Lindstrom G. 2009. Concentrations of organohalogen compounds and titres of antibodies to Epstein-Barr virus antigens and the risk for non-Hodgkin lymphoma. *Oncol Rep* 21(6): 1567-1576.
- HSDB. 2010. *Hazardous Substances Data Bank*. National Library of Medicine. <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB> and search on CAS number. Last accessed: 1/8/10.
- IARC. 1979. Hexachlorobenzene. In *Some Halogenated Hydrocarbons*. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 20. Lyon, France: International Agency for Research on Cancer. pp. 155-178.
- IARC. 1987. Hexachlorobenzene. In *Overall Evaluations of Carcinogenicity*. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, suppl. 7. Lyon, France: International Agency for Research on Cancer. pp. 219-220.
- IARC. 2001. Hexachlorobenzene. In *Some Thyrotropic Agents*. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 79. Lyon, France: International Agency for Research on Cancer. pp. 493-568.
- Itoh H, Iwasaki M, Hanaoka T, Kasuga Y, Yokoyama S, Onuma H, Nishimura H, Kusama R, Tsugane S. 2009. Serum organochlorines and breast cancer risk in Japanese women: a case-control study. *Cancer Causes Control* 20(5): 567-580.
- Iwasaki M, Inoue M, Sasazuki S, Kurahashi N, Itoh H, Usuda M, Tsugane S. 2008. Plasma organochlorine levels and subsequent risk of breast cancer among Japanese women: a nested case-control study. *Sci Total Environ* 402(2-3): 176-183.
- Kostyniak PJ, Stinson C, Greizerstein HB, Vena J, Buck G, Mendola P. 1999. Relation of Lake Ontario fish consumption, lifetime lactation, and parity to breast milk polychlorobiphenyl and pesticide concentrations. *Environ Res* 80(2 Pt 2): S166-S174.
- Lopez-Carrillo L, Lopez-Cervantes M, Torres-Sanchez L, Blair A, Cebrían ME, García RM. 2002. Serum levels of beta-hexachlorocyclohexane, hexachlorobenzene and polychlorinated biphenyls and breast cancer in Mexican women. *Eur J Cancer Prev* 11(2): 129-135.
- NIOSH. 1990. *National Occupational Exposure Survey (1981-83)*. National Institute for Occupational Safety and Health. Last updated: 7/1/90. <http://www.cdc.gov/noes/noes1/a1753sic.html>.
- Pavuk M, Cerhan JR, Lynch CF, Kocan A, Petrik J, Chovancova J. 2003. Case-control study of PCBs, other organochlorines and breast cancer in Eastern Slovakia. *J Expo Anal Environ Epidemiol* 13(4): 267-275.
- Schell LM, Hubicki LA, DeCaprio AP, Gallo MV, Ravenscroft J, Tarbell A, Jacobs A, David D, Worswick P. 2003. Organochlorines, lead, and mercury in Akwesasne Mohawk youth. *Environ Health Perspect* 111(7): 954-961.
- Smith AG, Cabral JR. 1980. Liver-cell tumours in rats fed hexachlorobenzene. *Cancer Lett* 11(2): 169-172.
- Spinelli JJ, Ng CH, Weber JP, Connors JM, Gascoyne RD, Lai AS, et al. 2007. Organochlorines and risk of non-Hodgkin lymphoma. *Int J Cancer* 121(12): 2767-2775.
- TRI. 2009. *TRI Explorer Chemical Report*. U.S. Environmental Protection Agency. Last updated: 3/19/09. <http://www.epa.gov/triexplorer> and select Hexachlorobenzene.
- USITC. 2010. *USITC Interactive Tariff and Trade DataWeb*. United States International Trade Commission. http://dataweb.usitc.gov/scripts/user_set.asp and search on HTS no. 290362. Last accessed: 1/8/10.
- Van Oostdam JC, Dewailly E, Gilman A, Hansen JC, Odland JO, Chashchin V, et al. 2004. Circumpolar maternal blood contaminant survey, 1994-1997 organochlorine compounds. *Sci Total Environ* 330(1-3): 55-70.