


Tetrachloroethylene

CAS No. 127-18-4

Reasonably anticipated to be a human carcinogen

First listed in the *Fifth Annual Report on Carcinogens* (1989)

Also known as perchloroethylene or perc



Carcinogenicity

Tetrachloroethylene is *reasonably anticipated to be a human carcinogen* based on sufficient evidence of carcinogenicity from studies in experimental animals.

Cancer Studies in Experimental Animals

Tetrachloroethylene caused tumors in two rodent species, at several different tissue sites, and by two different routes of exposure. Inhalation exposure to tetrachloroethylene caused benign and malignant liver tumors (hepatocellular adenoma and carcinoma) in mice of both sexes and mononuclear-cell leukemia in rats of both sexes. In male rats, it also increased the combined incidence of benign and malignant tubular-cell kidney tumors, which are rare in rats (NTP 1986). Liver tumors were also observed in mice of both sexes administered tetrachloroethylene by stomach tube (NTP 1977, IARC 1979, 1987).

Cancer Studies in Humans

The data available from epidemiological studies were inadequate to evaluate the relationship between human cancer and exposure specifically to tetrachloroethylene at the time it was listed in the *Fifth Annual Report on Carcinogens*. A number of cohort and case-control studies of occupational exposure to tetrachloroethylene had been conducted. Although tetrachloroethylene may have been the predominant solvent to which workers were exposed, co-exposure to other chemicals (in particular trichloroethylene) among workers in the drycleaning industry was common (IARC 1982, 1987).

Since tetrachloroethylene was listed *Fifth Annual Report on Carcinogens*, additional epidemiological studies have been identified. Several ecological studies assessed cancer outcomes among residents exposed to groundwater contaminated with tetrachloroethylene, among other chemicals. The International Agency for Research on Cancer (IARC 1995) concluded that there was limited evidence for the carcinogenicity of tetrachloroethylene in humans, based mainly on evidence of consistent associations between tetrachloroethylene exposure and esophageal cancer, cervical cancer, and non-Hodgkin lymphoma; however, confounding by exposure to other chemicals could not be ruled out, and the total numbers in the combined cohort studies were small.

Properties

Tetrachloroethylene is a halogenated alkene that exists at room temperature as a colorless liquid with a mildly sweet, ethereal odor (NTP 1986, HSDB 2009). It is only slightly soluble in water but is miscible with alcohol, ether, chloroform, benzene, and solvent hexane and soluble in most fixed and volatile oils. Tetrachloroethylene can be oxidized in air and sunlight and reacts with chemically active metals (e.g., barium or lithium) (IARC 1995). Physical and chemical properties of tetrachloroethylene are listed in the following table.

Property	Information
Molecular weight	165.8
Specific gravity	1.6227 at 20°C/4°C
Melting point	-22.3°C
Boiling point	121.3°C
Log K_{ow}	3.4
Water solubility	206 mg/L at 25°C
Vapor pressure	18.5 mm Hg at 25°C
Vapor density relative to air	5.7

Source: HSDB 2009.

Use

Tetrachloroethylene is used primarily as a cleaning solvent and as a chemical precursor for fluorocarbons. In the 1970s, domestic use patterns were as follows: 58% for drycleaning and textile processing, 18% for metal cleaning, 12% for chemical intermediates, and 12% for all other uses (IARC 1995). During the 1990s, tetrachloroethylene use in the drycleaning industry declined in order to meet stringent government regulations for workplace exposure. By 2002, uses were 15% for drycleaning, 10% for metal cleaning, 65% for chemical intermediates, and 10% for other uses (CMR 2002). Tetrachloroethylene also has been used as an insulating fluid and cooling gas in electrical transformers; in paint removers, printing inks, adhesive formulations, paper coatings, and leather treatments; in aerosol formulations, such as water repellents, automotive cleaners, silicone lubricants, and spot removers; as an extractant for pharmaceuticals; to remove soot from industrial boilers; and as an antihelminthic agent (IARC 1995).

Production

Tetrachloroethylene was first prepared in 1821, and commercial production in the United States began in 1925. Several commercial grades are available that differ in the amount and type of added stabilizers (e.g., amines, phenols, and epoxides). Annual production rose rapidly in the United States from 5,000 metric tons (1.1 million pounds) in 1941 to a peak of 347,000 metric tons (763 million pounds) in 1980. From 1980 to 1993, annual production declined by more than 60% (IARC 1995, ATSDR 1997), but from 1996 to 1999, U.S. demand (domestic production plus imports) increased from 280 million pounds to 318 million pounds. In 2002, the combined production capacity of the three U.S. manufacturers of tetrachloroethylene was 430 million pounds (CMR 2002). In 2009, tetrachloroethylene was produced by 15 manufacturers worldwide, including 3 U.S. manufacturers (SRI 2009), and was available from 115 suppliers, including 43 U.S. suppliers (ChemSources 2009).

U.S. imports of tetrachloroethylene were 132 million pounds in 1977, 38 million pounds in 1982, and 140 million pounds in 1985 (HSDB 2009). Imports averaged 80 million pounds from 1992 to 1996 and 36 million pounds from 1998 to 2000, reaching a maximum of 72.4 million pounds in 2003 (USITC 2009). In 2008, imports were 36.2 million pounds. U.S. exports of tetrachloroethylene totaled 64 million pounds in 1978, 54 million pounds in 1983, and 22 million pounds in 1985. Exports averaged 45 million pounds from 1992 to 1996, increasing to an average of 50 million pounds from 1998 to 2008 (HSDB 2009).

Exposure

Tetrachloroethylene is widely distributed in the environment, because it is released from many industrial processes and consumer products. The primary routes of potential human exposure to tetrachloroethylene are inhalation and ingestion of contaminated water or food. Dermal exposure also may occur, but is not important for the majority of the population (ATSDR 1997). According to the U.S. Environmental Protection Agency's Toxics Release Inventory, total en-

Environmental releases of tetrachloroethylene declined by almost 94% from 37.7 million pounds in 1988 to 2.2 million pounds in 2008. In 2008, most of the releases, from 225 facilities, were to air (TRI 2010). Numerous studies have detected tetrachloroethylene in the air in the United States in rural, urban, and industrial areas. Typical concentrations in rural and remote areas were in the low parts-per-trillion range, while concentrations in urban and industrial areas were in the high parts-per-trillion to low parts-per-billion range.

Studies conducted in New York City measured ambient indoor-air concentrations of tetrachloroethylene in apartments and in a day-care facility located in buildings containing drycleaning facilities (Schreiber *et al.* 2002). Concentrations of tetrachloroethylene were much higher in the buildings with drycleaning facilities than in buildings without such facilities. Tetrachloroethylene levels were measured in exhaled breath by personal monitoring devices and in samples of blood from individuals living in apartments over drycleaning facilities. Concentrations of tetrachloroethylene were elevated in samples from exposed individuals. The general population may also be exposed to tetrachloroethylene through use of coin-operated laundromats that contain drycleaning machines or through exposure to freshly drycleaned clothing. Studies show elevated concentrations of tetrachloroethylene in laundromats (even months after removal of the drycleaning machines). Tetrachloroethylene concentrations in homes with freshly drycleaned clothing stored in the closets may be 2 to 30 times higher than average background levels. In addition, workers in the drycleaning industry may carry tetrachloroethylene home on their person or clothes and therefore serve as a source of exposure of their families. In one study, indoor air concentrations of tetrachloroethylene in apartments where drycleaning workers lived were over 10 times the levels in other apartments in the same buildings, where the occupants were not employed by drycleaning facilities (ATSDR 1997).

Tetrachloroethylene may also be formed in small quantities during chlorination of water. EPA estimated that in 1985, 11.4 million people were exposed to tetrachloroethylene at concentrations of at least 0.5 µg/L and 874,000 were exposed to concentrations of at least 5 µg/L from municipal water supplies in the United States (IARC 1995). Contamination of drinking water with tetrachloroethylene was reported in the Cape Cod, Massachusetts, area in the late 1970s (Webler and Brown 1993). The chemical leached from the vinyl lining of asbestos-cement water distribution pipes. The highest level reported was 18 mg/L from a pipe in Falmouth, Massachusetts, but levels of 1,600 to 7,750 µg/L were reported for pipes running along dead-end streets (Wakeham *et al.* 1980, Aschengrau *et al.* 2003). Tetrachloroethylene has also been detected in rainwater, sea water, rivers, groundwater, commercial deionized charcoal-filtered water, dairy products, meats, oils and fats, beverages, fruits and vegetables, fresh bread, fish, shellfish, marine mammals, glues, printing inks, lubricants, stain and paint removers, and other consumer products (IARC 1995, ATSDR 1997). It has even been detected in snow in Antarctica (Zoccolillo *et al.* 2007). It was detected in 67 items in the U.S. Food and Drug Administration's Total Diet Study (FDA 2006).

Workers involved in drycleaning, metal degreasing, and fluorocarbon production are likely to be exposed to tetrachloroethylene. The National Occupational Exposure Survey (conducted from 1981 to 1983) estimated that 688,000 workers in 49,025 U.S. facilities potentially were exposed to tetrachloroethylene (NIOSH 1990). A 1994 survey prepared by industry estimated that 450,000 workers potentially were exposed (IARC 1995). Occupational exposure has trended lower over the past several decades. Typical tetrachloroethylene concentrations in workplace air at drycleaning facilities were 350 to 700 mg/m³ (about 50 to 100 ppm) in the 1970s and 70 to 350 mg/m³ (about 10 to

50 ppm) in the 1980s (IARC 1995). The highest exposures occur during loading and unloading of the drycleaning machines. More recent studies by the National Institute for Occupational Safety and Health indicated that exposure levels in the drycleaning industry were below the recommended occupational exposure guideline of 25 ppm (ATSDR 1997). In 2003, the mean concentration of tetrachloroethylene at U.S. drycleaning facilities was 3.8 ppm (Toraason *et al.* 2003).

Regulations

Coast Guard, Department of Homeland Security

Minimum requirements have been established for safe transport of tetrachloroethylene on ships and barges.

Consumer Product Safety Commission (CPSC)

Visual novelty devices containing tetrachloroethylene have labeling requirements.

Department of Transportation (DOT)

Tetrachloroethylene is considered a hazardous material and a marine pollutant, and special requirements have been set for marking, labeling, and transporting this material, including transporting it in tank cars.

Environmental Protection Agency (EPA)

Clean Air Act

National Emission Standards for Hazardous Air Pollutants: Listed as a hazardous air pollutant.

New Source Performance Standards: Manufacture of tetrachloroethylene is subject to certain provisions for the control of volatile organic compound emissions.

Urban Air Toxics Strategy: Identified as one of 33 hazardous air pollutants that present the greatest threat to public health in urban areas.

Clean Water Act

Effluent guidelines: Listed as a toxic pollutant.

Water quality criteria: Based on fish or shellfish and water consumption = 10 µg/L; based on fish or shellfish consumption only = 29 µg/L.

Comprehensive Environmental Response, Compensation, and Liability Act

Reportable quantity (RQ) = 100 lb.

Emergency Planning and Community Right-To-Know Act

Toxics Release Inventory: Listed substance subject to reporting requirements.

Resource Conservation and Recovery Act

Characteristic Hazardous Waste: Toxicity characteristic leaching procedure (TCLP) threshold = 0.7 mg/L.

Listed Hazardous Waste: Waste codes for which the listing is based wholly or partly on the presence of tetrachloroethylene = U210, F001, F002, F024, F025, K016, K019, K020, K073, K116, K150, K151. Listed as a hazardous constituent of waste.

Safe Drinking Water Act

Maximum contaminant level (MCL) = 0.005 mg/L.

Food and Drug Administration (FDA)

Maximum permissible level in bottled water = 0.005 mg/L.

Occupational Safety and Health Administration (OSHA)

While this section accurately identifies OSHA's legally enforceable PELs for this substance in 2010, specific PELs may not reflect the more current studies and may not adequately protect workers.

Permissible exposure limit (PEL) = 100 ppm.

Ceiling concentration = 200 ppm (5 min in any 3 h).

Acceptable peak exposure = 300 ppm.

Guidelines

American Conference of Governmental Industrial Hygienists (ACGIH)

Threshold limit value – time-weighted average (TLV-TWA) = 25 ppm.

Threshold limit value – short-term exposure limit (TLV-STEL) = 100 ppm.

Biological Exposure Index (BEI) (prior to shift) = 3 ppm in end-exhaled air; = 0.5 mg/L in blood.

National Institute for Occupational Safety and Health (NIOSH)

Recommends that workplace exposure levels of substance be minimized.

Immediately dangerous to life and health (IDLH) limit = 150 ppm.

Listed as a potential occupational carcinogen.

References

Aschengrau A, Rogers S, Ozonoff D. 2003. Perchloroethylene-contaminated drinking water and the risk of breast cancer: additional results from Cape Cod, Massachusetts, USA. *Environ Health Perspect* 111(2): 167-173.

ATSDR. 1997. *Toxicological Profile for Tetrachloroethylene*. Agency for Toxic Substances and Disease Registry. <http://www.atsdr.cdc.gov/toxprofiles/tp18.pdf>.

Report on Carcinogens, Fourteenth Edition

- ChemSources. 2009. *Chem Sources - Chemical Search*. Chemical Sources International. <http://www.chemsources.com/chemonline.html> and search on tetrachloroethylene. Last accessed: 8/18/09.
- CMR. 2002. Chemical profile – perchloroethylene. *Chem Mark Rep* 8/5/02. ICIS. <http://www.icis.com/Articles/2005/12/02/178388/chemical-profile-perchloroethylene.html>.
- FDA. 2006. *Total Diet Study Market Baskets 1991-3 through 2003-4*. U.S. Food and Drug Administration. <http://www.fda.gov/downloads/Food/FoodSafety/FoodContaminantsAdulteration/TotalDietStudy/UCM184304.pdf>.
- HSDB. 2009. *Hazardous Substances Data Bank*. National Library of Medicine. <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB> and search on CAS number. Last accessed: 8/18/09.
- IARC. 1979. Tetrachloroethylene. In *Some Halogenated Hydrocarbons*. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 20. Lyon, France: International Agency for Research on Cancer. pp. 491-514.
- IARC. 1982. Tetrachloroethylene. In *Chemicals, Industrial Processes and Industries Associated with Cancer in Humans*. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, suppl. 4. Lyon, France: International Agency for Research on Cancer. pp. 243-245.
- IARC. 1987. Tetrachloroethylene. In *Overall Evaluations of Carcinogenicity*. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, suppl. 7. Lyon, France: International Agency for Research on Cancer. pp. 355-357.
- IARC. 1995. Tetrachloroethylene. In *Dry Cleaning, Some Chlorinated Solvents and Other Industrial Chemicals*. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 63. Lyon, France: International Agency for Research on Cancer. pp. 159-221.
- NIOSH. 1990. *National Occupational Exposure Survey (1981-83)*. National Institute for Occupational Safety and Health. Last updated: 7/1/90. <http://www.cdc.gov/noes/noes1/54790sic.html>.
- NTP. 1977. *Bioassay of Tetrachloroethylene for Possible Carcinogenicity*. Technical Report Series no. 13. DHEW Publication No. (NIH) 77-813. Bethesda, MD: National Institutes of Health. 99 pp.
- NTP. 1986. *Toxicology and Carcinogenesis Studies of Tetrachloroethylene (Perchloroethylene) (CAS No. 127-18-4) in F344/N Rats and B6C3F₁ Mice (Inhalation Studies)*. Technical Report Series no. 311. NIH Publication No. 86-2587. Research Triangle Park, NC: National Toxicology Program. 199 pp.
- Schreiber JS, Hudnell HK, Geller AM, House DE, Aldous KM, et al. 2002. Apartment residents' and day care workers' exposures to tetrachloroethylene and deficits in visual contrast sensitivity. *Environ Health Perspect* 110(7): 655-664.
- SRI. 2009. *Directory of Chemical Producers*. Menlo Park, CA: SRI Consulting. Database edition. Last accessed: 8/18/09.
- Toraason M, Butler MA, Ruder A, Forrester C, Taylor L, Ashley DL, et al. 2003. Effect of perchloroethylene, smoking, and race on oxidative DNA damage in female dry cleaners. *Mutat Res* 539(1-2): 9-18.
- TRI. 2010. *TRI Explorer Chemical Report*. U.S. Environmental Protection Agency. <http://www.epa.gov/triexplorer> and select Tetrachloroethylene. Last accessed: 2/10.
- USITC. 2009. *USITC Interactive Tariff and Trade DataWeb*. United States International Trade Commission. http://dataweb.usitc.gov/scripts/user_set.asp and search on HTS no. 290323. Last accessed: 8/18/09.
- Wakeham SG, Davis AC, Witt RT, Tripp BW, Frew NM. 1980. Tetrachloroethylene contamination of drinking water by vinyl-coated asbestos-cement pipe. *Bull Environ Contam Toxicol* 25(4): 639-645.
- Webler T, Brown HS. 1993. Exposure to tetrachloroethylene via contaminated drinking water pipes in Massachusetts: a predictive model. *Arch Environ Health* 48(5): 293-297.
- Zoccolillo L, Amendola L, Cafaro C, Insogna S. 2007. Volatile chlorinated hydrocarbons in Antarctic superficial snow sampled during Italian ITASE expeditions. *Chemosphere* 67(10): 1897-1903.