

Received by email on October 11, 2016

From: Adria Huseh, National Pork Board



National Pork Board

1776 NW 114th Street, Clive, Iowa 50325

pork.org | Toll-free: 800-456-7675 | Phone: 515-223-2600 | Fax: 515-223-2646

October 11, 2016

Re: Volume 81, No. 175 – Request for information on consumption of red meat, processed meat and meat cooked at high temperatures:

Department of Health and Human Services: National Institutes of Health: Nominations to the National Toxicology Program for the Report on Carcinogens and Office of Health Assessment and Translation; Request for Information:

The National Pork Board appreciates the opportunity to provide evidence regarding the production and consumption of red and processed meat in the United States.

There are limited sources of data for red and processed meat exposure available to the National Toxicology Program's (NTP) request for information and all vary with regard to timeliness, accuracy and specificity. The most accessible sources of red and processed meat data available to the NTP's request are the Food Balance Sheets provided by FAO.¹ These data are gross measures of food availability and thus overestimate intake and provide little specificity with regard to consumption of further processed products. A recent publication comparing intake estimations using FAO data vs. Global Dietary Database (GDD), found world-wide estimates of red and processed meat intake to be 2.2 times higher with FAO as compared to the GDD.² The GDD appears to provide more precise intake data than the FAO, but the database is not publically available (i.e. limited to research group members), almost half of the dietary surveys used were collected prior to the year 2000, and food groups are broad and not clearly defined, thus limiting data specificity for red and processed meat. "Meat" is a broad food category that is ambiguous in nutritional epidemiology. Observational studies commonly report intake of "red meat" or "red and processed meat" without further definition and there is often no intake data to distinguish fresh meat from processed meat or lean meat from higher fat sources. "Processed meat" often reflects a combination of red meat and poultry-based products, but these are rarely reported separately in observational studies.³ For example, "processed meat" intake was recently estimated using the GDD, but the authors do not specify if this is red processed meat only or any processed meat.⁴ Reliance on GDD data that may be 15 or more years old, also limits the relevancy of this data to today's leaner beef and pork supply and changes in the formulation of processed meat to include less sodium and chemical preservatives.

¹ http://faostat3.fao.org/browse/FB/*/E

² Del Gobbo et al. 2015. Assessing global dietary habits: a comparison of national estimates from the FAO and the Global Dietary Database. *Am J Clin Nutr* doi:10.3945/ajcn.114.087403

³ McNeill SM and Van Elswyk, ME. Meat: Role in the Diet. *Encyclopedia of Food and Health, in press.*

⁴ Imamura et al., 2015. Dietary quality among men and women in 187 countries in 1990 and 2010: a systematic assessment. *Lancet Glob Health*, Mar;3(3):e132-42. doi: 10.1016/S2214-109X(14)70381-X.

Vision: Leading a world-class food industry. Responsible. Sustainable. Professional. Profitable.

Mission: The National Pork Board harnesses the resources of all producers to capture opportunity, address challenges and satisfy customers.



National Pork Board

1776 NW 114th Street, Clive, Iowa 50325

pork.org | Toll-free: 800-456-7675 | Phone: 515-223-2600 | Fax: 515-223-2646

As a member of the International Meat Secretariat (IMS), I am able to provide the most accurate production and intake estimations for red and processed meat for the United States. I've included an annotated bibliography of publically available publications and reports of red and processed meat intake. This information is the result of the U.S. food consumption survey data; National Health and Nutrition Examination Survey. For each food reported in NHANES 2003-2006, the USDA Food and Nutrient Database for Dietary Studies (FNDD) databases provided information on the amount of energy and approximately 60 nutrients or food constituents per 100 g of each food. The NHANES 2003-2006 respondents, 16,783 individuals, reported consumption of approximately 5,000 specific foods; each food was identified by USDA by a unique 8-digit food code. Assessed dietary intake by U.S. population age 2 years and older and the following sub-populations: children 2-5, males and females 6-11, 12-18, 19-39 and 40+ were derived from two, 24-hour dietary recalls for each subject. Pork-based foods were classified into two categories: processed pork and fresh pork.

Generally speaking, our findings can be summarized as follows:

- The average intake of total pork in the United States per capita is 29.2 g per day
- The average intake of fresh pork per capita is 7.5 g per day
- The average intake of lean pork per capita is 10.7 g per day
 - Per FDA's definition of lean, according to reference *21 CFR 101.62(e)(1)-(3)*, on seafood or game meat products that contain less than 10g total fat, 4.5g or less saturated fat, and less than 95mg cholesterol per RACC and per 100g
- The average intake of processed pork per capita is 21.7 g per day

There are studies linked to cooking meat at high temperatures which can produce heterocyclic amines (HCAs) and polycyclic aromatic hydrocarbons (PAHs). HCAs and PAHs exposure has been studied in animal models. However, the doses of HCAs and PAHs used in these studies were very high; equivalent to **thousands of times the doses that a person would consume** in a normal diet. Human studies have not established a definitive link between HCA and PAH exposure from cooked meats and cancer in humans. No specific guidelines for HCA/PAH consumption exist. The National Pork Board recommends not overcooking pork to achieve a tender and flavorful product. For pork cuts such as roasts, chops and tenderloin, cook to an internal temperature of 145 degrees F., plus a 3-minute rest. There are several ways to reduce HCAs and PAHs when grilling or pan frying meat that include; trim fat to reduce drips and flavor meats with marinades and rubs. Scientists believe the antioxidants in marinades block HCAs from forming. Herbs including basil, mint, sage and oregano may have similar effects. In addition, avoiding direct exposure of meat to an open flame or a hot metal surface, avoiding prolonged cooking times at high temperatures, continuously turning meat over on a high heat source and removing charred portions of meat can also reduce HCA and PAH exposure.

Vision: Leading a world-class food industry. Responsible. Sustainable. Professional. Profitable.

Mission: The National Pork Board harnesses the resources of all producers to capture opportunity, address challenges and satisfy customers.



National Pork Board

1776 NW 114th Street, Clive, Iowa 50325

pork.org | Toll-free: 800-456-7675 | Phone: 515-223-2600 | Fax: 515-223-2646

Research continues to show that not smoking, responsible alcohol consumption, maintaining a healthy weight and regular physical activity are much more important to cancer risk than eating any individual food. It is also important for individuals to adhere to screening recommendations, recommended clinical examinations and follow-up appointments with their medical professionals.

Again, I appreciate the opportunity to provide this evidence. Also, I would like to offer myself as a resource to the NTP group as they continue to deliberate their request for information on red and processed meat.

Adria Huseth, RD, LD, CPT

Manager of Nutrition Communications & Research

National Pork Board

ahuseth@pork.org

515-223-2632

Publication Citation

Murphy, M., Spungen, J., Bi, X., & Barraj, L. "Fresh and fresh lean pork are substantial sources of key nutrients when these products are consumed by adults in the United States." Nutrition Research/Volume 31.Issue 10 (2011): 776-783.

KEY FINDINGS FROM PUBLICATION:

Selection of fresh, lean pork cuts can help fulfill nutrient needs provided by the protein foods group while also helping to limit energy intake.

- Energy intakes by consumers of fresh lean pork were comparable to intakes by non-consumers of fresh lean pork who are presumably selecting alternate foods from the protein group.
- Fresh lean pork accounts for 23% of total protein intake and 25% or more of total intakes of selenium and thiamin while contributing only 7% of total energy intake and 9% of total fat intake on a day of consumption.
- Fresh lean pork accounts for 10% or more of total intakes of phosphorous, potassium, zinc and B vitamins including riboflavin, niacin, vitamin B6 and vitamin B12 on a day of consumption.

Americans are consuming fresh and fresh lean pork at levels within the total amount that is recommended in the protein foods group by current dietary guidance.

- On a day of consumption, US adults eat 3.2 ounces of fresh pork or 2.5 ounces of fresh lean pork, which are levels within the 5.5 ounce equivalents a day of total protein foods recommended for adults consuming 2,000 calories per day.

Vision: Leading a world-class food industry. Responsible. Sustainable. Professional. Profitable.

Mission: The National Pork Board harnesses the resources of all producers to capture opportunity, address challenges and satisfy customers.



National Pork Board

1776 NW 114th Street, Clive, Iowa 50325

pork.org | Toll-free: 800-456-7675 | Phone: 515-223-2600 | Fax: 515-223-2646

Energy-adjusted intakes of dietary fat by fresh and fresh lean pork consumers on a day of consumption are not different than those of non-consumers of these pork products.

- On a day of consumption, fresh pork accounted for 16% of total fat intake among consumers of fresh pork, while fresh lean pork accounted for 9% of total fat intake among consumers of fresh lean pork.
- Results of this study show no differences in calorie-adjusted intakes of total fat or saturated fat intake between fresh or fresh lean pork consumers and non-consumers of these products.

Selection of fresh, lean pork products by adults not currently eating these products could positively increase dietary variety without adversely affecting nutrient intake.

- Results of this study show that diets including fresh or fresh lean pork provide higher calorie-adjusted amounts of important nutrients like protein, selenium, thiamin and vitamin B6 compared to diets of adults not including these products, while still providing comparable energy-adjusted amounts of total and saturated fat. Dietary guidance recommends selection of lean or low-fat meats; fresh lean pork is consistent with this recommendation and therefore could help increase dietary variety.

Vision: Leading a world-class food industry. Responsible. Sustainable. Professional. Profitable.

Mission: The National Pork Board harnesses the resources of all producers to capture opportunity, address challenges and satisfy customers.

Citations:

1. Sindelar JJ, Milkowski AL. Human safety controversies surrounding nitrate and nitrite in the diet. *Nitric Oxide* 2012; 26: 259-66.
2. Assembly of Life Sciences (U.S.) Committee on Nitrite and Curing Agents in Food. *The Health Effects of Nitrate, Nitrite, and N-Nitroso Compounds*. Washington DC: National Academies, 1981.
3. Food and Agriculture Organization, World Health Organization (FAO/WHO). Nitrite (and potential endogenous formation of N-nitroso compounds). In: *Safety Evaluation of Certain Food Additives and Contaminants*. Geneva: World Health Organization, Joint FAO/WHO Expert Committee on Food Additives, WHO Food Additives Series No. 50, 2003. <http://inchem.org/documents/jecfa/jecmono/v50je05.htm>. Accessed June 17, 2012.
4. American Medical Association. Report 9 of the Council on Scientific Affairs (A-04). Labeling of Nitrite Content of Processed Foods. 2004. <http://www.ama-assn.org/resources/doc/csaph/a04csa9-fulltext.pdf>. Accessed June 17, 2012.
5. Hord NG, Tang Y, Bryan NS. Food sources of nitrates and nitrites: the physiologic context for potential health benefits. *Am J Clin Nutr* 2009; 90: 1-10.
6. Cassens RG. Residual nitrite in cured meat. *Food Technol*. 1997; 51: 53-5.
7. Nunez De Gonzalez MT, Osburn WN, Hardin MD, et al. Survey of residual nitrite and nitrate in conventional and organic/natural/uncured/indirectly cured meats available at retail in the United States. *J Agric. Food Chem* 2012; 60: 3981-90.
8. Clark L. 5 Health Scares You Can Ignore. July 5, 2012. <http://abcnews.go.com/Health/Wellness/health-scares-ignore/story?id=16712712>. Accessed July 5, 2012.
9. Cosby K, Partovi KS, Crawford JH, et al. Nitrite reduction to nitric oxide by deoxyhemoglobin vasodilates the human circulation. *Nature Med*. 2003; 9: 1498-1505.
10. Webb AJ, Patel N, Loukogeorgakis S, et al. Acute blood pressure lowering, vasoprotective, and antiplatelet properties of dietary nitrate via bioconversion to nitrite. *Hypertension* 2008; 51: 784-90.
11. S M Tabatabaei, L Fritschi, M W Knuiman, T Boyle, B J Iacopetta, C Platell and J S Heyworth. Meat consumption and cooking practices and the risk of colorectal cancer. *European Journal of Clinical Nutrition* (2011) 65, 668–675.
12. Sanjiv Agarwal, Victor L. Fulgoni III and Eric P. Berg. Association of lunch meat consumption with nutrient intake, diet quality and health risk factors in U.S. children and adults: NHANES 2007–2010. Agarwal et al. *Nutrition Journal* (2015) 14:128.
13. Dominik D. Alexander, PhD, MSPH, Douglas L. Weed, MD, PhD, Paula E. Miller, MPH, Muhima A. Mohamed, MPH. Red Meat and Colorectal Cancer: A Quantitative Update on the State of the Epidemiologic Science. *Journal of the American College of Nutrition*, Vol. 0, No. 0, 1–23 (2015).
14. Briseis Aschebrook-Kilfoy*, Amanda J. Cross, Rachael Z. Stolzenberg-Solomon, Arthur Schatzkin, Albert R. Hollenbeck, Rashmi Sinha, and Mary H. Ward. Pancreatic Cancer and Exposure to Dietary Nitrate and Nitrite in the NIH-AARP Diet and Health Study. *American Journal of Epidemiology*

Published by Oxford University Press on behalf of the Johns Hopkins Bloomberg School of Public Health 2011.

15. D Lightsey, National Council Against Health Fraud and Quackwatch. Department of Family and Consumer Education, Bakersfield College, Bakersfield, CA, USA; Glinn and Giordano Physical Therapy Inc., Bakersfield, CA, USA Comment on 'Red and processed meat consumption and risk of pancreatic cancer: meta-analysis of prospective studies. *British Journal of Cancer* advance online publication, 17 July 2012.
16. Luis D. Boada, Henríquez-Hernández, O.P. Luzardo. The impact of red and processed meat consumption on cancer and other health outcomes: Epidemiological evidences. *Food and Chemical Toxicology* Volume 92, June 2016, Pages 236–244.
17. Nadia M. Bastide, Fatima Chenni, Marc Audebert, Raphaelle L. Santarelli, Sylviane Tach, Nathalie Naud, Maryse Baradat, Isabelle Jouanin, Reggie Surya, Ditte A. Hobbs, Gunter G. Kuhnle, Isabelle Raymond-Letron, Françoise Gueraud, Denis E. Corpet, and Fabrice H.F. Pierre. A Central Role for Heme Iron in Colon Carcinogenesis Associated with Red Meat Intake. January 15, 2015 American Association for Cancer Research.
18. AMJ Gilsing, MP Weijenberg, RA Goldbohm, PC Dagnelie, PA van den Brandt and LJ Schouten. Vegetarianism, low meat consumption and the risk of lung, postmenopausal breast and prostate cancer in a population-based cohort study. *European Journal of Clinical Nutrition* (2016) 70, 723–729.
19. Eunjung Kim, Desire Coelho, François Blachier. Review of the association between meat consumption and risk of colorectal cancer. *Nutrition Research* (2013) 983–994.
20. Murphy, M., Spungen, J., Bi, X., & Barraj, L. "Fresh and fresh lean pork are substantial sources of key nutrients when these products are consumed by adults in the United States." *Nutrition Research*/Volume 31. Issue 10 (2011): 776-783.
21. Rodriguez NR. Introduction to Protein Summit 2.0: Continued exploration of the impact of high-quality protein on optimal health. *American Journal of Clinical Nutrition*. 2015.
22. Institute of Medicine, Food and Nutrition Board. *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fatty Acids, Cholesterol, Protein and Amino Acids*. Washington, DC: The National Academies Press, 2002/2005.
23. Leidy HJ, Clifton PM, Astrup A, Wycherley TP, Westerterp-Plantenga MS, Luscombe-Marsh ND, Woods SC, Mattes RD. The role of protein in weight loss and maintenance. *American Journal of Clinical Nutrition*. 2015.
24. Layman DK, Anthony TG, Rasmussen BB, Adams SH, Lynch CJ, Brinkworth GD, Davis TA. Defining meal requirements for protein to optimize metabolic roles of amino acids. *American Journal of Clinical Nutrition*. 2015.
25. Paddon-Jones D, Campbell WW, Jacques PF, Kritchevsky SB, Moore L, Rodriguez N, Van Loon LJC. Protein and healthy aging. *American Journal of Clinical Nutrition*. 2015.
26. Phillips SM, Fulgoni III VL, Heaney RP, Nicklas TA, Slavin JL, Weaver CM. Commonly consumed protein foods contribute to nutrient intake, diet quality, and nutrient adequacy. *American Journal of Clinical Nutrition*. 2015. 7 Rodriguez NR, Miller SL. Effective translation of current dietary guidance: Understanding and communicating the concepts of minimal and optimal levels of dietary protein. *American Journal of Clinical Nutrition*. 2015.

27. Meta-analysis of prospective studies of red meat consumption and colorectal cancer. Alexander DD, Weed DL, Cushing CA, Lowe KA. *Eur J Cancer Prev*. 2011 May 2. [Epub ahead of print]
28. Red meat and colorectal cancer: a critical summary of prospective epidemiologic studies. Alexander DD, Cushing CA. *Obes Rev*. 2011 May;12(5):e472-93. doi: 10.1111/j.1467-789X.2010.00785.x.
29. Processed meat and colorectal cancer: a quantitative review of prospective epidemiologic studies. Alexander DD, Miller AJ, Cushing CA, Lowe KA. *Eur J Cancer Prev*. 2010 Sep;19(5):328-41.
30. Meta-analysis of animal fat or animal protein intake and colorectal cancer. Alexander DD, Cushing CA, Lowe KA, Scourman B, Roberts MA. *Am J Clin Nutr* 2009; 89(5):1402–1409.
31. Alexander, D.D. Red Meat and Processed Meat Consumption and Cancer: A Technical Summary of the Epidemiologic Evidence, 2010.
32. Kushi, L.H. et al. American Cancer Society Guidelines on Nutrition and Physical Activity for Cancer Prevention: Reducing the Risk of Cancer With Healthy Food Choices and Physical Activity. *CA: A Cancer Journal for Clinicians* 2006; 56:254-281.
33. National Toxicology Program Board of Scientific Counselors Technical Reports Review Subcommittee, May 18, 2000 Summary Minutes.
34. Nicolas Stettler, Mary M Murphy, Leila M Barraij, Kimberly M Smith, Rexford S Ahima. Systematic review of clinical studies related to pork intake and metabolic syndrome or its components. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*. Volume 2013:6; 347-357.
35. *Mortality in Vegetarians and Non-vegetarians: Detailed Findings From a Collaborative Analysis of 5 Prospective Studies*. Key & others, *AJCN*, 70:516S-24S, 1999.
36. *Health Effects of Vegetarian and Vegan Diets*. Key & others, *Proc.Nutr.Soc*. 65:35-41, 2006.
37. Fulgoni VL. Current protein intake in America: analysis of the National Health and Nutrition Examination Survey, 2003-2004. *Am J Clin Nutr* 2008; 87(suppl): 1554s-57s.
38. Hord NG, Tang Y, Bryan S. Food sources of nitrates and nitrites: the physiologic content for potential health benefits. *Am J Clin Nutr* 2009; 90: 1-10.
39. Gannon MC, Nuttall FQ, Saeed A, Jordan K, Hoover H. An increase in dietary protein improves the blood glucose response in persons with type 2 diabetes. *Am J Clin Nutr* 2003; 78: 671–672.
40. Gannon MC, Nuttall FQ. Effect of a high-protein, low-carbohydrate diet on blood glucose control in people with type 2 diabetes. *Diabetes* 2004; 53:2375-2382.
41. Parker B, Noakes M, Luscombe N, Clifton P. Effect of a high-protein, high-monounsaturated fat weight loss diet on glycemic control and lipid levels in type 2 diabetes. *Diabetes Care* 2002; 25: 425-430.
42. Wheeler, M. L., Fineberg, S. E., Fineberg, N. S., Gibson, R. G. & Hackward, L. L. Animal versus plant protein meals in individuals with type 2 diabetes and microalbuminuria: effects on renal, glycemic, and lipid parameters. *Diabetes Care* 2002; 25:1277-1282.
43. Gannon MC, Nuttall JA, Damberg G, Gupta V, Nuttall FQ. Effect of protein ingestion on the glucose appearance rate in people with type 2 diabetes. *J Clin Endocrinol Metab* 2001; 86:1040–1047.
44. Mikkelsen PB, Toubro S, Astrup A. Effect of fat-reduced diets on 24-h energy expenditure: comparisons between animal protein, vegetable protein, and carbohydrate. *Am J Clin Nutr* 2000;72:1135–41.

On-going Research:

45. Kevin C. Maki, Effects of a lean pork-containing, high-protein breakfast on indices of satiety and metabolic health in men and women with pre-diabetes.
46. Wayne Campbell Purdue University, Effects of incorporating red meat into a Mediterranean-style dietary pattern on cardiometabolic and emotional well-being.