F1_M3

C Number:	C20005
Lock Date:	10/10/2006
Cage Range:	ALL
Date Range:	ALL
Reasons For Removal:	ALL
Removal Date Range:	ALL
Treatment Groups:	Include ALL
Study Gender:	Both
TDMSE Version:	2.0.0

HISTORICAL CONTROL STUDIES
From October 2007 Report

SAME ROUTE

88123-07 -- Formamide (GAVAGE WATER)
95011-06 -- 5-(Hydroxymethyl)-2-furfural (GAVAGE WATER)

ALL ROUTES

88138-04 -- Benzophenone (ORAL FEED)
99027-04 -- Chromium picolinate monohydrate (ORAL FEED)
05174-06 -- Cresols (ORAL FEED)
96011-06 -- Cumene (INHALATION AIR)
97003-08 -- 1,2-Dibromo-2,4-dicyanobutane (SKIN ETHANOL)
93020-08 -- Diisopropylcarbodiimide (SKIN ETHANOL)
88004-06 -- Divinylbenzene (INHALATION AIR)
88123-07 -- Formamide (GAVAGE WATER)
95011-06 -- 5-(Hydroxymethyl)-2-furfural (GAVAGE WATER)
88105-04 -- Isoeugenol (GAVAGE CORN OIL)
97006-02 -- Methyl isobutyl ketone (INHALATION AIR)
91005-08 -- Methylene blue trihydrate (GAVAGE METHYLCELLULOSE)
92013-06 -- 4-Methylimidazole (ORAL FEED)
88006-04 -- alpha-Methylstyrene (INHALATION AIR)
99023-04 -- beta-Myrcene (GAVAGE CORN OIL)
97008-06 -- Propargyl alcohol (INHALATION AIR)
20114-06 -- Sodium dichromate dihydrate (VI) (ORAL WATER)
96019-06 -- Water disinfection byproducts (Bromochloroacetic acid) (ORAL WATER)
96017-06 -- Water disinfection byproducts (Dibromoacetic acid) (ORAL WATER)
96015-06 -- Water disinfection byproducts (Dibromoacetonitrile) (ORAL WATER)

FOR ALL DOSES THE TUMOR RATES IN THE FOLLOWING TISSUES/ORGANS ARE BASED ON NUMBER OF TISSUES EXAMINED. IN OTHER TISSUES/ORGANS RATES ARE BASED ON THE NUMBER OF ANIMALS NECROPSIED.

Adrenal Cortex
Adrenal Medulla
Bone Marrow
Epididymis
Heart
Islets, Pancreatic
Kidney
Liver
Lung
Nose
Ovary
Pituitary Gland
Prostate
Salivary Glands
Spleen
Testes
Thyroid Gland

SUMMARY OF STATISTICALLY SIGNIFICANT (P<=.05) RESULTS IN THE ANALYSIS OF Ginseng

MALE MICE

Organ	Morphology
Intestine Small: Site Unspecified	Carcinoma or Adenoma
Kidney: Renal Tubule	Adenoma
Lung	Alveolar/Bronchiolar Adenoma
	Alveolar/Bronchiolar Carcinoma or Alveolar/Bronchiolar Adenoma
All Organs	Malignant Lymphoma: Histiocytic, Lymphocytic, Mixed, NOS, or Undifferentiated Cell Type
	Benign Tumors

FEMALE MICE

Organ	Morphology
Ovary	Cystadenoma
Pituitary Gland: Pars Distalis or Unspecified Site	Adenoma
Skin	Fibrous Histiocytoma
	Sarcoma
All Organs	Hemangiosarcoma
	Hemangiosarcoma or Hemangioma
	Malignant Tumors

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Adrenal Cortex Adenoma				
TUMOR RATES				
OVERALL (a)	1/50 (2%)	2/50 (4%)	4/50 (8%)	2/50 (4%)
POLY-3 RATE (b)	1/43.85	2/44.2	4/46.18	2/42.27
POLY-3 PERCENT (g)	2.3%	4.5%	8.7%	4.7%
TERMINAL (d)	1/32 (3%)	2/32 (6%)	4/38 (11%)	2/32 (6%)
FIRST INCIDENCE	730 (T)	730 (T)	730 (T)	730 (T)
HC TUMORS SAME RTE	7/99 (7%)			
HC TUMORS ALL RTES	64/943 (7%)			
STATISTICAL TESTS				
POLY 3	P=0.362	P=0.503	P=0.195	P=0.487
POLY 1.5	P=0.366	P=0.501	P=0.189	P=0.489
POLY 6	P=0.361	P=0.504	P=0.203	P=0.487
COCH-ARM / FISHERS	P=0.387	P=0.500	P=0.181	P=0.500
MAX-ISO-POLY-3	P=0.289	P=0.283	P=0.099	P=0.273
HISTCONT SAME RTE	P=0.381	P=0.395	P=0.138	P=0.381
HISTCONT ALL RTES	P=0.337	P=0.355	P=0.114	P=0.341

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Males			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Bone **Osteosarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/43.85	0/44.2	0/46.18	0/42.27
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/32 (0%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	2/949 (0%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Males			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Harderian Gland Adenoma				
TUMOR RATES	#	#	#	#
OVERALL (a)	11/50 (22%)	7/50 (14%)	11/50 (22%)	15/50 (30%)
POLY-3 RATE (b)	11/46.02	7/45.7	11/48.25	15/43.99
POLY-3 PERCENT (g)	23.9%	15.3%	22.8%	34.1%
TERMINAL (d)	5/32 (16%)	3/32 (9%)	5/38 (13%)	11/32 (34%)
FIRST INCIDENCE	562	542	486	483
HC TUMORS SAME RTE	17/100 (17%)			
HC TUMORS ALL RTES	123/949 (13%)			
STATISTICAL TESTS				
POLY 3	P=0.083	P=0.220N	P=0.546N	P=0.201
POLY 1.5	P=0.087	P=0.219N	P=0.564N	P=0.213
POLY 6	P=0.080	P=0.223N	P=0.525N	P=0.189
COCH-ARM / FISHERS	P=0.102	P=0.218N	P=0.595N	P=0.247
MAX-ISO-POLY-3	P=0.091	P=0.150N	P=0.449N	P=0.146
HISTCONT SAME RTE	P=0.099	P=1.000	P=0.383	P=0.143
HISTCONT ALL RTES	P=0.002**	P=0.551	P=0.139	P=0.020*

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Harderian Gland **Carcinoma or Adenoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	12/50 (24%)	7/50 (14%)	11/50 (22%)	15/50 (30%)
POLY-3 RATE (b)	12/46.15	7/45.7	11/48.25	15/43.99
POLY-3 PERCENT (g)	26%	15.3%	22.8%	34.1%
TERMINAL (d)	5/32 (16%)	3/32 (9%)	5/38 (13%)	11/32 (34%)
FIRST INCIDENCE	562	542	486	483
HC TUMORS SAME RTE	20/100 (20%)			
HC TUMORS ALL RTES	146/949 (15%)			
STATISTICAL TESTS				
POLY 3	P=0.120	P=0.156N	P=0.452N	P=0.271
POLY 1.5	P=0.125	P=0.155N	P=0.469N	P=0.285
POLY 6	P=0.115	P=0.161N	P=0.432N	P=0.254
COCH-ARM / FISHERS	P=0.145	P=0.154N	P=0.500N	P=0.326
MAX-ISO-POLY-3	P=0.117	P=0.103N	P=0.361N	P=0.206
HISTCONT SAME RTE	P=0.159	P=1.000	P=0.579	P=0.192
HISTCONT ALL RTES	P=0.016*	P=1.000	P=0.290	P=0.052

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Intestine Small: Jejunum				
Carcinoma				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	1/50 (2%)	0/50 (0%)	2/50 (4%)
POLY-3 RATE (b)	0/43.85	1/44.2	0/46.18	2/42.52
POLY-3 PERCENT (g)	0%	2.3%	0%	4.7%
TERMINAL (d)	0/32 (0%)	1/32 (3%)	0/38 (0%)	1/32 (3%)
FIRST INCIDENCE	---	730 (T)	---	666
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	8/949 (1%)			
STATISTICAL TESTS				
POLY 3	P=0.137	P=0.502	(e)	P=0.230
POLY 1.5	P=0.137	P=0.501	(e)	P=0.231
POLY 6	P=0.138	P=0.502	(e)	P=0.231
COCH-ARM / FISHERS	P=0.140	P=0.500	(e)	P=0.247
MAX-ISO-POLY-3	P=0.077	P=0.161	(e)	P=0.076
HISTCONT SAME RTE	P=0.104	P=0.119	(e)	P=0.106
HISTCONT ALL RTES	P=0.088	P=0.192	(e)	P=0.086

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Intestine Small: Site Unspecified				
Carcinoma				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	1/50 (2%)	0/50 (0%)	2/50 (4%)
POLY-3 RATE (b)	0/43.85	1/44.2	0/46.18	2/42.52
POLY-3 PERCENT (g)	0%	2.3%	0%	4.7%
TERMINAL (d)	0/32 (0%)	1/32 (3%)	0/38 (0%)	1/32 (3%)
FIRST INCIDENCE	---	730 (T)	---	666
HC TUMORS SAME RTE	2/100 (2%)			
HC TUMORS ALL RTES	12/949 (1%)			
STATISTICAL TESTS				
POLY 3	P=0.137	P=0.502	(e)	P=0.230
POLY 1.5	P=0.137	P=0.501	(e)	P=0.231
POLY 6	P=0.138	P=0.502	(e)	P=0.231
COCH-ARM / FISHERS	P=0.140	P=0.500	(e)	P=0.247
MAX-ISO-POLY-3	P=0.077	P=0.161	(e)	P=0.076
HISTCONT SAME RTE	P=0.104	P=0.226	(e)	P=0.106
HISTCONT ALL RTES	P=0.088	P=0.192	(e)	P=0.086

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Males			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Intestine Small: Site Unspecified				
Carcinoma or Adenoma				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	2/50 (4%)	1/50 (2%)	3/50 (6%)
POLY-3 RATE (b)	0/43.85	2/44.2	1/46.18	3/42.91
POLY-3 PERCENT (g)	0%	4.5%	2.2%	7%
TERMINAL (d)	0/32 (0%)	2/32 (6%)	1/38 (3%)	1/32 (3%)
FIRST INCIDENCE	---	730 (T)	730 (T)	617
HC TUMORS SAME RTE	2/100 (2%)			
HC TUMORS ALL RTES	20/949 (2%)			
STATISTICAL TESTS				
POLY 3	P=0.100	P=0.239	P=0.510	P=0.115
POLY 1.5	P=0.100	P=0.238	P=0.506	P=0.115
POLY 6	P=0.103	P=0.239	P=0.516	P=0.116
COCH-ARM / FISHERS	P=0.104	P=0.247	P=0.500	P=0.121
MAX-ISO-POLY-3	P=0.060	P=0.077	P=0.172	P=0.038*
HISTCONT SAME RTE	P=0.081	P=0.108	P=0.241	P=0.051
HISTCONT ALL RTES	P=0.068	P=0.089	P=0.206	P=0.040*

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Males			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Islets, Pancreatic
Carcinoma or Adenoma

TUMOR RATES

OVERALL (a)	2/50 (4%)	1/50 (2%)	1/50 (2%)	1/50 (2%)
POLY-3 RATE (b)	2/44.12	1/44.26	1/46.76	1/42.27
POLY-3 PERCENT (g)	4.5%	2.3%	2.1%	2.4%
TERMINAL (d)	0/32 (0%)	0/32 (0%)	0/38 (0%)	1/32 (3%)
FIRST INCIDENCE	695	715	549	730 (T)
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	11/940 (1%)			

STATISTICAL TESTS

POLY 3	P=0.415N	P=0.499N	P=0.480N	P=0.515N
POLY 1.5	P=0.414N	P=0.500N	P=0.488N	P=0.512N
POLY 6	P=0.413N	P=0.499N	P=0.471N	P=0.517N
COCH-ARM / FISHERS	P=0.409N	P=0.500N	P=0.500N	P=0.500N
MAX-ISO-POLY-3	P=0.433N	P=0.280N	P=0.269N	P=0.297N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Males			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Kidney: Renal Tubule Adenoma				
TUMOR RATES				
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	2/50 (4%)
POLY-3 RATE (b)	0/43.85	0/44.2	0/46.18	2/42.27
POLY-3 PERCENT (g)	0%	0%	0%	4.7%
TERMINAL (d)	0/32 (0%)	0/32 (0%)	0/38 (0%)	2/32 (6%)
FIRST INCIDENCE	---	---	---	730 (T)
HC TUMORS SAME RTE	1/100 (1%)			
HC TUMORS ALL RTES	10/945 (1%)			
STATISTICAL TESTS				
POLY 3	P=0.044*	(e)	(e)	P=0.229
POLY 1.5	P=0.044*	(e)	(e)	P=0.230
POLY 6	P=0.044*	(e)	(e)	P=0.229
COCH-ARM / FISHERS	P=0.046*	(e)	(e)	P=0.247
MAX-ISO-POLY-3	P=0.032*	(e)	(e)	P=0.076
HISTCONT SAME RTE	P=0.043*	(e)	(e)	P=0.106
HISTCONT ALL RTES	P=0.036*	(e)	(e)	P=0.086

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Kidney: Renal Tubule
Carcinoma or Adenoma

TUMOR RATES

OVERALL (a)	1/50 (2%)	0/50 (0%)	0/50 (0%)	2/50 (4%)
POLY-3 RATE (b)	1/43.85	0/44.2	0/46.18	2/42.27
POLY-3 PERCENT (g)	2.3%	0%	0%	4.7%
TERMINAL (d)	1/32 (3%)	0/32 (0%)	0/38 (0%)	2/32 (6%)
FIRST INCIDENCE	730 (T)	---	---	730 (T)
HC TUMORS SAME RTE	1/100 (1%)			
HC TUMORS ALL RTES	13/945 (1%)			

STATISTICAL TESTS

POLY 3	P=0.242	P=0.498N	P=0.490N	P=0.487
POLY 1.5	P=0.243	P=0.499N	P=0.494N	P=0.489
POLY 6	P=0.243	P=0.498N	P=0.484N	P=0.487
COCH-ARM / FISHERS	P=0.246	P=0.500N	P=0.500N	P=0.500
MAX-ISO-POLY-3	P=0.130	P=0.158N	P=0.158N	P=0.273
HISTCONT SAME RTE	P=0.147	(e)	(e)	P=0.141
HISTCONT ALL RTES	P=0.149	(e)	(e)	P=0.225

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Liver **Hemangiosarcoma**				
TUMOR RATES				
OVERALL (a)	2/50 (4%)	4/50 (8%)	1/50 (2%)	3/50 (6%)
POLY-3 RATE (b)	2/43.99	4/44.91	1/46.18	3/43.07
POLY-3 PERCENT (g)	4.6%	8.9%	2.2%	7%
TERMINAL (d)	1/32 (3%)	2/32 (6%)	1/38 (3%)	1/32 (3%)
FIRST INCIDENCE	695	542	730 (T)	558
HC TUMORS SAME RTE	2/100 (2%)			
HC TUMORS ALL RTES	27/948 (3%)			
STATISTICAL TESTS				
POLY 3	P=0.537	P=0.347	P=0.483N	P=0.490
POLY 1.5	P=0.535	P=0.342	P=0.490N	P=0.490
POLY 6	P=0.542	P=0.351	P=0.475N	P=0.495
COCH-ARM / FISHERS	P=0.544	P=0.339	P=0.500N	P=0.500
MAX-ISO-POLY-3	P=0.504	P=0.210	P=0.271N	P=0.317
HISTCONT SAME RTE	P=0.281	P=0.093	P=1.000	P=0.107
HISTCONT ALL RTES	P=0.290	P=0.115	P=1.000	P=0.238

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Males			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Liver **Hepatocellular Adenoma**				
TUMOR RATES				
OVERALL (a)	34/50 (68%)	29/50 (58%)	33/50 (66%)	34/50 (68%)
POLY-3 RATE (b)	34/47.46	29/46.97	33/47.34	34/45.77
POLY-3 PERCENT (g)	71.6%	61.8%	69.7%	74.3%
TERMINAL (d)	26/32 (81%)	20/32 (63%)	28/38 (74%)	23/32 (72%)
FIRST INCIDENCE	339	542	549	429
HC TUMORS SAME RTE	57/100 (57%)			
HC TUMORS ALL RTES	465/948 (49%)			
STATISTICAL TESTS				
POLY 3	P=0.297	P=0.205N	P=0.508N	P=0.478
POLY 1.5	P=0.312	P=0.214N	P=0.508N	P=0.488
POLY 6	P=0.303	P=0.184N	P=0.496N	P=0.498
COCH-ARM / FISHERS	P=0.401	P=0.204N	P=0.500N	P=0.585N
MAX-ISO-POLY-3	P=0.343	P=0.148N	P=0.414N	P=0.384
HISTCONT SAME RTE	P=0.160	P=0.592	P=0.329	P=0.237
HISTCONT ALL RTES	P=0.054	P=0.397	P=0.229	P=0.171

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Males			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Liver **Hepatocellular Carcinoma**				
TUMOR RATES				
OVERALL (a)	14/50 (28%)	14/50 (28%)	15/50 (30%)	12/50 (24%)
POLY-3 RATE (b)	14/44.62	14/45.56	15/46.69	12/43.38
POLY-3 PERCENT (g)	31.4%	30.7%	32.1%	27.7%
TERMINAL (d)	11/32 (34%)	9/32 (28%)	13/38 (34%)	6/32 (19%)
FIRST INCIDENCE	617	576	632	563
HC TUMORS SAME RTE	26/100 (26%)			
HC TUMORS ALL RTES	260/948 (27%)			
STATISTICAL TESTS				
POLY 3	P=0.399N	P=0.564N	P=0.559	P=0.441N
POLY 1.5	P=0.396N	P=0.576N	P=0.536	P=0.439N
POLY 6	P=0.394N	P=0.548N	P=0.587	P=0.434N
COCH-ARM / FISHERS	P=0.364N	P=0.588N	P=0.500	P=0.410N
MAX-ISO-POLY-3	P=0.559N	P=0.473N	P=0.469	P=0.352N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Liver
Hepatocellular Carcinoma or Hepatoblastoma

TUMOR RATES

OVERALL (a)	14/50 (28%)	15/50 (30%)	15/50 (30%)	13/50 (26%)
POLY-3 RATE (b)	14/44.62	15/45.56	15/46.69	13/43.38
POLY-3 PERCENT (g)	31.4%	32.9%	32.1%	30%
TERMINAL (d)	11/32 (34%)	10/32 (31%)	13/38 (34%)	7/32 (22%)
FIRST INCIDENCE	617	576	632	563
HC TUMORS SAME RTE	28/100 (28%)			
HC TUMORS ALL RTES	285/948 (30%)			

STATISTICAL TESTS

POLY 3	P=0.464N	P=0.527	P=0.559	P=0.535N
POLY 1.5	P=0.461N	P=0.514	P=0.536	P=0.532N
POLY 6	P=0.459N	P=0.545	P=0.587	P=0.529N
COCH-ARM / FISHERS	P=0.426N	P=0.500	P=0.500	P=0.500N
MAX-ISO-POLY-3	P=0.642N	P=0.436	P=0.469	P=0.442N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Liver
Hepatocellular Carcinoma or Hepatocellular Adenoma

TUMOR RATES

OVERALL (a)	39/50 (78%)	38/50 (76%)	38/50 (76%)	39/50 (78%)
POLY-3 RATE (b)	39/47.94	38/48	38/47.85	39/45.97
POLY-3 PERCENT (g)	81.3%	79.2%	79.4%	84.8%
TERMINAL (d)	29/32 (91%)	25/32 (78%)	31/38 (82%)	27/32 (84%)
FIRST INCIDENCE	339	542	549	429
HC TUMORS SAME RTE	72/100 (72%)			
HC TUMORS ALL RTES	611/948 (65%)			

STATISTICAL TESTS

POLY 3	P=0.342	P=0.495N	P=0.508N	P=0.426
POLY 1.5	P=0.378	P=0.518N	P=0.509N	P=0.454
POLY 6	P=0.328	P=0.437N	P=0.489N	P=0.434
COCH-ARM / FISHERS	P=0.523	P=0.500N	P=0.500N	P=0.595N
MAX-ISO-POLY-3	P=0.435	P=0.389N	P=0.400N	P=0.320
HISTCONT SAME RTE	P=0.122	P=0.342	P=0.331	P=0.197
HISTCONT ALL RTES	P=0.069	P=0.289	P=0.284	P=0.193

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Liver
Hepatocellular Carcinoma, Hepatocellular Adenoma, or Hepatoblastoma

TUMOR RATES

OVERALL (a)	39/50 (78%)	38/50 (76%)	38/50 (76%)	39/50 (78%)
POLY-3 RATE (b)	39/47.94	38/48	38/47.85	39/45.97
POLY-3 PERCENT (g)	81.3%	79.2%	79.4%	84.8%
TERMINAL (d)	29/32 (91%)	25/32 (78%)	31/38 (82%)	27/32 (84%)
FIRST INCIDENCE	339	542	549	429
HC TUMORS SAME RTE	73/100 (73%)			
HC TUMORS ALL RTES	622/948 (66%)			

STATISTICAL TESTS

POLY 3	P=0.342	P=0.495N	P=0.508N	P=0.426
POLY 1.5	P=0.378	P=0.518N	P=0.509N	P=0.454
POLY 6	P=0.328	P=0.437N	P=0.489N	P=0.434
COCH-ARM / FISHERS	P=0.523	P=0.500N	P=0.500N	P=0.595N
MAX-ISO-POLY-3	P=0.435	P=0.389N	P=0.400N	P=0.320
HISTCONT SAME RTE	P=0.108	P=0.343	P=0.331	P=0.183
HISTCONT ALL RTES	P=0.102	P=0.328	P=0.323	P=0.225

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Lung **Alveolar/Bronchiolar Adenoma**				
TUMOR RATES				
OVERALL (a)	9/50 (18%)	3/50 (6%)	6/50 (12%)	16/50 (32%)
POLY-3 RATE (b)	9/43.91	3/44.26	6/46.53	16/44.44
POLY-3 PERCENT (g)	20.5%	6.8%	12.9%	36%
TERMINAL (d)	8/32 (25%)	2/32 (6%)	5/38 (13%)	9/32 (28%)
FIRST INCIDENCE	714	715	632	558
HC TUMORS SAME RTE	20/100 (20%)			
HC TUMORS ALL RTES	180/948 (19%)			
STATISTICAL TESTS				
POLY 3	P=0.008**	P=0.056N	P=0.246N	P=0.080
POLY 1.5	P=0.007**	P=0.058N	P=0.263N	P=0.076
POLY 6	P=0.009**	P=0.054N	P=0.224N	P=0.092
COCH-ARM / FISHERS	P=0.009**	P=0.061N	P=0.288N	P=0.083
MAX-ISO-POLY-3	P=0.004**	P=0.029N*	P=0.173N	P=0.051
HISTCONT SAME RTE	P=0.005**	P=1.000	P=1.000	P=0.069
HISTCONT ALL RTES	P=0.004**	P=1.000	P=1.000	P=0.056

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Lung
Alveolar/Bronchiolar Carcinoma

TUMOR RATES

OVERALL (a)	3/50 (6%)	3/50 (6%)	5/50 (10%)	3/50 (6%)
POLY-3 RATE (b)	3/44.68	3/44.71	5/46.25	3/42.67
POLY-3 PERCENT (g)	6.7%	6.7%	10.8%	7%
TERMINAL (d)	1/32 (3%)	1/32 (3%)	4/38 (11%)	2/32 (6%)
FIRST INCIDENCE	562	576	714	617
HC TUMORS SAME RTE	13/100 (13%)			
HC TUMORS ALL RTES	91/948 (10%)			

STATISTICAL TESTS

POLY 3	P=0.505	P=0.662N	P=0.375	P=0.641
POLY 1.5	P=0.512	P=0.662	P=0.369	P=0.646
POLY 6	P=0.502	P=0.663	P=0.384	P=0.639
COCH-ARM / FISHERS	P=0.537	P=0.661N	P=0.357	P=0.661N
MAX-ISO-POLY-3	P=0.555	P=0.499N	P=0.250	P=0.476
HISTCONT SAME RTE	P=0.687	P=1.000	P=0.350	P=0.629
HISTCONT ALL RTES	P=0.654	P=1.000	P=0.310	P=0.603

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Lung
Alveolar/Bronchiolar Carcinoma or Alveolar/Bronchiolar Adenoma

TUMOR RATES

OVERALL (a)	12/50 (24%)	6/50 (12%)	10/50 (20%)	19/50 (38%)
POLY-3 RATE (b)	12/44.74	6/44.77	10/46.6	19/44.83
POLY-3 PERCENT (g)	26.8%	13.4%	21.5%	42.4%
TERMINAL (d)	9/32 (28%)	3/32 (9%)	8/38 (21%)	11/32 (34%)
FIRST INCIDENCE	562	576	632	558
HC TUMORS SAME RTE	32/100 (32%)			
HC TUMORS ALL RTES	260/948 (27%)			

STATISTICAL TESTS

POLY 3	P=0.014*	P=0.091N	P=0.362N	P=0.089
POLY 1.5	P=0.014*	P=0.093N	P=0.378N	P=0.086
POLY 6	P=0.016*	P=0.089N	P=0.341N	P=0.100
COCH-ARM / FISHERS	P=0.017*	P=0.096N	P=0.405N	P=0.097
MAX-ISO-POLY-3	P=0.011*	P=0.054N	P=0.279N	P=0.058
HISTCONT SAME RTE	P=0.013*	P=1.000	P=1.000	P=0.079
HISTCONT ALL RTES	P=0.011*	P=1.000	P=1.000	P=0.064

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Pituitary Gland: Pars Distalis or Unspecified Site
Adenoma

TUMOR RATES

OVERALL (a)	1/50 (2%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	1/43.85	0/44.2	0/46.18	0/42.27
POLY-3 PERCENT (g)	2.3%	0%	0%	0%
TERMINAL (d)	1/32 (3%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	730 (T)	---	---	---
HC TUMORS SAME RTE	1/98 (1%)			
HC TUMORS ALL RTES	3/924 (0%)			

STATISTICAL TESTS

POLY 3	P=0.299N	P=0.498N	P=0.490N	P=0.507N
POLY 1.5	P=0.301N	P=0.499N	P=0.494N	P=0.506N
POLY 6	P=0.298N	P=0.498N	P=0.484N	P=0.507N
COCH-ARM / FISHERS	P=0.306N	P=0.500N	P=0.500N	P=0.500N
MAX-ISO-POLY-3	P=0.136N	P=0.158N	P=0.158N	P=0.168N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Skin
Basal or Sq. Cell Carcinoma, Carcinoma, Basosq. Tumor (M or B), Basal Cell Adenoma, Adenoma, Papilloma, Sq Papilloma, Keratoacanthoma, Trichoepithelioma

TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	2/50 (4%)	0/50 (0%)
POLY-3 RATE (b)	0/43.85	0/44.2	2/46.18	0/42.27
POLY-3 PERCENT (g)	0%	0%	4.3%	0%
TERMINAL (d)	0/32 (0%)	0/32 (0%)	2/38 (5%)	0/32 (0%)
FIRST INCIDENCE	---	---	730 (T)	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	2/949 (0%)			

STATISTICAL TESTS				
POLY 3	P=0.569	(e)	P=0.249	(e)
POLY 1.5	P=0.574	(e)	P=0.244	(e)
POLY 6	P=0.566	(e)	P=0.255	(e)
COCH-ARM / FISHERS	P=0.595	(e)	P=0.247	(e)
MAX-ISO-POLY-3	P=0.269	(e)	P=0.087	(e)
HISTCONT SAME RTE	P=0.356	(e)	P=0.110	(e)
HISTCONT ALL RTES	P=0.189	(e)	P=0.004**	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Males			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Skin
Fibroma, Fibrosarcoma, Sarcoma, Myxoma, Myxosarcoma, or Fibrous Histiocytoma

TUMOR RATES	#	#	#	#
OVERALL (a)	1/50 (2%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	1/43.85	0/44.2	0/46.18	0/42.27
POLY-3 PERCENT (g)	2.3%	0%	0%	0%
TERMINAL (d)	1/32 (3%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	730 (T)	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	12/949 (1%)			

STATISTICAL TESTS				
POLY 3	P=0.299N	P=0.498N	P=0.490N	P=0.507N
POLY 1.5	P=0.301N	P=0.499N	P=0.494N	P=0.506N
POLY 6	P=0.298N	P=0.498N	P=0.484N	P=0.507N
COCH-ARM / FISHERS	P=0.306N	P=0.500N	P=0.500N	P=0.500N
MAX-ISO-POLY-3	P=0.136N	P=0.158N	P=0.158N	P=0.168N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Skin **Fibrosarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	1/50 (2%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	1/43.85	0/44.2	0/46.18	0/42.27
POLY-3 PERCENT (g)	2.3%	0%	0%	0%
TERMINAL (d)	1/32 (3%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	730 (T)	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	0/949 (0%)			
STATISTICAL TESTS				
POLY 3	P=0.299N	P=0.498N	P=0.490N	P=0.507N
POLY 1.5	P=0.301N	P=0.499N	P=0.494N	P=0.506N
POLY 6	P=0.298N	P=0.498N	P=0.484N	P=0.507N
COCH-ARM / FISHERS	P=0.306N	P=0.500N	P=0.500N	P=0.500N
MAX-ISO-POLY-3	P=0.136N	P=0.158N	P=0.158N	P=0.168N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Skin
Fibrosarcoma, Sarcoma, Myxosarcoma, or Fibrous Histiocytoma

TUMOR RATES	#	#	#	#
OVERALL (a)	1/50 (2%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	1/43.85	0/44.2	0/46.18	0/42.27
POLY-3 PERCENT (g)	2.3%	0%	0%	0%
TERMINAL (d)	1/32 (3%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	730 (T)	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	9/949 (1%)			

STATISTICAL TESTS				
POLY 3	P=0.299N	P=0.498N	P=0.490N	P=0.507N
POLY 1.5	P=0.301N	P=0.499N	P=0.494N	P=0.506N
POLY 6	P=0.298N	P=0.498N	P=0.484N	P=0.507N
COCH-ARM / FISHERS	P=0.306N	P=0.500N	P=0.500N	P=0.500N
MAX-ISO-POLY-3	P=0.136N	P=0.158N	P=0.158N	P=0.168N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Males			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Skin **Fibrous Histiocytoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/43.85	0/44.2	0/46.18	0/42.27
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/32 (0%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	5/949 (1%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Skin **Sarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/43.85	0/44.2	0/46.18	0/42.27
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/32 (0%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	4/949 (0%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Males			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Skin **Schwannoma Malignant**				
TUMOR RATES	#	#	#	#
OVERALL (a)	1/50 (2%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	1/43.99	0/44.2	0/46.18	0/42.27
POLY-3 PERCENT (g)	2.3%	0%	0%	0%
TERMINAL (d)	0/32 (0%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	695	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	1/949 (0%)			
STATISTICAL TESTS				
POLY 3	P=0.299N	P=0.499N	P=0.490N	P=0.508N
POLY 1.5	P=0.301N	P=0.500N	P=0.494N	P=0.506N
POLY 6	P=0.297N	P=0.499N	P=0.486N	P=0.509N
COCH-ARM / FISHERS	P=0.306N	P=0.500N	P=0.500N	P=0.500N
MAX-ISO-POLY-3	P=0.137N	P=0.158N	P=0.158N	P=0.168N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Skin **Squamous Cell Papilloma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	2/50 (4%)	0/50 (0%)
POLY-3 RATE (b)	0/43.85	0/44.2	2/46.18	0/42.27
POLY-3 PERCENT (g)	0%	0%	4.3%	0%
TERMINAL (d)	0/32 (0%)	0/32 (0%)	2/38 (5%)	0/32 (0%)
FIRST INCIDENCE	---	---	730 (T)	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	1/949 (0%)			
STATISTICAL TESTS				
POLY 3	P=0.569	(e)	P=0.249	(e)
POLY 1.5	P=0.574	(e)	P=0.244	(e)
POLY 6	P=0.566	(e)	P=0.255	(e)
COCH-ARM / FISHERS	P=0.595	(e)	P=0.247	(e)
MAX-ISO-POLY-3	P=0.269	(e)	P=0.087	(e)
HISTCONT SAME RTE	P=0.356	(e)	P=0.110	(e)
HISTCONT ALL RTES	P=0.137	(e)	P=0.002**	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Hemangioma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	1/50 (2%)	1/50 (2%)	1/50 (2%)	1/50 (2%)
POLY-3 RATE (b)	1/43.85	1/44.2	1/46.18	1/42.27
POLY-3 PERCENT (g)	2.3%	2.3%	2.2%	2.4%
TERMINAL (d)	1/32 (3%)	1/32 (3%)	1/38 (3%)	1/32 (3%)
FIRST INCIDENCE	730 (T)	730 (T)	730 (T)	730 (T)
HC TUMORS SAME RTE	2/100 (2%)			
HC TUMORS ALL RTES	7/949 (1%)			
STATISTICAL TESTS				
POLY 3	P=0.620	P=0.759N	P=0.749N	P=0.753
POLY 1.5	P=0.622	P=0.760N	P=0.753N	P=0.754
POLY 6	P=0.620	P=0.758N	P=0.743N	P=0.753
COCH-ARM / FISHERS	P=0.634	P=0.753N	P=0.753N	P=0.753N
MAX-ISO-POLY-3	P=0.731	P=0.497N	P=0.485N	P=0.489
HISTCONT SAME RTE	P=0.835	P=0.642	P=1.000	P=0.627
HISTCONT ALL RTES	P=0.211	P=0.274	P=0.281	P=0.267

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Hemangiosarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	4/50 (8%)	4/50 (8%)	2/50 (4%)	5/50 (10%)
POLY-3 RATE (b)	4/44.49	4/44.91	2/46.52	5/43.87
POLY-3 PERCENT (g)	9%	8.9%	4.3%	11.4%
TERMINAL (d)	2/32 (6%)	2/32 (6%)	1/38 (3%)	2/32 (6%)
FIRST INCIDENCE	578	542	637	429
HC TUMORS SAME RTE	5/100 (5%)			
HC TUMORS ALL RTES	60/949 (6%)			
STATISTICAL TESTS				
POLY 3	P=0.441	P=0.638N	P=0.317N	P=0.491
POLY 1.5	P=0.439	P=0.641N	P=0.325N	P=0.490
POLY 6	P=0.446	P=0.637N	P=0.307N	P=0.496
COCH-ARM / FISHERS	P=0.446	P=0.643N	P=0.339N	P=0.500
MAX-ISO-POLY-3	P=0.404	P=0.494N	P=0.190N	P=0.355
HISTCONT SAME RTE	P=0.165	P=0.212	P=1.000	P=0.129
HISTCONT ALL RTES	P=0.443	P=0.473	P=1.000	P=0.301

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Hemangiosarcoma or Hemangioma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	5/50 (10%)	5/50 (10%)	3/50 (6%)	6/50 (12%)
POLY-3 RATE (b)	5/44.49	5/44.91	3/46.52	6/43.87
POLY-3 PERCENT (g)	11.2%	11.1%	6.5%	13.7%
TERMINAL (d)	3/32 (9%)	3/32 (9%)	2/38 (5%)	3/32 (9%)
FIRST INCIDENCE	578	542	637	429
HC TUMORS SAME RTE	7/100 (7%)			
HC TUMORS ALL RTES	67/949 (7%)			
STATISTICAL TESTS				
POLY 3	P=0.446	P=0.625N	P=0.332N	P=0.490
POLY 1.5	P=0.443	P=0.627N	P=0.342N	P=0.488
POLY 6	P=0.451	P=0.623N	P=0.320N	P=0.495
COCH-ARM / FISHERS	P=0.451	P=0.630N	P=0.357N	P=0.500
MAX-ISO-POLY-3	P=0.432	P=0.493N	P=0.216N	P=0.365
HISTCONT SAME RTE	P=0.260	P=0.341	P=1.000	P=0.215
HISTCONT ALL RTES	P=0.263	P=0.351	P=1.000	P=0.200

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Histiocytic Sarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	1/50 (2%)	0/50 (0%)	1/50 (2%)	0/50 (0%)
POLY-3 RATE (b)	1/43.85	0/44.2	1/46.38	0/42.27
POLY-3 PERCENT (g)	2.3%	0%	2.2%	0%
TERMINAL (d)	1/32 (3%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	730 (T)	---	679	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	5/949 (1%)			
STATISTICAL TESTS				
POLY 3	P=0.410N	P=0.498N	P=0.748N	P=0.507N
POLY 1.5	P=0.410N	P=0.499N	P=0.753N	P=0.506N
POLY 6	P=0.410N	P=0.498N	P=0.741N	P=0.507N
COCH-ARM / FISHERS	P=0.405N	P=0.500N	P=0.753N	P=0.500N
MAX-ISO-POLY-3	P=0.270N	P=0.158N	P=0.484N	P=0.168N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

All Organs
Malignant Lymphoma: Histiocytic, Lymphocytic, Mixed, NOS, or Undifferentiated Cell Type

TUMOR RATES	#	#	#	#
OVERALL (a)	4/50 (8%)	1/50 (2%)	2/50 (4%)	0/50 (0%)
POLY-3 RATE (b)	4/44.17	1/44.2	2/46.75	0/42.27
POLY-3 PERCENT (g)	9.1%	2.3%	4.3%	0%
TERMINAL (d)	2/32 (6%)	1/32 (3%)	1/38 (3%)	0/32 (0%)
FIRST INCIDENCE	666	730 (T)	554	---
HC TUMORS SAME RTE	4/100 (4%)			
HC TUMORS ALL RTES	26/949 (3%)			

STATISTICAL TESTS				
POLY 3	P=0.054N	P=0.178N	P=0.311N	P=0.066N
POLY 1.5	P=0.054N	P=0.179N	P=0.321N	P=0.065N
POLY 6	P=0.053N	P=0.178N	P=0.300N	P=0.066N
COCH-ARM / FISHERS	P=0.052N	P=0.181N	P=0.339N	P=0.059N
MAX-ISO-POLY-3	P=0.027N*	P=0.083N	P=0.187N	P=0.024N*
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Osteosarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/43.85	0/44.2	0/46.18	0/42.27
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/32 (0%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	2/949 (0%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Osteosarcoma or Osteoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/43.85	0/44.2	0/46.18	0/42.27
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/32 (0%)	0/32 (0%)	0/38 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	2/949 (0%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Benign Tumors**				
TUMOR RATES	#	#	#	#
OVERALL (a)	38/50 (76%)	34/50 (68%)	40/50 (80%)	44/50 (88%)
POLY-3 RATE (b)	38/48.6	34/47.2	40/49.09	44/47.99
POLY-3 PERCENT (g)	78.2%	72%	81.5%	91.7%
TERMINAL (d)	26/32 (81%)	24/32 (75%)	31/38 (82%)	29/32 (91%)
FIRST INCIDENCE	339	542	486	429
HC TUMORS SAME RTE	73/100 (73%)			
HC TUMORS ALL RTES	634/949 (67%)			
STATISTICAL TESTS				
POLY 3	P=0.019*	P=0.318N	P=0.438	P=0.050*
POLY 1.5	P=0.018*	P=0.301N	P=0.421	P=0.050*
POLY 6	P=0.024*	P=0.327N	P=0.469	P=0.060
COCH-ARM / FISHERS	P=0.032*	P=0.252N	P=0.405	P=0.096
MAX-ISO-POLY-3	P=0.024*	P=0.241N	P=0.339	P=0.027*
HISTCONT SAME RTE	P=0.032*	P=1.000	P=0.457	P=0.035*
HISTCONT ALL RTES	P=0.026*	P=0.581	P=0.293	P=0.028*

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Malignant Tumors**				
TUMOR RATES	#	#	#	#
OVERALL (a)	26/50 (52%)	22/50 (44%)	24/50 (48%)	22/50 (44%)
POLY-3 RATE (b)	26/46.26	22/46.62	24/48.15	22/45.37
POLY-3 PERCENT (g)	56.2%	47.2%	49.8%	48.5%
TERMINAL (d)	17/32 (53%)	13/32 (41%)	17/38 (45%)	12/32 (38%)
FIRST INCIDENCE	562	542	554	429
HC TUMORS SAME RTE	54/100 (54%)			
HC TUMORS ALL RTES	482/949 (51%)			
STATISTICAL TESTS				
POLY 3	P=0.322N	P=0.251N	P=0.339N	P=0.296N
POLY 1.5	P=0.325N	P=0.262N	P=0.369N	P=0.299N
POLY 6	P=0.313N	P=0.243N	P=0.304N	P=0.289N
COCH-ARM / FISHERS	P=0.299N	P=0.274N	P=0.421N	P=0.274N
MAX-ISO-POLY-3	P=0.376N	P=0.190N	P=0.271N	P=0.230N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Males		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Malignant and Benign Tumors**				
TUMOR RATES	#	#	#	#
OVERALL (a)	46/50 (92%)	44/50 (88%)	46/50 (92%)	47/50 (94%)
POLY-3 RATE (b)	46/49.14	44/48.24	46/50	47/48.19
POLY-3 PERCENT (g)	93.6%	91.2%	92%	97.5%
TERMINAL (d)	30/32 (94%)	29/32 (91%)	34/38 (90%)	31/32 (97%)
FIRST INCIDENCE	339	542	486	429
HC TUMORS SAME RTE	90/100 (90%)			
HC TUMORS ALL RTES	812/949 (86%)			
STATISTICAL TESTS				
POLY 3	P=0.212	P=0.473N	P=0.532N	P=0.315
POLY 1.5	P=0.198	P=0.454N	P=0.583N	P=0.309
POLY 6	P=0.253	P=0.470N	P=0.461N	P=0.366
COCH-ARM / FISHERS	P=0.319	P=0.370N	P=0.643N	P=0.500
MAX-ISO-POLY-3	P=0.223	P=0.320N	P=0.376N	P=0.159
HISTCONT SAME RTE	P=0.297	P=0.604	P=0.564	P=0.223
HISTCONT ALL RTES	P=0.108	P=0.419	P=0.378	P=0.189

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Adrenal Cortex Adenoma				
TUMOR RATES				
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	0/43.48	0/43.18	0/40.8
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/99 (0%)			
HC TUMORS ALL RTES	3/994 (0%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Bone **Osteosarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	2/50 (4%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	2/44.91	0/43.18	0/40.8
POLY-3 PERCENT (g)	0%	4.5%	0%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	449	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	3/999 (0%)			
STATISTICAL TESTS				
POLY 3	P=0.439N	P=0.228	(e)	(e)
POLY 1.5	P=0.430N	P=0.232	(e)	(e)
POLY 6	P=0.445N	P=0.222	(e)	(e)
COCH-ARM / FISHERS	P=0.405N	P=0.247	(e)	(e)
MAX-ISO-POLY-3	P=0.297N	P=0.076	(e)	(e)
HISTCONT SAME RTE	(h)	(h)	(e)	(e)
HISTCONT ALL RTES	(h)	(h)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Harderian Gland Adenoma				
TUMOR RATES	#	#	#	#
OVERALL (a)	7/50 (14%)	7/50 (14%)	9/50 (18%)	8/50 (16%)
POLY-3 RATE (b)	7/47.08	7/43.88	9/43.75	8/40.8
POLY-3 PERCENT (g)	14.9%	16%	20.6%	19.6%
TERMINAL (d)	5/38 (13%)	6/31 (19%)	6/34 (18%)	8/32 (25%)
FIRST INCIDENCE	697	618	654	729 (T)
HC TUMORS SAME RTE	11/100 (11%)			
HC TUMORS ALL RTES	102/999 (10%)			
STATISTICAL TESTS				
POLY 3	P=0.290	P=0.558	P=0.332	P=0.381
POLY 1.5	P=0.318	P=0.580	P=0.336	P=0.413
POLY 6	P=0.275	P=0.532	P=0.337	P=0.358
COCH-ARM / FISHERS	P=0.409	P=0.613N	P=0.393	P=0.500
MAX-ISO-POLY-3	P=0.443	P=0.444	P=0.247	P=0.293
HISTCONT SAME RTE	P=0.142	P=0.304	P=0.155	P=0.185
HISTCONT ALL RTES	P=0.038*	P=0.319	P=0.126	P=0.175

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Harderian Gland **Carcinoma or Adenoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	7/50 (14%)	7/50 (14%)	9/50 (18%)	8/50 (16%)
POLY-3 RATE (b)	7/47.08	7/43.88	9/43.75	8/40.8
POLY-3 PERCENT (g)	14.9%	16%	20.6%	19.6%
TERMINAL (d)	5/38 (13%)	6/31 (19%)	6/34 (18%)	8/32 (25%)
FIRST INCIDENCE	697	618	654	729 (T)
HC TUMORS SAME RTE	15/100 (15%)			
HC TUMORS ALL RTES	128/999 (13%)			
STATISTICAL TESTS				
POLY 3	P=0.290	P=0.558	P=0.332	P=0.381
POLY 1.5	P=0.318	P=0.580	P=0.336	P=0.413
POLY 6	P=0.275	P=0.532	P=0.337	P=0.358
COCH-ARM / FISHERS	P=0.409	P=0.613N	P=0.393	P=0.500
MAX-ISO-POLY-3	P=0.443	P=0.444	P=0.247	P=0.293
HISTCONT SAME RTE	P=0.271	P=0.594	P=0.193	P=0.231
HISTCONT ALL RTES	P=0.221	P=0.521	P=0.275	P=0.339

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Intestine Small: Jejunum Carcinoma				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	0/43.48	0/43.18	0/40.8
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	4/999 (0%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Intestine Small: Site Unspecified				
Carcinoma				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	0/43.48	0/43.18	0/40.8
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	5/999 (1%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Intestine Small: Site Unspecified **Carcinoma or Adenoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	0/43.48	0/43.18	0/40.8
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	8/999 (1%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Islets, Pancreatic **Carcinoma or Adenoma**				
TUMOR RATES				
OVERALL (a)	1/50 (2%)	0/50 (0%)	1/50 (2%)	0/50 (0%)
POLY-3 RATE (b)	1/46.91	0/43.48	1/43.18	0/40.8
POLY-3 PERCENT (g)	2.1%	0%	2.3%	0%
TERMINAL (d)	1/38 (3%)	0/31 (0%)	1/34 (3%)	0/32 (0%)
FIRST INCIDENCE	729 (T)	---	729 (T)	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	12/984 (1%)			
STATISTICAL TESTS				
POLY 3	P=0.419N	P=0.515N	P=0.742	P=0.528N
POLY 1.5	P=0.417N	P=0.509N	P=0.744	P=0.520N
POLY 6	P=0.418N	P=0.522N	P=0.742	P=0.533N
COCH-ARM / FISHERS	P=0.405N	P=0.500N	P=0.753N	P=0.500N
MAX-ISO-POLY-3	P=0.309N	P=0.177N	P=0.476	P=0.192N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Kidney: Renal Tubule Adenoma				
TUMOR RATES				
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	0/43.48	0/43.18	0/40.8
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	0/990 (0%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Kidney: Renal Tubule
Carcinoma or Adenoma

TUMOR RATES

OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	0/43.48	0/43.18	0/40.8
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	0/990 (0%)			

STATISTICAL TESTS

POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Liver **Hemangiosarcoma**				
TUMOR RATES				
OVERALL (a)	0/50 (0%)	1/50 (2%)	2/50 (4%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	1/43.48	2/43.18	0/40.8
POLY-3 PERCENT (g)	0%	2.3%	4.6%	0%
TERMINAL (d)	0/38 (0%)	1/31 (3%)	2/34 (6%)	0/32 (0%)
FIRST INCIDENCE	---	729 (T)	729 (T)	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	8/996 (1%)			
STATISTICAL TESTS				
POLY 3	P=0.620	P=0.485	P=0.219	(e)
POLY 1.5	P=0.639	P=0.491	P=0.221	(e)
POLY 6	P=0.607	P=0.478	P=0.219	(e)
COCH-ARM / FISHERS	P=0.616N	P=0.500	P=0.247	(e)
MAX-ISO-POLY-3	P=0.343	P=0.158	P=0.076	(e)
HISTCONT SAME RTE	P=0.441	P=0.121	P=0.106	(e)
HISTCONT ALL RTES	P=0.400	P=0.189	P=0.048*	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Liver **Hepatocellular Adenoma**				
TUMOR RATES				
OVERALL (a)	21/50 (42%)	20/50 (40%)	17/50 (34%)	15/50 (30%)
POLY-3 RATE (b)	21/47.66	20/43.9	17/43.23	15/41.75
POLY-3 PERCENT (g)	44.1%	45.6%	39.3%	35.9%
TERMINAL (d)	18/38 (47%)	17/31 (55%)	16/34 (47%)	12/32 (38%)
FIRST INCIDENCE	578	648	715	555
HC TUMORS SAME RTE	20/100 (20%)			
HC TUMORS ALL RTES	270/996 (27%)			
STATISTICAL TESTS				
POLY 3	P=0.204N	P=0.527	P=0.403N	P=0.285N
POLY 1.5	P=0.178N	P=0.581	P=0.380N	P=0.245N
POLY 6	P=0.214N	P=0.460	P=0.417N	P=0.310N
COCH-ARM / FISHERS	P=0.103N	P=0.500N	P=0.268N	P=0.149N
MAX-ISO-POLY-3	P=0.330N	P=0.443	P=0.331N	P=0.231N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Liver **Hepatocellular Carcinoma**				
TUMOR RATES				
OVERALL (a)	2/50 (4%)	5/50 (10%)	4/50 (8%)	1/50 (2%)
POLY-3 RATE (b)	2/46.91	5/43.95	4/43.31	1/41
POLY-3 PERCENT (g)	4.3%	11.4%	9.2%	2.4%
TERMINAL (d)	2/38 (5%)	2/31 (7%)	2/34 (6%)	0/32 (0%)
FIRST INCIDENCE	729 (T)	633	711	677
HC TUMORS SAME RTE	6/100 (6%)			
HC TUMORS ALL RTES	101/996 (10%)			
STATISTICAL TESTS				
POLY 3	P=0.352N	P=0.190	P=0.301	P=0.547N
POLY 1.5	P=0.332N	P=0.200	P=0.305	P=0.535N
POLY 6	P=0.363N	P=0.180	P=0.302	P=0.554N
COCH-ARM / FISHERS	P=0.273N	P=0.218	P=0.339	P=0.500N
MAX-ISO-POLY-3	P=0.256N	P=0.109	P=0.183	P=0.331N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Liver **Hepatocellular Carcinoma or Hepatoblastoma**				
TUMOR RATES				
OVERALL (a)	2/50 (4%)	5/50 (10%)	4/50 (8%)	1/50 (2%)
POLY-3 RATE (b)	2/46.91	5/43.95	4/43.31	1/41
POLY-3 PERCENT (g)	4.3%	11.4%	9.2%	2.4%
TERMINAL (d)	2/38 (5%)	2/31 (7%)	2/34 (6%)	0/32 (0%)
FIRST INCIDENCE	729 (T)	633	711	677
HC TUMORS SAME RTE	7/100 (7%)			
HC TUMORS ALL RTES	104/996 (10%)			
STATISTICAL TESTS				
POLY 3	P=0.352N	P=0.190	P=0.301	P=0.547N
POLY 1.5	P=0.332N	P=0.200	P=0.305	P=0.535N
POLY 6	P=0.363N	P=0.180	P=0.302	P=0.554N
COCH-ARM / FISHERS	P=0.273N	P=0.218	P=0.339	P=0.500N
MAX-ISO-POLY-3	P=0.256N	P=0.109	P=0.183	P=0.331N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Liver
Hepatocellular Carcinoma or Hepatocellular Adenoma

TUMOR RATES

OVERALL (a)	23/50 (46%)	21/50 (42%)	19/50 (38%)	16/50 (32%)
POLY-3 RATE (b)	23/47.66	21/44.24	19/43.31	16/41.95
POLY-3 PERCENT (g)	48.3%	47.5%	43.9%	38.1%
TERMINAL (d)	20/38 (53%)	17/31 (55%)	17/34 (50%)	12/32 (38%)
FIRST INCIDENCE	578	633	711	555
HC TUMORS SAME RTE	23/100 (23%)			
HC TUMORS ALL RTES	329/996 (33%)			

STATISTICAL TESTS

POLY 3	P=0.172N	P=0.553N	P=0.417N	P=0.225N
POLY 1.5	P=0.149N	P=0.500N	P=0.393N	P=0.191N
POLY 6	P=0.177N	P=0.556	P=0.429N	P=0.243N
COCH-ARM / FISHERS	P=0.082N	P=0.420N	P=0.272N	P=0.109N
MAX-ISO-POLY-3	P=0.277N	P=0.469N	P=0.343N	P=0.181N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Liver
Hepatocellular Carcinoma, Hepatocellular Adenoma, or Hepatoblastoma

TUMOR RATES

OVERALL (a)	23/50 (46%)	21/50 (42%)	19/50 (38%)	16/50 (32%)
POLY-3 RATE (b)	23/47.66	21/44.24	19/43.31	16/41.95
POLY-3 PERCENT (g)	48.3%	47.5%	43.9%	38.1%
TERMINAL (d)	20/38 (53%)	17/31 (55%)	17/34 (50%)	12/32 (38%)
FIRST INCIDENCE	578	633	711	555
HC TUMORS SAME RTE	24/100 (24%)			
HC TUMORS ALL RTES	330/996 (33%)			

STATISTICAL TESTS

POLY 3	P=0.172N	P=0.553N	P=0.417N	P=0.225N
POLY 1.5	P=0.149N	P=0.500N	P=0.393N	P=0.191N
POLY 6	P=0.177N	P=0.556	P=0.429N	P=0.243N
COCH-ARM / FISHERS	P=0.082N	P=0.420N	P=0.272N	P=0.109N
MAX-ISO-POLY-3	P=0.277N	P=0.469N	P=0.343N	P=0.181N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Lung **Alveolar/Bronchiolar Adenoma**				
TUMOR RATES				
OVERALL (a)	6/50 (12%)	3/50 (6%)	2/50 (4%)	4/50 (8%)
POLY-3 RATE (b)	6/46.91	3/43.48	2/43.18	4/40.8
POLY-3 PERCENT (g)	12.8%	6.9%	4.6%	9.8%
TERMINAL (d)	6/38 (16%)	3/31 (10%)	2/34 (6%)	4/32 (13%)
FIRST INCIDENCE	729 (T)	729 (T)	729 (T)	729 (T)
HC TUMORS SAME RTE	6/100 (6%)			
HC TUMORS ALL RTES	47/998 (5%)			
STATISTICAL TESTS				
POLY 3	P=0.384N	P=0.281N	P=0.161N	P=0.460N
POLY 1.5	P=0.376N	P=0.265N	P=0.158N	P=0.435N
POLY 6	P=0.385N	P=0.300N	P=0.161N	P=0.476N
COCH-ARM / FISHERS	P=0.341N	P=0.243N	P=0.134N	P=0.370N
MAX-ISO-POLY-3	P=0.300N	P=0.185N	P=0.096N	P=0.341N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Lung
Alveolar/Bronchiolar Carcinoma

TUMOR RATES

OVERALL (a)	1/50 (2%)	3/50 (6%)	1/50 (2%)	0/50 (0%)
POLY-3 RATE (b)	1/46.91	3/44.05	1/43.19	0/40.8
POLY-3 PERCENT (g)	2.1%	6.8%	2.3%	0%
TERMINAL (d)	1/38 (3%)	2/31 (7%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	729 (T)	550	725	---
HC TUMORS SAME RTE	3/100 (3%)			
HC TUMORS ALL RTES	43/998 (4%)			

STATISTICAL TESTS

POLY 3	P=0.242N	P=0.283	P=0.742	P=0.528N
POLY 1.5	P=0.231N	P=0.292	P=0.744	P=0.520N
POLY 6	P=0.249N	P=0.273	P=0.742	P=0.533N
COCH-ARM / FISHERS	P=0.200N	P=0.309	P=0.753N	P=0.500N
MAX-ISO-POLY-3	P=0.200N	P=0.146	P=0.476	P=0.192N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Lung
Alveolar/Bronchiolar Carcinoma or Alveolar/Bronchiolar Adenoma

TUMOR RATES

OVERALL (a)	7/50 (14%)	6/50 (12%)	3/50 (6%)	4/50 (8%)
POLY-3 RATE (b)	7/46.91	6/44.05	3/43.19	4/40.8
POLY-3 PERCENT (g)	14.9%	13.6%	7%	9.8%
TERMINAL (d)	7/38 (18%)	5/31 (16%)	2/34 (6%)	4/32 (13%)
FIRST INCIDENCE	729 (T)	550	725	729 (T)
HC TUMORS SAME RTE	8/100 (8%)			
HC TUMORS ALL RTES	86/998 (9%)			

STATISTICAL TESTS

POLY 3	P=0.221N	P=0.548N	P=0.193N	P=0.346N
POLY 1.5	P=0.209N	P=0.529N	P=0.189N	P=0.323N
POLY 6	P=0.226N	P=0.571N	P=0.192N	P=0.362N
COCH-ARM / FISHERS	P=0.169N	P=0.500N	P=0.159N	P=0.262N
MAX-ISO-POLY-3	P=0.296N	P=0.430N	P=0.124N	P=0.252N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Ovary **Cystadenoma**				
TUMOR RATES				
OVERALL (a)	7/50 (14%)	2/50 (4%)	3/50 (6%)	0/50 (0%)
POLY-3 RATE (b)	7/46.91	2/43.48	3/43.18	0/40.8
POLY-3 PERCENT (g)	14.9%	4.6%	7%	0%
TERMINAL (d)	7/38 (18%)	2/31 (7%)	3/34 (9%)	0/32 (0%)
FIRST INCIDENCE	729 (T)	729 (T)	729 (T)	---
HC TUMORS SAME RTE	4/100 (4%)			
HC TUMORS ALL RTES	39/977 (4%)			
STATISTICAL TESTS				
POLY 3	P=0.010N**	P=0.098N	P=0.193N	P=0.013N*
POLY 1.5	P=0.009N**	P=0.090N	P=0.189N	P=0.011N*
POLY 6	P=0.010N**	P=0.108N	P=0.192N	P=0.014N*
COCH-ARM / FISHERS	P=0.008N**	P=0.080N	P=0.159N	P=0.006N**
MAX-ISO-POLY-3	P=0.007N**	P=0.056N	P=0.124N	P=0.007N**
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Pituitary Gland: Pars Distalis or Unspecified Site
Adenoma

TUMOR RATES

OVERALL (a)	1/50 (2%)	4/50 (8%)	5/50 (10%)	3/49 (6%)
POLY-3 RATE (b)	1/46.91	4/43.62	5/44.23	3/39.84
POLY-3 PERCENT (g)	2.1%	9.2%	11.3%	7.5%
TERMINAL (d)	1/38 (3%)	2/31 (7%)	3/34 (9%)	2/31 (7%)
FIRST INCIDENCE	729 (T)	704	444	719
HC TUMORS SAME RTE	5/99 (5%)			
HC TUMORS ALL RTES	72/974 (7%)			

STATISTICAL TESTS

POLY 3	P=0.232	P=0.157	P=0.088	P=0.250
POLY 1.5	P=0.253	P=0.166	P=0.089	P=0.262
POLY 6	P=0.219	P=0.148	P=0.090	P=0.241
COCH-ARM / FISHERS	P=0.320	P=0.181	P=0.102	P=0.301
MAX-ISO-POLY-3	P=0.180	P=0.078	P=0.042*	P=0.134
HISTCONT SAME RTE	P=0.243	P=0.109	P=0.057	P=0.190
HISTCONT ALL RTES	P=0.209	P=0.089	P=0.045*	P=0.160

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Skin
Basal or Sq. Cell Carcinoma, Carcinoma, Basosq. Tumor (M or B), Basal Cell Adenoma, Adenoma, Papilloma, Sq Papilloma, Keratoacanthoma, Trichoepithelioma

TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	1/50 (2%)
POLY-3 RATE (b)	0/46.91	0/43.48	0/43.18	1/40.8
POLY-3 PERCENT (g)	0%	0%	0%	2.5%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	1/32 (3%)
FIRST INCIDENCE	---	---	---	729 (T)
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	5/999 (1%)			

STATISTICAL TESTS				
POLY 3	P=0.184	(e)	(e)	P=0.472
POLY 1.5	P=0.188	(e)	(e)	P=0.480
POLY 6	P=0.183	(e)	(e)	P=0.467
COCH-ARM / FISHERS	P=0.198	(e)	(e)	P=0.500
MAX-ISO-POLY-3	P=0.123	(e)	(e)	P=0.158
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Skin
Fibroma, Fibrosarcoma, Sarcoma, Myxoma, Myxosarcoma, or Fibrous Histiocytoma

TUMOR RATES	#	#	#	#
OVERALL (a)	4/50 (8%)	3/50 (6%)	1/50 (2%)	5/50 (10%)
POLY-3 RATE (b)	4/47.66	3/43.78	1/43.39	5/41.47
POLY-3 PERCENT (g)	8.4%	6.9%	2.3%	12.1%
TERMINAL (d)	1/38 (3%)	2/31 (7%)	0/34 (0%)	2/32 (6%)
FIRST INCIDENCE	590	648	673	644
HC TUMORS SAME RTE	1/100 (1%)			
HC TUMORS ALL RTES	46/999 (5%)			

STATISTICAL TESTS				
POLY 3	P=0.356	P=0.546N	P=0.209N	P=0.414
POLY 1.5	P=0.366	P=0.528N	P=0.204N	P=0.436
POLY 6	P=0.353	P=0.568N	P=0.211N	P=0.400
COCH-ARM / FISHERS	P=0.404	P=0.500N	P=0.181N	P=0.500
MAX-ISO-POLY-3	P=0.242	P=0.394N	P=0.112N	P=0.298
HISTCONT SAME RTE	P=0.114	P=0.248	P=0.569	P=0.131
HISTCONT ALL RTES	P=0.183	P=0.455	P=1.000	P=0.124

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Skin **Fibrosarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	3/50 (6%)	0/50 (0%)	0/50 (0%)	2/50 (4%)
POLY-3 RATE (b)	3/47.45	0/43.48	0/43.18	2/40.8
POLY-3 PERCENT (g)	6.3%	0%	0%	4.9%
TERMINAL (d)	1/38 (3%)	0/31 (0%)	0/34 (0%)	2/32 (6%)
FIRST INCIDENCE	590	---	---	729 (T)
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	18/999 (2%)			
STATISTICAL TESTS				
POLY 3	P=0.524N	P=0.135N	P=0.137N	P=0.568N
POLY 1.5	P=0.529N	P=0.128N	P=0.134N	P=0.549N
POLY 6	P=0.516N	P=0.144N	P=0.138N	P=0.582N
COCH-ARM / FISHERS	P=0.531N	P=0.121N	P=0.121N	P=0.500N
MAX-ISO-POLY-3	P=0.168N	P=0.052N	P=0.053N	P=0.393N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

Skin
Fibrosarcoma, Sarcoma, Myxosarcoma, or Fibrous Histiocytoma

TUMOR RATES	#	#	#	#
OVERALL (a)	4/50 (8%)	3/50 (6%)	1/50 (2%)	5/50 (10%)
POLY-3 RATE (b)	4/47.66	3/43.78	1/43.39	5/41.47
POLY-3 PERCENT (g)	8.4%	6.9%	2.3%	12.1%
TERMINAL (d)	1/38 (3%)	2/31 (7%)	0/34 (0%)	2/32 (6%)
FIRST INCIDENCE	590	648	673	644
HC TUMORS SAME RTE	1/100 (1%)			
HC TUMORS ALL RTES	46/999 (5%)			

STATISTICAL TESTS				
POLY 3	P=0.356	P=0.546N	P=0.209N	P=0.414
POLY 1.5	P=0.366	P=0.528N	P=0.204N	P=0.436
POLY 6	P=0.353	P=0.568N	P=0.211N	P=0.400
COCH-ARM / FISHERS	P=0.404	P=0.500N	P=0.181N	P=0.500
MAX-ISO-POLY-3	P=0.242	P=0.394N	P=0.112N	P=0.298
HISTCONT SAME RTE	P=0.114	P=0.248	P=0.569	P=0.131
HISTCONT ALL RTES	P=0.183	P=0.455	P=1.000	P=0.124

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Skin **Fibrous Histiocytoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	1/50 (2%)	0/50 (0%)	0/50 (0%)	3/50 (6%)
POLY-3 RATE (b)	1/47.12	0/43.48	0/43.18	3/41.47
POLY-3 PERCENT (g)	2.1%	0%	0%	7.2%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	673	---	---	644
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	7/999 (1%)			
STATISTICAL TESTS				
POLY 3	P=0.080	P=0.516N	P=0.517N	P=0.261
POLY 1.5	P=0.080	P=0.510N	P=0.515N	P=0.272
POLY 6	P=0.081	P=0.524N	P=0.518N	P=0.256
COCH-ARM / FISHERS	P=0.086	P=0.500N	P=0.500N	P=0.309
MAX-ISO-POLY-3	P=0.040*	P=0.178N	P=0.180N	P=0.139
HISTCONT SAME RTE	P=0.054	(e)	(e)	P=0.100
HISTCONT ALL RTES	P=0.032*	(e)	(e)	P=0.025*

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Skin **Sarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	3/50 (6%)	1/50 (2%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	3/43.78	1/43.39	0/40.8
POLY-3 PERCENT (g)	0%	6.9%	2.3%	0%
TERMINAL (d)	0/38 (0%)	2/31 (7%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	648	673	---
HC TUMORS SAME RTE	1/100 (1%)			
HC TUMORS ALL RTES	22/999 (2%)			
STATISTICAL TESTS				
POLY 3	P=0.427N	P=0.107	P=0.484	(e)
POLY 1.5	P=0.411N	P=0.111	P=0.486	(e)
POLY 6	P=0.438N	P=0.102	P=0.485	(e)
COCH-ARM / FISHERS	P=0.366N	P=0.121	P=0.500	(e)
MAX-ISO-POLY-3	P=0.267N	P=0.038*	P=0.158	(e)
HISTCONT SAME RTE	(h)	(h)	(h)	(e)
HISTCONT ALL RTES	(h)	(h)	(h)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Skin **Schwannoma Malignant**				
TUMOR RATES	#	#	#	#
OVERALL (a)	2/50 (4%)	0/50 (0%)	1/50 (2%)	2/50 (4%)
POLY-3 RATE (b)	2/47.56	0/43.48	1/43.18	2/41.08
POLY-3 PERCENT (g)	4.2%	0%	2.3%	4.9%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	1/34 (3%)	0/32 (0%)
FIRST INCIDENCE	596	---	729 (T)	677
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	3/999 (0%)			
STATISTICAL TESTS				
POLY 3	P=0.443	P=0.258N	P=0.534N	P=0.640
POLY 1.5	P=0.448	P=0.249N	P=0.528N	P=0.655
POLY 6	P=0.443	P=0.269N	P=0.537N	P=0.629
COCH-ARM / FISHERS	P=0.469	P=0.247N	P=0.500N	P=0.691N
MAX-ISO-POLY-3	P=0.390	P=0.095N	P=0.317N	P=0.443
HISTCONT SAME RTE	P=0.148	(e)	P=0.347	P=0.169
HISTCONT ALL RTES	P=0.007**	(e)	P=0.157	P=0.022*

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Skin **Squamous Cell Papilloma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	0/43.48	0/43.18	0/40.8
POLY-3 PERCENT (g)	0%	0%	0%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	---	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	1/999 (0%)			
STATISTICAL TESTS				
POLY 3	(e)	(e)	(e)	(e)
POLY 1.5	(e)	(e)	(e)	(e)
POLY 6	(e)	(e)	(e)	(e)
COCH-ARM / FISHERS	(e)	(e)	(e)	(e)
MAX-ISO-POLY-3	(e)	(e)	(e)	(e)
HISTCONT SAME RTE	(e)	(e)	(e)	(e)
HISTCONT ALL RTES	(e)	(e)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Uterus **Polyp Stromal**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	2/50 (4%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	0/43.48	2/43.18	0/40.8
POLY-3 PERCENT (g)	0%	0%	4.6%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	2/34 (6%)	0/32 (0%)
FIRST INCIDENCE	---	---	729 (T)	---
HC TUMORS SAME RTE	1/100 (1%)			
HC TUMORS ALL RTES	29/999 (3%)			
STATISTICAL TESTS				
POLY 3	P=0.530	(e)	P=0.219	(e)
POLY 1.5	P=0.548	(e)	P=0.221	(e)
POLY 6	P=0.519	(e)	P=0.219	(e)
COCH-ARM / FISHERS	P=0.595	(e)	P=0.247	(e)
MAX-ISO-POLY-3	P=0.262	(e)	P=0.076	(e)
HISTCONT SAME RTE	P=0.347	(e)	P=0.106	(e)
HISTCONT ALL RTES	P=0.304	(e)	P=0.086	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
Uterus				
Sarcoma Stromal or Polyp Stromal				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	0/50 (0%)	2/50 (4%)	1/50 (2%)
POLY-3 RATE (b)	0/46.91	0/43.48	2/43.18	1/41.62
POLY-3 PERCENT (g)	0%	0%	4.6%	2.4%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	2/34 (6%)	0/32 (0%)
FIRST INCIDENCE	---	---	729 (T)	410
HC TUMORS SAME RTE	1/100 (1%)			
HC TUMORS ALL RTES	32/999 (3%)			
STATISTICAL TESTS				
POLY 3	P=0.210	(e)	P=0.219	P=0.476
POLY 1.5	P=0.219	(e)	P=0.221	P=0.482
POLY 6	P=0.206	(e)	P=0.219	P=0.472
COCH-ARM / FISHERS	P=0.246	(e)	P=0.247	P=0.500
MAX-ISO-POLY-3	P=0.179	(e)	P=0.076	P=0.158
HISTCONT SAME RTE	P=0.241	(e)	P=0.106	P=0.223
HISTCONT ALL RTES	P=0.207	(e)	P=0.086	P=0.189

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Hemangioma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	1/50 (2%)	1/50 (2%)	2/50 (4%)	1/50 (2%)
POLY-3 RATE (b)	1/46.91	1/43.5	2/43.18	1/40.8
POLY-3 PERCENT (g)	2.1%	2.3%	4.6%	2.5%
TERMINAL (d)	1/38 (3%)	0/31 (0%)	2/34 (6%)	1/32 (3%)
FIRST INCIDENCE	729 (T)	725	729 (T)	729 (T)
HC TUMORS SAME RTE	2/100 (2%)			
HC TUMORS ALL RTES	15/999 (2%)			
STATISTICAL TESTS				
POLY 3	P=0.532	P=0.744	P=0.471	P=0.729
POLY 1.5	P=0.547	P=0.750	P=0.475	P=0.737
POLY 6	P=0.524	P=0.735	P=0.471	P=0.723
COCH-ARM / FISHERS	P=0.591	P=0.753N	P=0.500	P=0.753N
MAX-ISO-POLY-3	P=0.559	P=0.478	P=0.265	P=0.462
HISTCONT SAME RTE	P=0.607	P=0.632	P=0.315	P=0.614
HISTCONT ALL RTES	P=0.457	P=0.526	P=0.204	P=0.509

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Hemangiosarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	2/50 (4%)	3/50 (6%)	1/50 (2%)
POLY-3 RATE (b)	0/46.91	2/44.28	3/43.18	1/40.8
POLY-3 PERCENT (g)	0%	4.5%	7%	2.5%
TERMINAL (d)	0/38 (0%)	1/31 (3%)	3/34 (9%)	1/32 (3%)
FIRST INCIDENCE	---	428	729 (T)	729 (T)
HC TUMORS SAME RTE	7/100 (7%)			
HC TUMORS ALL RTES	43/999 (4%)			
STATISTICAL TESTS				
POLY 3	P=0.360	P=0.225	P=0.105	P=0.472
POLY 1.5	P=0.382	P=0.230	P=0.106	P=0.480
POLY 6	P=0.347	P=0.218	P=0.104	P=0.467
COCH-ARM / FISHERS	P=0.444	P=0.247	P=0.121	P=0.500
MAX-ISO-POLY-3	P=0.203	P=0.076	P=0.038*	P=0.158
HISTCONT SAME RTE	P=0.272	P=0.106	P=0.051	P=0.223
HISTCONT ALL RTES	P=0.235	P=0.086	P=0.040*	P=0.189

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Hemangiosarcoma or Hemangioma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	1/50 (2%)	3/50 (6%)	5/50 (10%)	2/50 (4%)
POLY-3 RATE (b)	1/46.91	3/44.3	5/43.18	2/40.8
POLY-3 PERCENT (g)	2.1%	6.8%	11.6%	4.9%
TERMINAL (d)	1/38 (3%)	1/31 (3%)	5/34 (15%)	2/32 (6%)
FIRST INCIDENCE	729 (T)	428	729 (T)	729 (T)
HC TUMORS SAME RTE	9/100 (9%)			
HC TUMORS ALL RTES	58/999 (6%)			
STATISTICAL TESTS				
POLY 3	P=0.340	P=0.285	P=0.083	P=0.452
POLY 1.5	P=0.364	P=0.294	P=0.086	P=0.464
POLY 6	P=0.325	P=0.274	P=0.083	P=0.443
COCH-ARM / FISHERS	P=0.438	P=0.309	P=0.102	P=0.500
MAX-ISO-POLY-3	P=0.213	P=0.147	P=0.041*	P=0.255
HISTCONT SAME RTE	P=0.284	P=0.208	P=0.056	P=0.357
HISTCONT ALL RTES	P=0.246	P=0.175	P=0.044*	P=0.317

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Females		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Histiocytic Sarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	1/50 (2%)	1/50 (2%)	2/50 (4%)
POLY-3 RATE (b)	0/46.91	1/43.71	1/43.26	2/41.52
POLY-3 PERCENT (g)	0%	2.3%	2.3%	4.8%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	1/32 (3%)
FIRST INCIDENCE	---	668	709	477
HC TUMORS SAME RTE	2/100 (2%)			
HC TUMORS ALL RTES	25/999 (3%)			
STATISTICAL TESTS				
POLY 3	P=0.129	P=0.486	P=0.484	P=0.211
POLY 1.5	P=0.134	P=0.491	P=0.486	P=0.217
POLY 6	P=0.126	P=0.480	P=0.484	P=0.207
COCH-ARM / FISHERS	P=0.153	P=0.500	P=0.500	P=0.247
MAX-ISO-POLY-3	P=0.121	P=0.158	P=0.158	P=0.076
HISTCONT SAME RTE	P=0.163	P=0.223	P=0.223	P=0.106
HISTCONT ALL RTES	P=0.139	P=0.189	P=0.189	P=0.086

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**

All Organs
Malignant Lymphoma: Histiocytic, Lymphocytic, Mixed, NOS, or Undifferentiated Cell Type

TUMOR RATES	#	#	#	#
OVERALL (a)	9/50 (18%)	12/50 (24%)	6/50 (12%)	8/50 (16%)
POLY-3 RATE (b)	9/47.57	12/45.12	6/43.18	8/42.06
POLY-3 PERCENT (g)	18.9%	26.6%	13.9%	19%
TERMINAL (d)	4/38 (11%)	8/31 (26%)	6/34 (18%)	4/32 (13%)
FIRST INCIDENCE	677	508	729 (T)	493
HC TUMORS SAME RTE	18/100 (18%)			
HC TUMORS ALL RTES	213/999 (21%)			

STATISTICAL TESTS				
POLY 3	P=0.401N	P=0.263	P=0.360N	P=0.601
POLY 1.5	P=0.377N	P=0.283	P=0.348N	P=0.583N
POLY 6	P=0.415N	P=0.238	P=0.370N	P=0.584
COCH-ARM / FISHERS	P=0.296N	P=0.312	P=0.288N	P=0.500N
MAX-ISO-POLY-3	P=0.391N	P=0.195	P=0.271N	P=0.495
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Osteosarcoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	2/50 (4%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	2/44.91	0/43.18	0/40.8
POLY-3 PERCENT (g)	0%	4.5%	0%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	449	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	4/999 (0%)			
STATISTICAL TESTS				
POLY 3	P=0.439N	P=0.228	(e)	(e)
POLY 1.5	P=0.430N	P=0.232	(e)	(e)
POLY 6	P=0.445N	P=0.222	(e)	(e)
COCH-ARM / FISHERS	P=0.405N	P=0.247	(e)	(e)
MAX-ISO-POLY-3	P=0.297N	P=0.076	(e)	(e)
HISTCONT SAME RTE	(h)	(h)	(e)	(e)
HISTCONT ALL RTES	(h)	(h)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Osteosarcoma or Osteoma**				
TUMOR RATES	#	#	#	#
OVERALL (a)	0/50 (0%)	2/50 (4%)	0/50 (0%)	0/50 (0%)
POLY-3 RATE (b)	0/46.91	2/44.91	0/43.18	0/40.8
POLY-3 PERCENT (g)	0%	4.5%	0%	0%
TERMINAL (d)	0/38 (0%)	0/31 (0%)	0/34 (0%)	0/32 (0%)
FIRST INCIDENCE	---	449	---	---
HC TUMORS SAME RTE	0/100 (0%)			
HC TUMORS ALL RTES	5/999 (1%)			
STATISTICAL TESTS				
POLY 3	P=0.439N	P=0.228	(e)	(e)
POLY 1.5	P=0.430N	P=0.232	(e)	(e)
POLY 6	P=0.445N	P=0.222	(e)	(e)
COCH-ARM / FISHERS	P=0.405N	P=0.247	(e)	(e)
MAX-ISO-POLY-3	P=0.297N	P=0.076	(e)	(e)
HISTCONT SAME RTE	(h)	(h)	(e)	(e)
HISTCONT ALL RTES	(h)	(h)	(e)	(e)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Benign Tumors**				
TUMOR RATES	#	#	#	#
OVERALL (a)	33/50 (66%)	30/50 (60%)	29/50 (58%)	25/50 (50%)
POLY-3 RATE (b)	33/47.83	30/44.99	29/44.58	25/41.79
POLY-3 PERCENT (g)	69%	66.7%	65.1%	59.8%
TERMINAL (d)	28/38 (74%)	23/31 (74%)	24/34 (71%)	21/32 (66%)
FIRST INCIDENCE	578	550	444	555
HC TUMORS SAME RTE	43/100 (43%)			
HC TUMORS ALL RTES	508/999 (51%)			
STATISTICAL TESTS				
POLY 3	P=0.198N	P=0.493N	P=0.427N	P=0.241N
POLY 1.5	P=0.157N	P=0.429N	P=0.414N	P=0.184N
POLY 6	P=0.217N	P=0.570N	P=0.417N	P=0.283N
COCH-ARM / FISHERS	P=0.063N	P=0.339N	P=0.268N	P=0.078N
MAX-ISO-POLY-3	P=0.308N	P=0.403N	P=0.345N	P=0.191N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

		Females		
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Malignant Tumors**				
TUMOR RATES	#	#	#	#
OVERALL (a)	15/50 (30%)	27/50 (54%)	17/50 (34%)	20/50 (40%)
POLY-3 RATE (b)	15/48.77	27/48.24	17/44.5	20/44.55
POLY-3 PERCENT (g)	30.8%	56%	38.2%	44.9%
TERMINAL (d)	6/38 (16%)	15/31 (48%)	10/34 (29%)	9/32 (28%)
FIRST INCIDENCE	590	428	521	410
HC TUMORS SAME RTE	42/100 (42%)			
HC TUMORS ALL RTES	495/999 (50%)			
STATISTICAL TESTS				
POLY 3	P=0.258	P=0.009**	P=0.296	P=0.115
POLY 1.5	P=0.290	P=0.010**	P=0.314	P=0.131
POLY 6	P=0.247	P=0.007**	P=0.285	P=0.109
COCH-ARM / FISHERS	P=0.432	P=0.013*	P=0.415	P=0.201
MAX-ISO-POLY-3	P=0.103	P=0.005**	P=0.236	P=0.088
HISTCONT SAME RTE	P=0.141	P=0.006**	P=0.331	P=0.123
HISTCONT ALL RTES	P=0.119	P=0.005**	P=0.291	P=0.101

STATISTICAL ANALYSIS OF PRIMARY TUMORS IN MICE(B6C3F1)
TERMINAL SACRIFICE AT 105 WEEKS

	Females			
DOSE	**0 MG/KG**	**1250 MG/KG**	**2500 MG/KG**	**5000 MG/KG**
All Organs **Malignant and Benign Tumors**				
TUMOR RATES	#	#	#	#
OVERALL (a)	42/50 (84%)	38/50 (76%)	35/50 (70%)	36/50 (72%)
POLY-3 RATE (b)	42/49.4	38/48.73	35/45.56	36/45.11
POLY-3 PERCENT (g)	85%	78%	76.8%	79.8%
TERMINAL (d)	31/38 (82%)	24/31 (77%)	26/34 (77%)	24/32 (75%)
FIRST INCIDENCE	578	428	444	410
HC TUMORS SAME RTE	70/100 (70%)			
HC TUMORS ALL RTES	746/999 (75%)			
STATISTICAL TESTS				
POLY 3	P=0.336N	P=0.257N	P=0.222N	P=0.344N
POLY 1.5	P=0.276N	P=0.240N	P=0.194N	P=0.276N
POLY 6	P=0.356N	P=0.298N	P=0.242N	P=0.380N
COCH-ARM / FISHERS	P=0.106N	P=0.227N	P=0.077N	P=0.114N
MAX-ISO-POLY-3	P=0.339N	P=0.182N	P=0.161N	P=0.259N
HISTCONT SAME RTE	(h)	(h)	(h)	(h)
HISTCONT ALL RTES	(h)	(h)	(h)	(h)

LEGEND

(a)	Number of tumor-bearing animals/number of animals examined at site.
(b)	Number of tumor-bearing animals/Poly-3 number
(d)	Observed incidence at terminal kill.
(e)	Value of Statistic cannot be computed.
(f)	Beneath the control incidence are the P-values associated with the trend test. Beneath the dosed group incidence are theP-values corresponding to pairwise comparisons between the controls and that dosed group.
(g)	Poly-3 adjusted lifetime tumor incidence.
(h)	Value of Historical Controls statistic is only calculated when Poly-3 trend is positive and when there is more than one tumor in the treatment groups.
(I)	Interim sacrifice
(T)	Terminal sacrifice
#	Tumor rates based on numbers of animals necropsied.
*	To the right of any statistical result, indicates significance at (P<=0.05).
**	To the right of any statistical result, indicates significance at (P<=0.01).
N	Indicates a negative trend for all tests
	The Cochran-Armitage and Fishers exact tests compare directly the overall incidence rates.

*** END OF REPORT ***