




Historical Control Body Weights for Gavage/Corn Oil Studies Male Rats/Harlan Sprague–Dawley 1 study




Female Rats/Harlan Sprague–Dawley 1 study



Historical Control Body Weights for Inhalation/Air Studies Male Rats/Harlan Sprague–Dawley 1 study




Female Rats/Harlan Sprague–Dawley 1 study




Historical Control Body Weights for Oral/Feed Studies


Male Rats/Harlan Sprague–Dawley 1 study




Female Rats/Harlan Sprague–Dawley 1 study



Historical Control Body Weights for Perinatal Postnatal Gavage/Methylcellulose Studies Male Rats/Harlan Sprague–Dawley 2 studies




Female Rats/Harlan Sprague–Dawley 2 studies




Historical Control Body Weights for Perinatal Postnatal/Feed Studies


Male Rats/Harlan Sprague–Dawley 5 studies




Female Rats/Harlan Sprague–Dawley 5 studies




Historical Control Body Weights for Perinatal Postnatal/Water Studies Male Rats/Harlan Sprague–Dawley 1 study



Female Rats/Harlan Sprague–Dawley 1 study



Historical Control Body Weights for Whole Body Exposure/Air Studies Male Rats/Harlan Sprague–Dawley 1 study



Female Rats/Harlan Sprague–Dawley 1 study

